

# Quality Changes During Postharvest Handling Chain of Black Tiger Shrimp ARTIKEL SEMINAR faperta 2015-29 juni

*by* Andi Noor Asikin

---

**Submission date:** 20-Apr-2020 07:58AM (UTC+0700)

**Submission ID:** 1302010757

**File name:** n\_of\_Black\_Tiger\_Shrimp\_ARTIKEL\_SEMINAR\_faperta\_2015-29\_juni.pdf (404.61K)

**Word count:** 5132

**Character count:** 26354

# Quality Changes During Postharvest Handling Chain of Black Tiger Shrimp (*Penaeus Monodon* Fab. 1897) from Brackish Water Pond in Mahakam Delta Region: Case Study

Andi Noor Asikin<sup>1\*</sup>, Sahala Hutabarat<sup>2</sup>, YS.Darmanto<sup>2</sup>, Slamet Budi Prayitno<sup>2</sup>

<sup>1</sup>Faculty of Fisheries and Marine Sciences Mulawarman University, Jl. Gunung Tabur Kampus Gunung Kelua Samarinda 75119, Indonesia. Asikin63@yahoo.com

<sup>2</sup>Faculty of Fisheries and Marine Sciences Diponegoro University, Jl. Prof. Soedarto, SH. Kampus UNDIP Tembalang Semarang 50275, Indonesia.

## Abstract

Shrimps are more perishable food and their quality are depend on some of factors including harvest condition, handling process, storage duration and temperature during handling. Most of farmer and collector in Mahakam delta are not aware about the postharvest handling so that caused low quality of shrimp. The aim of this study were to investigate of biochemical quality and microbiological content on the postharvest handling chain of shrimp in Mahakam delta region. The study were carried out in three sub districts namely Muara Jawa, Anggana and Muara Badak. Three small ponds around 2-5 Ha in each sub districts were used as observation unit. The shrimp samples were taken at ponds, farmer point and depot. The biochemical parameters observed were moisture content, pH, TVB-N, indole and microbial parameter was Total Plate Count (TPC). The result showed that moisture content of shrimp was significant differences ( $P < 0.05$ ) at pond, farmer point and depot in Muara Jawa, Anggana and Muara Badak. The pH content found was not significant differences ( $P > 0.05$ ) at ponds, farmer point and depot in Muara Jawa, Anggana and Muara Badak sub districts. The TVB-N content of shrimp was significant differences ( $P < 0.05$ ) at all of handling stages in three sub districts. The indole content of shrimp was found only at depot level in three sub districts. The TPC content found was not significant level on all handling stages in three sub districts studied area. From the research found, it can be concluded that the quality of postharvest handling chain of black tiger shrimp in Mahakam delta were still meet the requirements, however they still need improvement so that the shrimp quality will increase more.

Key words: Change, Quality, postharvest, handling chain, pond, Mahakam Delta.

## I. Introduction

Mahakam Delta is located in the jurisdiction of Kutai Kartanegara district that is the largest contributor of exporting black tiger shrimp to East Kalimantan. Mahakam delta has been known as a centre of good shrimp culture in Indonesia, especially Muara Jawa, Anggana and Muara Badak sub districts. Due to its unique condition and it was very good mangrove forest that intensively converted into brackish water ponds and favourable environment for shrimp culture and tradisional shrimp farming. In 2011, the production of black tiger shrimp from Kutai Kartanegara district contributed as much as 13.259 tons (65%) for shrimp exports of East Kalimantan. These exports were consisted of 6156 tons of shrimp, white shrimp 2786 tons and 4316 tons of other shrimp (Marine and Fisheries Agency of Kutai Kartanegara, 2012). The main cultivated species in this region is *Penaeus monodon*, more commonly referred to as black tiger shrimp. Consumers today are critical and require the provision of safe and healthy food products. Exportable shrimp requires special care to retain as much as particable to original physical appearance, odor and organoleptic condition. It must be free from dirt, filth, pathogenic organism, uncertified chemicals and any antibiotic event in the minute's quality. The importing countries particularly the USA, UE and Japan are highly conscious about food hygiene and safety (Paul, S. *et al.*, 2010; Hassan, M.N. *et al.*, 2012).

Postharvest handling chain of shrimp practices in Mahakam delta region consist of three stages: harvesting by farmer at the pond, postharvest handling by farmer at the farmer point and postharvest handling by collectors at depot and then bring to processing plant. Long period of postharvest handling chain, use of inadequate of ice, bad water quality for handling process caused low quality of shrimp. In the region, most of the shrimp farmers and collectors are not aware on the importance of the postharvest handling. The shrimps which brought to processing plant sometimes do not meet the criteria of standard shrimp quality at processing plant. The shrimps sold to the processing plant were not always accepted even in some condition it will be rejected. The Shrimps were severely degraded during handling and transportation, around 20-30% was below minimum accepted processing plant quality (Asikin, A.N., 2012; Asikin, A.N. *et al.*, 2012). The longtime of postharvest handling chain to processing plant caused quality change of shrimps. In this postharvest handling chain, shrimps pass one stage to another stage with the elapse of time, time duration because the inner biochemical degradation of shrimp body in the different stage (Ali, M.Y., *et al.*, 2013). They are exposed to ambient temperature for a long time from capture to receiving at the depot. Ice was used in depots only when a bulk quantity was gathered and finally transported to the plants (Hassan, M.N., *et al.*, 2012). Serious quality deterioration has been reported at this stage (Kamal *et al.*, 2000). Hossain *et al.* (2010), quality of processing product very varies depend on quality of raw material. Therefore, to produce good quality shrimp and shrimp products, the raw shrimp should be wholesome. Moreover, the wholesomeness of finished products depends on the harvesting, postharvest care, transportation, handling and sanitation and hygiene aspects at farms, depots and industry levels (Faridullah, M., *et al.*, 2008). Some causes of postharvest quality losses from farm until processing plant have been reported. Paul, *et al.* (2010), but they did not present any lab-based practical on quality change each stage. Other researchers, Ali, M.Y., *et al.* (2013<sup>a</sup>), Ali, M.Y., *et al.* (2013<sup>b</sup>) have already carried out in this respect, but they only determined of protein losses in the value chain. So far, there are no researcher has been conducted on quality of shrimp during postharvest handling chain. Therefore, the aim of this study were to investigate of biochemical quality and microbiological content on the postharvest handling chain of shrimp in Mahakam delta region. The improvement of postharvest handling is necessary in order to produce excellent shrimp quality in Indonesia, especially in Mahakam delta region which could produce more competitive quality.

16

## II. Materials and Methods

### 2.1. Study area

The study was conducted at three subdistricts: sub district of Muara Jawa in Muara Pegah village, sub district of Anggana in Muara Pantuan village and sub district of Muara Badak in Saloapi village in Mahakam delta region. Two small ponds around 2-5 Ha in each subdistricts were used in unit observation. This size was selected since it was more manageable than bigger ponds.

### 2.2. Sampling collection

Black tiger shrimp (*P. monodon*) were obtained from three different sub districts in Mahakam delta : Muara Jawa, Anggana and Muara Badak (Fig. 1). The sample of 150 shrimps on average of 25-45 g were taken from each subdistrict on different time and the shrimp samples were divided into three stages of handling :

#### 1. Ponds

The 50 samples of whole shrimps were taken immediately after harvesting.

#### 2. Farmer point

After harvesting, farmer brought shrimp to farmer point. In this point, the farmer usually practice sorting (separate shrimp from fish and foreign materials), washing, beheading and icing (water chilling system). The farmers were used pond water/well water or potable water to handling of shrimp. In farmer point level, shrimps stored for 1-2 days and then shrimps brought to depot.

The 50 samples of headless shrimps were taken from farmer point.

#### 3. Depot

In depot level, the collector (depot owner call collector) applied sorting (freshness and size), washing and icing (water chilling system). Owner depot call collector. In this level, the shrimp stored for 2-3 days depend on volume of shrimp and then brought to the processing plant. The samples of 50 headless shrimps were taken from depot level.

All samples immediately placed into containers and added crushed ice with ratio is 1:1 (shrimps:ice) and brought to laboratory and stored for one day at refrigerator before being analyzed.




Fig.1. Map of Kutai Kartanegara Regency Showing Sampling Sites of Shrimp (*P. Monodon* F) at Ponds of Mahakam Delta Region. Blue line: Bordery of Mahakam delta region. (Source: Laboratory of Karto-Hidrooceanografi, Faculty of Fishery and Marine Science, Mulawarman University).

## 2.3. Biochemical quality determination

### 2.3.1. Determination of moisture content (AOAC, 1995)

Weighed accurately about 5 g of the shrimp muscles in a moisture dish with slip on cover. Dried in an air oven (Mimmert UNB 500) at 110-1°C for 5 hours. The dishes were then put and cool in a dessicator. The dish weighed quickly. Return the dish with the cover to and dry for another ½ hour. Cool in the dessicator and weigh again. Repeat until successive weights do not differ more than 1 mg.

### 2.3.2. Determination of TVB-N (Antonacopoulos and Vincke, 1989)

Determination of TVB-N was carried out by adding 10 g blended shrimp muscle sample in 300 ml distilled water in kjeldhal flask 500 ml. After addition of 2 g magnesium oxide (MgO) and 2-3 drops anti bumping granules and anti foaming silicone, the vessel is inserted into the preheated steam generator and the distillation is started immediately (steam flow approx. 10 ml/min). The



distillate is collected in the receiver, containing 10 ml boric acid and 8 drops Tashiro indicator, filled up to approx. 100 ml with distilled water. Distillation time is exactly 12 min: 10 min with condenser outlet immersed, 2 min above the surface of the distillation. Titration the distillate containing volatile basic nitrogen with 0.1 N HCL until the change colour point. Calculation:  $\text{Volume ml 0.1 N HCL} \times 14 = \text{mg TVB-N}/100\text{g}$ .

### 2.3.3. Determination of pH (AOAC, 2005)

pH of shrimp was measured at room temperature by ten (10) g samples were homogenized and mixed thoroughly with 100 ml distilled water and pH determined using the pH meter 330i (WTW, Germany).

### 2.3.4. Determination of Indole (Snellings *et al.*, 2003)

Indole was extracted from shrimp muscle by grinding 20 g shrimp with 50 ml toluene and 5 ml 5% TCA for 1 min. The puree was centrifuged for 30 min at 3500 rpm to separate the toluene from the shrimp pulp. The toluene layer was decanted and filtered through a 0.45  $\mu\text{m}$  ringe filter into a beaker containing anhydrous  $\text{N}_2\text{SO}_4$ . The absorbance of MeOH (as solvent) at bottom layer was measured with a Hach DR/2000 spectrophotometer at 567 nm. Unadulterated shrimp were used as the blank for the absorbance measurements. A Cary 50 Conc UV/VIS spectrophotometer was used in experiments.

### 2.4. Determination of Microbiological (TPC) (BSN, 2006a)

Twenty five g of the shrimp muscle tissue were aseptically taken and crushed in a mortar with a pestle. Shrimp homogenates were prepared by adding 9 mL of sterile distilled water to 1 g of the crushed samples. Thereafter, 1 mL each of homogenized solutions was inoculated using spread plate method into nutrient agar and incubated at 25°C for 24 hours. The total plate count was determined on the nutrient agar. After incubation the colonies were counted using colony forming unit and cfu/g was calculated.

### 2.5. Statistical Analysis

Results are expressed as mean  $\pm$  standard deviation. The significant ( $P < 0.05$ ) of the parameters were assessed by one way ANOVA test. SPSS (PASW), for windows (Version 22) was used for statistical analysis.

## III. Results and Discussion

### 3.1. Change of moisture content

The moisture content of postharvest shrimp tends to increase during handling process. The moisture content of the shrimp can be affected by postharvest handling of shrimp started in ponds. The moisture content of shrimp at different handling stages in Muara Jawa, Anggana, Muara Badak sub districts are represented in Fig.1. The moisture content was increased gradually during handling from ponds to depot varied significantly ( $P < 0.05$ ). The moisture content of fresh shrimps, prior to handling by farmers in three sub districts ranged of between 76.28%-78.02%. However, after handling by farmers at farmer point and collectors at depot, shrimp moisture content increased to 78.81- 81.59%. In Muara Jawa, moisture content of shrimp was 77.64%, 77.59% and 81.59% at ponds, farmer point and depot respectively. The moisture content of shrimp tends to increase during handling stages, except at farmer point. In Anggana, moisture content of shrimp was increased from ponds to depot. The moisture content was found as 78.02%, 79.42% and 80.87% at ponds, farmer point and depot respectively. In Muara Badak, the moisture content of shrimp was increased from pond to depot, where the higher moisture content was found at depot as 78.81% than at ponds and farmer point as 76.28% and 78.16% respectively. On an average basis moisture shrimp content increased by approximately 3.10% from ponds to depots in Mahakam delta region. Method of handling can influence shrimp quality. In generally, farmers and collectors in Mahakam delta region stored the shrimp at chilling water system. Chilling methods used greatly affect the moisture content of shrimp during handling. Hence, soaking shrimp in chilling water system can cause water uptake by shrimp meat or rehydration. Rachmawati *et al.* (2007), states that the increase in the moisture content

in the fish during storage in line with the continuing process of deterioration. The higher moisture content in the shrimp, the faster the shrimp decay process. In addition, soaking in ice plus water can cause shrimp softening and the loss of some nutrients in shrimp (Jeyasekaran *et al.*, 2006).




Fig.2. Moisture Content (%) of Shrimp at Different Stages of Postharvest Handling Chain in Three Sub districts of Mahakam Delta Region.

### 3.2. Changes of pH content

The results showed that the pH value of shrimp fluctuated from pond until depot at Muara Jawa, Anggana and Muara Badak sub districts (Fig. 1). In Muara Jawa, pH content of shrimp was not significantly different ( $P>0.05$ ), whereas in Anggana and Muara Badak sub districts were significant differences ( $P<0.05$ ) from ponds, farmer point and depot level. The pH content fluctuated was 6.89, 6.86 and 6.89 at ponds, farmer point and depots respectively in Muara Jawa sub district. In Anggana, the pH content found was 6.96, 6.55 and 6.69 at ponds, farmer point and depot respectively. In Muara Badak, the pH content was reorded as 6.91, 6.67 and 6.72 at ponds, farmer point and depot respectively. Nevertheless, the pH content of shrimp from Muara Jawa was higher than to other districts. The pH content of shrimp from the three sub districts of the present study agrees with Mohamed-Ali, F.H. (2011), the pH of marine products varies from 10 to 7.2 depending on conditions at the time of harvest, but the normal pH varies from 5.8 to 6.5. Walker and Betts (2000) reported that, ultimate pH of meat was significant for resistance to spoilage because most bacteria grow optimally at about pH 7 and not well below pH 4. The changes of pH value was caused by biochemical processes occuring during handling. After the death of shrimp, glycolysis occurs anaerobically. The process produces lactic acid lowering the pH value of shrimp. Value of pH increases after handling by the collectors (depot). Increasing pH value caused by the activities of proteolytic enzymes that degrade proteins result in volatile bases such as ammonia, methylamine compounds and trimethylamine (Abu-Bakar *et al.*, 2008).




Fig 3. pH Content of Shrimp at Different Stages of Postharvest Handling Chain at Three Subdistricts of Mahakam Delta Region.

### 3.3. Change of TVB-N content

One of parameter indicating level of freshness of fishery products is Total Volatile Base (TVB-N), which is volatile basic compounds consisting of ammonia, methylamine compounds, dimetilamin and trimethylamine, and other volatile basic compounds that contain nitrogen. The content of TVB-N shrimp continues to increase in line with storage time (Fig 3). The TVB-N content of shrimp at all stages of handling was significantly different at three sub districts ( $P < 0.05$ ). In Muara Jawa, the TVB-N content was recorded as 18.67(mgN/100g), 20.02(mgN/100g), 23.95(mgN/100g) at ponds, farmer point and depot respectively. In Anggana, the TVB-N content was 16.85(mgN/100g), 18.22(mgN/100g) and 19.93 (mgN/100g) at ponds, farmer point and depot. In Muara Badak, the TVB-N content of shrimp was detected as 9.92(mgN/100g), 11.41(mgN/100g) and 12.09(mgN/100g) at ponds, farmer point and depot. At farmer point level, TVB-N content of shrimp ranged of between 9.92 to 18.67(mgN/100g). TVB-N value of this study agree with Ali, M.Y., *et al.* (2013<sup>b</sup>), at farm level when shrimp were fresh, TVB-N content ranged between 15.30-19.91 (mgN/100g). In this study, on an average basis TVB-N content increased approximately 3.51 (mgN/100g) from pond level to depot level. However, The lowest TVB-N content of shrimp until the end of handling stages were found in Muara Badak sub district followed by shrimp from Anggana and Muara Jawa. The highest TVB-N content was found on shrimps from ponds to depot in Muara Jawa sub district. After harvesting (Muara Badak and Anggana), the shrimps were soaked in ice water immediately, so that can slow down the process of autolysis and microbial activity, except Muara Jawa subs district. Handling process by farmer used pond water or river water, without/inadequate of ice and low sanitation that can increased number of bacteria on the shrimps. The higher the content of TVB-N, the higher the rate of quality deterioration of shrimp (Zeng *et al.*, 2005). According to Nirmal and Benjakul (2011), TVB content significantly correlate with microbial content. Increased TVB content during handling process is considered to be caused by the degradation of protein or its derivatives by microbe activity that continues process of quality deterioration (Zeng *et al.*, 2005). The process of decline of the quality of the shrimp depends on the temperature of the water during soaking. The lower the temperature of ice water, the slower quality deterioration process occur (Jeyasekaran *et al.*, 2006). However, TVB-N content of shrimp from three subdistricts are still under TVB-N content limit for shrimp. According to Cobb *et al.* (1973) the limit is 30 mg N/100 g.




Fig 4. TVB-N Contents of Shrimp at Different Stages of Postharvest Handling Chain at Three Sub districts of Mahakam Delta Region.

### 3.4. Change of indole content

Indole content of shrimp at handling stages from 3 sub districts showed that indole has not been detected either in the pond or in the farmer point. Indole was first detected when the shrimp in depot (Table 1). The highest indole content were found in shrimp from Muara Jawa sub district followed the shrimps from two other sub districts. This condition indicates that the handling process of the shrimp at the depot has not done well; it means that deterioration has occurred. However, the indole value in shrimp from the 3 sub districts still in acceptable level of freshness, because the content of indole in shrimp that can be accepted by the US-FDA is 25  $\mu$ g/100g (Benner *et al.*, 2003). Indole can be used as an index of freshness of the shrimp. Indole enzymatically is produced through metabolism of proteins by bacteria. Proteolytic bacteria will hydrolyze the protein into free amino

acids using proteinase and peptidase enzymes. Furthermore, the amino acid tryptophan residues would undergo decarboxylation and deamination reactions catalyzed by the enzyme tryptophanase bacteria producing indole (Snellings *et al.*, 2003).

Table 1. Indole Content (mg/100g) of Shrimps at Different Stages of Postharvest Handling Chain in Three Sub districts of Mahakam Delta Region.

Handling stages	Subdistricts of Mahakam Delta Region		
	Muara Jawa	Anggana	Muara Badak
Ponds	ND	ND	ND
Farmer point	ND	ND	ND
Depot	1.25	1.12	1.10

ND = Not Detected

### 3.5. Change of Total Plate Count (TPC)

The TPC content of shrimp at different handling stages in three sub districts are presented in Fig 5. The statistically not significant variations ( $P>0.05$ ) in the content of shrimp TPC were observed among studied samples. In Muara Jawa, the TPC content was 5.17cfu/g, 5.23cfu/g and 5.30cfu/g at ponds, farmer point and depot respectively. In Anggana, the TPC content was found as 5.45cfu/g, 5.23cfu/g and 5.46cfu/g at ponds, farmer point and depot respectively. In Muara Badak, TPC content was recorded as 5.08cfu/g, 5.07cfu/g and 5.25cfu/g at ponds, farmer point and depot. However, the TPC content of shrimp from brackish water ponds in Muara Badak until the end of handling is the lowest compared to two other sub districts. Bacteria counts on brackish water *P. monodon* immediately after harvesting ranged from 2.23 to 4.20 (cfu/g) at 20°C and 3.36 to 4.20 (cfu/g) at 37°C (Reilly *et al.*, 1985). Harvesting temperature at ponds of study area was recorded at range 28-40°C. Decrease in TPC content of shrimp occurred at farmer point whereas farmers removed heads of shrimps (behead). Shrimp beheading is intended to remove most of bacteria, especially bacteria at head of shrimp. Novak (1973) stated that, the head occupies 40% of the total weight of shrimp and contains about 75% of bacteria. Increasing TPC content in shrimp could be caused by increase amount of bacteria that had already existed (depot). It is caused by inadequate ice amount used by farmers to lower temperature during handling. In addition, cross-contamination occurs between workers, equipment and water used during the handling process. The river water used by farmers in the sub district of Muara Jawa and Anggana can increase the content of bacteria (pond to depot). While the farmers and collector in the sub district of Muara Badak use well water/taps during the process of handling. Therefore, TPC content of shrimp from Muara Badak subdistrict was lowest compared to others subdistrict since at ponds until depot.




Fig. 5. TPC Contents of Shrimp at Different Stages of Postharvest Handling Chain in Three Subdistricts of Mahakam Delta Region.

According to Casani *et al.* (2006), microorganisms and organic matter in the water must be removed to prevent the growth of microorganisms that can cause spoilage and contamination of pathogenic bacteria. Also according to Hatha *et al.*, (2003), microbial content of the shrimp will continue to increase during transport and storage. The content of TPC in postharvest shrimp from



ponds in Mahakam delta are generally lower than the requirements according to SNI 7388:2009, TPC content must be less than  $5 \times 10^5$  or log 5.70 cfu / g (BSN, 2009).

### Conclusion

Postharvest handling stages at pond, farmer point and depot in the subdistricts of Muara Jawa, Anggana and Muara Badak were difference on quality of shrimp produced. The moisture content of shrimp from Muara Badak was lower than two sub districts until the depot. The pH content of shrimp not significance in three subdistricts, while TVB-N content of shrimp was lower on shrimp from Muara Badak compared to others until the end of handling chain. Indole content of shrimp was detected at depot of three districts. However, shrimp content from Muara Badak sub district was lower compared to shrimp from Muara Jawa and Anggana sub districts. The TPC content of shrimp from Muara Badak sub district was lowest at the end of handling stages (depot). In general, quality of shrimp from ponds until at depots still meet the requirements. In order to maintain good quality of shrimp from brackish water pond at Mahakam delta, the postharvest handling process by farmers and collectors in every stages of the handling chain has to be applied cold chain method, using good quality of water and good sanitation.

### Acknowledgements

Authors thank to the Fisheries and Marine agency, Kukar for supported financial for this research. Authors also thank to Iwan Suyatna as head of Laboratory of Karto-Hidrooceanografi, Faculty of Fishery and Marine Science Mulawarman University for contributing in the design mapping, Bagus Fajar Pamungkas for laboratory assistance.

### References

- [AOAC] Association of Analytical Chemistry. 2005. *pH of Acidified Foods*. Official Method 981.12.
- [AOAC] Association of Official Analytical Chemist. 1995. Official Method of Analysis of the Association of Official Analytical Chemist. Virginia.
- [BSN] National Bureau of Standards. 2006<sup>a</sup>. Indonesia National Standard, ISO 01-2332.3-2006. *Assesment Methods for Fishery Microbiology. Determination of Total Aerobic Plate Count*. National Standardization Agency. Jakarta.
- [BSN] National Bureau of Standards. 2006<sup>b</sup>. Indonesia National Standard, ISO 01-2728.1-2006. *Fresh Shrimp Part 1: Specification, Part 2: Raw Material Requirements and Part 3: Handling and Processing*. National Standardization Agency. Jakarta.
- Abu-Bakr, A, F. Salleh, AB, Razak, C.N.A. 2008. *Biochemical Changes of Fresh and Preserved Freshwater Prawn (Macrobrachium rosenbergii) During Storage*. International foog Research Journal. Vol. **15** (2) :181-190.
- Ali, M.Y., Z. Mahmud, Md. A. Rashed, M. Khanom and Md. G. Sarwer. 2013<sup>a</sup>. *Protein Losses Due To Post-Harvest Handling of Shrimp (Penaeus monodon) In The Value Chain of Khulna Region in Bangladesh*. International Journal of Engineering and Applied Sciences. **3** (1):56-63.
- Ali, M.Y., Z. Mahmud, Md. A. Rashed, M. Khanom and Md. G. Sarwer. 2013<sup>b</sup>. *Post-Harvest Quality Losses of Shrimp (Penaeus monodon) In The Value Chain of Southwestren Region in Bangladesh*. International Journal of Scientific Knowledge. **3** (2):36-43.
- Antonacopoulos N, Vyncke W. 1989. *Determination of volatile basic nitrogen in fish: a third collaborative study by the West European Fish Technologists' Association (WEFTA)*. Z Lebensm Unters Forsch, 189:309-316.
- Asikin, N.A. 2012. *Pattern and Handling Strategy of Postharvest Tiger Shrimp (Penaeus monodon F) From Brakish Water Pond in the Mahakam Delta Region*. Desertation. Doktor Programm of Coastal Management, Diponegoro University. 235 pp.
- Asikin, N.A., S. Hutabarat, YS. Darmanto, S. B. Prayitno. 2012. *Strategy of Qulity Improvement of Pond Shrimp Postharvest Manajement (Penaues monodon F) in Mahakam Delta (Case Study)*. International Journal of Science and Engeneering (IJSE). Vol. **3** (2):1-40.
- Benner, JR., RA, Staruszkiewicz, WF, Rogers, PL, Otwell, WS. 2003. *Evaluation of putrescine, cadaverine, and indole as chemical indicators of decomposition in Penaeid Shrimp*. Journal of Food Science **68**: 2178-2185.

- Casani, S., T. Leth, Susan. K. 2006. Water Reuse in a Shrimp Processing Line: Safety considerations using a HACCP approach. Elsevier. Food Control. **17**:540-550.
- Cobb, BF, Vanderzant, C., Thompson, CA, Custer, CS. 1973. *Chemical characteristics, bacterial counts and potential self-life of shrimp from various locations on the northwestern Gulf of Mexico*. J. Milk Food Technol. **36**: 463-468.
- Department of Marine and Fisheries Kukar. 2012. Aquaculture Statistical Report of 2011. Department of Marine and Fisheries Kutai Kartanegara District. Tenggarong. 32 h.
- Faridullah, M., N. Haider, M. Kamal and M. N. Islam. 2008. *Status of Penaeus monodon Farms, Depots and Processing Industries of Some Selected Areas of Banglades*. Progress. Agric. **19** (1) : 125-134.
- Hassan, M.N., M. Rahman, M.M. Hossain, A.A.K.M. Nowsad and M.B. Hossain. 2012. *Post-Harvest Handling and marketing of Shrimp and Prawn in South-Western Region of Bangladesh*. World Journal of Fish and Marine Sciences. **4** (6):651-656.
- Hatha Muhamed. A.A., T.K. Maqbool, S. Suresh Kumar. 2003. *Microbial Quality of Shrimp Products of Export Trade Produced from Shrimp Aquaculture*. International Journal of Food Microbiology. **82**:231-221.
- Jeyasekaran, G., Ganesan, P., Anandaraj, R., Shakila, RJ, Sukumar, D. 2006. *Quantitative and qualitative studies on the bacteriological quality of Indian white shrimp (Penaeus indicus) stored in dry ice*. Food Microbiology **23**: 526-533.
- Kamal, M., M.M. Rahman, L. Yasmin, S.U. Ahmed and M.N. Islam. 2000. *Studies on the post-mortem changes in shrimp and prawn during ice storage : II. Biochemical aspect of quality changes*. Bangladesh J. Fisheries res. **4**:91-96.
- Mohamed-Ali, H.F. 2011. *Quality Evaluation of Some Imported Fresh and Frozen Seafood*. Advance Journal of Food Science and Technology. **3** (1):83-88.
- Nirmal, N.P. and Benjakul, S. 2010. *Effect of catechin and ferulic acid on melanosis and quality of Pacific white shrimp subjected to freeze-thawing prior refrigerated storage*. Food Control. **21**: 1263-1271.
- Novak, A.F. 1973. Microbiological Consideration in the Handling and Processing of Crustacean shellfish in: C.O. Chichaster and HD. Graman (Eds). *Microbiological Safety of Fishery Product*: 59-73. Academy Press, New York-London.
- Paul, S., M.S. Reza, F.H. Sikha, M.N.A. Khan and M. Kamal. 2010. *Culture Practices and Quality Losses of Shrimp and Prawn at Different Stages of Handling and Transportation in Bangladesh*. Int. J. BioRes. **1**(4):07-13.
- R. Miget. 2010. *Shellfish Handling Practices-Shrimp and molluscs*. SRAC Publication No. 4902. Southern Regional Aquaculture Centre. The United States Department of Agriculture, National Institute of Food and Agriculture. 6 p.
- Rachmawati. N., R. Riyanto, F. Ariyani. 2007. *Formaldehyde Formation In Tiger Grouper (Ephinephelus fuscoguttatus) During Cold Storage Temperature*. Postharvest Journal of Marine and Fisheries, and Biotechnology. Vol. **2** (2) :137-145.
- Reilly, A., M. A. Bernarte and E. Dangla. 1985. *Quality Changes in Brackish Water Prawn (Penaeus Monodon) During Storage at Ambient Temperature in Ice and After Delays in Icing*. Reilly, A. (ed). FAO Fish Rep. (317) Suppl. :133-145. Republic of Bangladesh.
- Snellings, SL, Takenaka, NE, Kim-Hayes, Y., Miller, DW. 2003. *Rapid Colorimetric Method To Detect Indole in Shrimp With Gas Chromatography Mass Spectrometry Confirmation*. Journal of Food Science **68**: 1548-1553.
- Walker, S. J. and Betts, G. 2000. Factors Effecting the Microflora of Chilled Foods In: Chilled Foods. Stringer, M. and C. Dennis (Eds) Woodhead publishing. London, pp: 157-166.
- Zeng, QZ, Thorarinsdottir, KA, Olafsdottir, G. 2005. *Quality Changes of Shrimp (Pandalus Borealis) Stored Under Different Cooling Conditions*. Journal of Food Science **70**: 459-466.

# Quality Changes During Postharvest Handling Chain of Black Tiger Shrimp

## ARTIKEL SEMINAR faperta 2015-29 juni

### ORIGINALITY REPORT

17%

SIMILARITY INDEX

9%

INTERNET SOURCES

11%

PUBLICATIONS

5%

STUDENT PAPERS

### PRIMARY SOURCES

1

[dasnetgroup.com](http://dasnetgroup.com)

Internet Source

2%

2

Nicolaus Antonacopoulos. "Determination of volatile basic nitrogen in fish: A third collaborative study by the West European Fish Technologists' Association (WEFTA)", Zeitschrift für Lebensmittel-Untersuchung und -Forschung, 10/1989

Publication

2%

3

S.L. Akintola, S.B. Bakare. "Microbiological Changes in Freshwater Prawn (*Macrobrachium vollenhovenii*, Herklots 1857) Stored in Ice", American Journal of Food Technology, 2011

Publication

2%

4

S.L. Snellings. "Rapid Colorimetric Method to Detect Indole in Shrimp with Gas Chromatography Mass Spectrometry Confirmation", Journal of Food Science, 5/2003

Publication

1%

5	Mohamed A. Seif El-Yazal. "Changes in Metabolic Processes during Break Dormancy in Apple Buds under Foliar-Applied Garlic Extract", International Journal for Empirical Education and Research, 2018 Publication	1 %
6	Submitted to The University of the South Pacific Student Paper	1 %
7	banglajol.info Internet Source	1 %
8	eaas-journal.org Internet Source	1 %
9	Submitted to Universitas Mulawarman Student Paper	1 %
10	Submitted to Massey University Student Paper	1 %
11	www.mohfw.nic.in Internet Source	1 %
12	fssai.gov.in Internet Source	1 %
13	Swastawati, Fronthea, Eko Susanto, Bambang Cahyono, and Wahyu Aji Trilaksono. "Quality Characteristic and Lysine Available of Smoked Fish", APCBEE Procedia, 2012. Publication	<1 %



14	repository.unmul.ac.id	Internet Source	<1 %
15	alcalc.oxfordjournals.org	Internet Source	<1 %
16	ijoeear.com	Internet Source	<1 %
17	"Supplement A, July 2017", Journal of Food Protection, 2017	Publication	<1 %
18	maxwellsci.com	Internet Source	<1 %

Exclude quotes      On

Exclude bibliography      On

Exclude matches

< 12 words