

# The effect of deforestation and climate change on all-cause mortality and unsafe work conditions due to heat exposure in Berau, Indonesia: a modelling study

Nicholas H Wolff, Lucas R Vargas Zeppetello, Luke A Parsons, Ike Aggraeni, David S Battisti, Kristie L Ebi, Edward T Game, Timm Kroeger, Yuta J Masuda, June T Spector



## Summary

**Background** Previous studies focusing on urban, industrialised regions have found that excess heat exposure can increase all-cause mortality, heat-related illnesses, and occupational injuries. However, little research has examined how deforestation and climate change can adversely affect work conditions and population health in low latitude, industrialising countries.

**Methods** For this modelling study we used data at 1 km<sup>2</sup> resolution to compare forest cover and temperature conditions in the Berau regency, Indonesia, between 2002 and 2018. We used spatially explicit satellite, climate model, and population data to estimate the effects of global warming, between 2002 and 2018 and after applying 1·0°C, 1·5°C, and 2·0°C of global warming to 2018 temperatures, on all-cause mortality and unsafe work conditions in the Berau regency, Indonesia.

**Findings** Between 2002 and 2018, 4375 km<sup>2</sup> of forested land in Berau was cleared, corresponding to approximately 17% of the entire regency. Deforestation increased mean daily maximum temperatures by 0·95°C (95% CI 0·97–0·92;  $p < 0·0001$ ). Mean daily temperatures increased by a population-weighted 0·86°C, accounting for an estimated 7·3–8·5% of all-cause mortality (or 101–118 additional deaths per year) in 2018. Unsafe work time increased by 0·31 h per day (95% CI 0·30–0·32;  $p < 0·0001$ ) in deforested areas compared to 0·03 h per day (0·03–0·04;  $p < 0·0001$ ) in areas that maintained forest cover. With 2·0°C of additional future global warming, relative to 2018, deforested areas could experience an estimated 17–20% increase in all-cause mortality (corresponding to an additional 236–282 deaths per year) and up to 5 h of unsafe work per day.

**Interpretation** Heat exposure from deforestation and climate change has already started affecting populations in low latitude, industrialising countries, and future global warming indicates substantial health impacts in these regions. Further research should examine how deforestation is currently affecting the health and wellbeing of local communities.

**Funding** University of Washington Population Health Initiative.

**Copyright** © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

## Introduction

Heat exposure is a major human health risk, especially for people living in low latitude tropical countries where the ability to adapt to higher temperatures (ie, adaptive capacity) can be hindered by inadequate access to water, electricity, and other infrastructure. In the general population, particularly older people, very young children, and those with chronic diseases, heat-related mortality is projected to increase as a result of climate change.<sup>1,2</sup> These mortality effects are projected to be particularly pronounced in south-east Asia.<sup>3</sup> In working populations, including young and otherwise healthy workers, excess ambient heat exposure, combined with internal heat generated from physical labour, can lead to increased morbidity and mortality,<sup>4</sup> decreased productivity,<sup>5,6</sup> and increased risks of traumatic injury.<sup>7</sup> A growing body of research indicates that in tropical countries, both climate

change and deforestation are increasing temperatures and heat exposure,<sup>4,8–11</sup> but the combined risks of these changes have so far been underappreciated. For example, forest clearing in tropical countries can cause immediate increases in local temperatures of up to 8°C and exacerbate diurnal temperature variation.<sup>11</sup> The amount of warming can scale with larger deforested patches (>100 km<sup>2</sup>),<sup>12</sup> and the effects of warming can extend up to 50 km beyond deforested sites.<sup>13</sup> Yet, little is known about how warming associated with deforestation affects human health at large geographical scales (eg, >10 000 km<sup>2</sup>), or how these risks are likely to change in the future due to climate change.

Research on the human health risks of climate change has largely focused on urban areas in industrialised settings and on the potential health effects associated with heatwaves, other extreme weather events, and vector-borne

*Lancet Planet Health* 2021

Published Online  
November 11, 2021  
[https://doi.org/10.1016/S2542-5196\(21\)00279-5](https://doi.org/10.1016/S2542-5196(21)00279-5)

For the Bahasa translation of the abstract see [Online for appendix 1](#)

The Nature Conservancy, Brunswick, ME, USA (N H Wolff PhD); Department of Atmospheric Sciences (L R V Zeppetello MS, Prof D S Battisti, PhD), Department of Environmental and Occupational Health Sciences (Prof K L Ebi PhD, J T Spector MD), Department of Global Health (Prof K L Ebi), and Department of Medicine (J T Spector), University of Washington, Seattle, WA, USA; Nicholas School of the Environment, Duke University, Durham, NC, USA (L A Parsons PhD); Faculty of Public Health, Mulawarman University, Samarinda, Indonesia (I Aggraeni PhD); The Nature Conservancy, Brisbane, QLD, Australia (E T Game PhD); The Nature Conservancy, Arlington, VA, USA (T Kroeger PhD); Global Science, The Nature Conservancy, Arlington, VA, USA (Y J Masuda PhD)

Correspondence to:  
Dr Yuta J Masuda, Global Science, The Nature Conservancy, Arlington, VA 22203–1606, USA  
[yumasuda@tnc.org](mailto:yumasuda@tnc.org)

### Research in context

#### Evidence before this study

Although no formal literature search was done before undertaking the study, there is a vast body of published literature indicating that the effect of increasing temperatures on human health and wellbeing is a growing concern. Previous studies have found that excess heat exposure can increase all-cause mortality, heat-related illnesses, and occupational injuries, and can decrease productivity. Past work, however, has largely only examined heat events in urban, industrialised country settings. Little to no research has examined how deforestation, which causes substantial local temperature increases and also contributes to climate change, can adversely affect work conditions and population health in low latitude, industrialising countries.

#### Added value of this study

To the best of our knowledge, our modelling study is the first to estimate the effects of deforestation and climate change on all-cause mortality and unsafe work conditions from increases in heat exposure in some of the least resilient populations to climate change—populations in low latitude, industrialising countries. A major barrier to understanding and addressing how local (ie, deforestation) and global (ie, climate change) factors driving temperature increases affect the health of these populations is the paucity of representative human health data needed to understand the impact of these changes over large temporal and spatial scales. Our study overcomes this barrier by using spatially explicit data on tree cover change, land surface temperatures, and population distribution. We used established thresholds of heat health for worker safety to estimate the number of

lost safe work hours in a day and used established heat-mortality slopes to estimate changes in all-cause mortality from changes in mean daily temperatures. Finally, we used realistic projections for future temperature increases to assess the impact of climate change on people living in low latitude, industrialising countries.

#### Implications of all the available evidence

Our results indicate that heat exposure from deforestation has already significantly increased all-cause mortality and decreased safe work hours. Future temperature increases from climate change, even with the assumption of no further deforestation, will probably lead to an even more serious public health concern. Taken together, these results highlight the major challenge of the combined effects of deforestation and climate change for populations in low latitude, industrialising countries. Threats to health caused by increasing heat exposure will affect entire households and communities. Older people, very young children, and those with chronic diseases are particularly vulnerable to heat-related mortality, while reduced productivity among otherwise healthy workers will compound these impacts. Our findings point to an urgent need for action for the approximately 800 million people living in the world's tropical forest nations—a population that is expected to substantially increase by 2050. Importantly, these populations contribute the least to climate change but will bear a disproportionate amount of its adverse effects. Policy makers should therefore identify where and to what extent the local cooling effect of trees can address challenges to human health and wellbeing from increasing temperatures in low latitude, industrialising countries.

diseases.<sup>14–16</sup> Less research has been done to explore how environmental change that exacerbates climate change—such as deforestation—can have immediate and substantial local and regional effects on heat exposure<sup>13</sup> and human health for rural populations living in tropical countries. These populations are identified as being the least resilient to the effects of climate change<sup>10</sup> and are expected to experience substantial temperature increases,<sup>17</sup> indicating that they are at especially high risk of experiencing substantial adverse health effects. A primary barrier to understanding and addressing how local (ie, deforestation) and global (ie, climate change) factors driving temperature increases impact the health of these populations is the paucity of representative human health data needed to understand the impact of these changes over large temporal and spatial scales. Yet, when considered alongside recommendations such as those from the Intergovernmental Panel on Climate Change,<sup>18</sup> this information could have substantial implications for protecting human health as trade-offs between economic welfare, human health, and the natural environment are considered in the implementation of climate adaptation and mitigation recommendations.

The aim of this study was to estimate the effects of deforestation and climate change on all-cause mortality and unsafe work conditions due to heat exposure in the tropics. We focused our analysis on the Berau regency in Indonesia (hereafter referred to as Berau), which is emblematic of tropical forest conditions in other countries experiencing pressures on forests from expansion of agriculture, palm oil production, mining, and other activities.<sup>19</sup> We used satellite, climate model, and population data to quantify changes in heat exposure resulting from deforestation; estimate the effects of deforestation and climate change on all-cause mortality due to heat exposure; and estimate the effects and implications of deforestation and climate change on unsafe work conditions, assessed as changes in hours of work deemed unsafe by established occupational health guidelines for heat stress.

## Methods

### Study design and timeframe

For this modelling study we did a spatially explicit analysis of all-cause mortality and unsafe work conditions in Berau (appendix 2 p 6), comparing remotely sensed surface

See Online for appendix

temperature between 2002 and 2018 (observed), and after applying 1.0°C, 1.5°C, and 2.0°C of global warming relative to the 2008–27 mean, hereafter referred to as the present or 2018 baseline climate. We focused our analysis on 2002 and 2018 because the El Niño Southern Oscillation probably had minimal impacts on local climate in Indonesia during these years.<sup>20</sup> This approach allowed us to isolate the impacts of increasing atmospheric greenhouse gas concentrations and deforestation on local temperature changes. We chose the 1.0°C warming threshold relative to the 2018 baseline climate because this threshold corresponds to the Paris Climate Accords goal of limiting warming to less than 2.0°C relative to pre-industrial levels (approximately 1.0°C of warming relative to pre-industrial levels has already occurred). We also explored the impacts of an additional 1.5°C to 2.0°C of warming relative to present-day conditions because this is an increasingly likely outcome for the planet.<sup>21,22</sup> Our goal was to capture the relative additional effects of climate change on heat exposure (in terms of work hours lost and mortality) and thus we kept population, land use, and other factors constant. Our climate impacts should therefore be interpreted as conservative estimates of potential future impacts. Schematics of the study design and primary study components from spatially explicit data are presented in appendix 2 (p 6), and mirror analytical approaches that estimate environmental impacts on populations over large spatial scales.<sup>23–27</sup> An extended version of the methods is available in appendix 2 (p 1).

This study used spatially explicit data on climate, tree cover, population distributions, and data collected via the Global Burden of Disease, Injuries, and Risk Factors Study 2017 (GBD 2017). Analyses of these data did not require ethics approval.

### Data sources and approach

To estimate historical forest loss and gain, we re-gridded version 1.6 of Hansen and colleagues'<sup>28</sup> dataset (hereafter referred to as H13), a 30 m spatial resolution dataset based on data provided by Landsat satellite missions, to a 1×1 km resolution. The H13 dataset provides each grid cell's fractional forest cover in 2000 and the year during which forest loss (the year when cover goes from >0% to 0%) or forest gain (the year when cover goes from 0 to >50%) occurred.<sup>28</sup>


To estimate historical heat exposure (daily temperature and hourly heat index), we obtained values for surface daytime and night-time temperature at 1×1 km resolution using Moderate Resolution Imaging Spectroradiometer (MODIS) satellite observations.<sup>29</sup> In forested regions, the MODIS satellite detects top-of-canopy temperatures, rather than temperatures near the forest floor. However, once the forest has been removed, the satellite observes temperature at the land surface. Temperatures near the forest floor are systematically lower than those at the top of the canopy,<sup>30</sup> so our estimates of temperature change associated with deforestation are conservative.

We obtained the diurnal cycle of land surface temperature (“skt” variable) and relative humidity over Berau using hourly data from the fifth generation of the European Centre for Medium-Range Weather Forecast atmospheric reanalysis data product (ERA5)<sup>31</sup> for 2002 and 2018. Using the MODIS observations as the maximum and minimum values of the sinusoidal cycle, we created continuous diurnal cycles for each day based on the two temperatures observed by the MODIS satellite (appendix 2 p 8).

We estimated daily relative humidity at a 1×1 km resolution by assuming that the amount of water vapour in the atmosphere (specific humidity) is constant over Berau at daily timescales. This assumption, combined with our diurnal cycles of temperature, generated a relative humidity diurnal cycle that closely matched the one found in ERA5 over Berau. We then generated hourly estimates of heat indices using Rothfus's modification of Steadman's work,<sup>32</sup> which has also been used in recent studies estimating safe work hours for farm workers in the USA.<sup>27</sup>

To estimate future heat exposure, we used atmospheric surface air temperature output from global climate models that participated in the Coupled Model Intercomparison Project Phase 5 (CMIP5)<sup>33</sup> to estimate the average warming of Berau relative to global warming. We used output from all 39 CMIP5 models that did the RCP8.5 experiment (end-of-century radiative forcing of 8.5 W/m<sup>2</sup> relative to pre-industrial conditions), in which 21st century greenhouse gas emissions are not curtailed (appendix 2 p 12). Although we used RCP8.5 to calculate this relationship, the coefficient for warming is nearly identical in other warming experiments.

To estimate all-cause mortality, we calculated the population-weighted mean change in daily temperature for 2002–18 and for additional future global warming of 1.0°C, 1.5°C, and 2.0°C for Berau by overlaying our estimated daily mean temperature change with LandScan 2017 data,<sup>34</sup> which provide spatially explicit population data at 1×1 km resolution (appendix 2 p 14). We then used estimated relationships of heat-attributable and all-cause (Philippines) and non-external (Vietnam) heat-attributable excess mortality reported by Lee and colleagues<sup>35</sup> for the Philippines and Vietnam to construct estimates of changes in heat-related excess mortality for Berau. We selected Vietnam and the Philippines as sources of country-specific heat-related excess mortality curves for Berau because annual mean temperatures in these countries are similar to those of Berau (appendix 2 pp 16, 18) and no such curve exists for Indonesia nor can it be estimated because of the lack of daily mortality data,<sup>1,36,37</sup> and because these countries are similar to Berau with regard to key drivers of heat-related mortality (appendix 2 p 16).<sup>35,38</sup> Lee and colleagues'<sup>35</sup> heat-mortality slopes represent the estimated percentage-point increase in heat-attributable excess mortality (heat-attributable mortality divided by non-heat-attributable mortality) per °C increase in mean daily temperature.



**Figure 1: Forest cover and annual mean maximum daily land surface temperature in 2002 and 2018**

Each pixel equals 1 km<sup>2</sup> (N=25 053 pixels). Significant forest cover loss occurred between 2002 and 2018, as shown in panels A, C, and E, with a few regions of forest gain. Panels B, D, and F show that regions of greatest warming between 2002 and 2018 correspond to forest loss (red contour) and regions of greatest cooling corresponded to forest gain (blue contour).

For more on GBD cause of death data see <http://ghdx.healthdata.org/gbd-results-tool>

Mortality data for East Kalimantan were obtained from the GBD cause of death data for 2017. We used GBD mortality data for East Kalimantan because these province-level data are the highest-resolution GBD data available for Indonesia and because they provide both the all-cause and the non-external mortality rates needed for applying Lee and colleagues<sup>35</sup> heat-mortality slopes for the Philippines and Vietnam. Berau publishes only all-cause mortality data, which show an all-cause mortality rate comparable to East Kalimantan's overall mortality rate (appendix 2 p 19). We used rates of all-cause (596 deaths per 100 000 people) and non-external mortality (552 deaths per 100 000 people) in 2017 for East Kalimantan after confirming that the year was not an outlier. Notably, external causes of death comprise those due to injury or poisoning or other external causes (International Classification of Diseases, Tenth Revision, Clinical Modification [ICD-10] cause-of-death codes beginning with S, T, V, X or Y).

To assess lost safe work time due to unsafe work conditions, we used an implementation of the American Conference of Governmental Industrial Hygienists (ACGIH) Threshold Limit Value (TLV), intended for computing time-weighted average exposure levels and adapted for use with the heat index assuming sun exposure,<sup>39</sup> to compute the amount of time considered unsafe (ie, lost safe work time) in each hour (work-recovery cycle). Guidance for heat exposure, such as from WHO and the ACGIH, is based on maintaining the core body temperature within a safe range (eg, within 1.0°C of normal [37.0°C]).<sup>40</sup> As defined in the ACGIH TLV,<sup>41</sup> we assumed acclimatisation, recovery and rest in the shade, regular single-layer work clothes, and 415 W metabolic rate work, based on the literature for heavy physical work in agriculture or construction.<sup>41,42</sup> We compared lost safe work time in each day in 2002 and 2018, and after applying 1.0°C, 1.5°C, and 2.0°C of global warming relative to present conditions. In the analysis of future lost safe work time, we assumed no further deforestation or population changes and used the shape of the diurnal cycles of the heat index derived from the MODIS observations.

### Analyses

To analyse the relationship between deforestation and heat exposure, we computed 1×1 km pixel-level changes in temperature between 2002 and 2018, composited by geographical locations in Berau that experienced forest loss and maintained forest cover. These high-resolution data illustrate spatial relationships between the amount of forest cover and annual mean maximum daily land surface temperature in 2002 and 2018, and the difference between these years. We created histograms of mean daily maximum temperature differences between 2002 and 2018 for areas that kept and lost forest cover and computed the percentage of pixels with various degrees of warming that were co-located with pixels experiencing forest loss.

We also analysed the effects of deforestation and climate change on all-cause mortality due to heat exposure. Lee and colleagues<sup>35</sup> estimated heat-mortality slopes for the Philippines (9.15 percentage points per °C) with all-cause mortality data and for Vietnam (11.82 percentage points per °C) with non-external mortality data. We multiplied these slopes by the 2002–18 change in population-weighted mean annual temperature in Berau to estimate the percentage-point change in heat-attributable excess mortality due to the 2002–18 temperature change. We multiplied the GBD all-cause and non-external mortality rates in 2017 for East Kalimantan by Berau's 2018 population<sup>43</sup> from Statistics Indonesia of Berau (Badan Pusat Statistik Kabupaten Berau) to estimate Berau's all-cause and non-external mortality in 2018 (details of mortality and population data are provided in appendix 2 p 19). Because the 2018 mortality numbers already reflect the mortality impact from the 2002–18 temperature change, we calculated corrected, counterfactual (ie,

without the 2002–18 temperature change) mortality numbers for Berau, as follows:

$$\text{Actual mortality} / (1 + \text{percentage-point increase}_{\text{mortality}})$$

where percentage-point increase<sub>mortality</sub> is equal to the product of the heat-mortality slope and the observed mean population-weighted temperature change in Berau during 2002 and 2018. We then obtained the estimated 2018 mortality attributable to the change in mean daily temperature during 2002–18, as the difference between Berau's 2018 counterfactual all-cause and non-external mortality. Future all-cause mortality from an additional 1.0°C, 1.5°C, and 2.0°C of global warming was calculated by applying the local warming multiplier to estimate future population-weighted temperature changes in Berau. This was then used to estimate future changes in all-cause mortality.

To assess the impact of changes in lost safe work time due to heat exposure from deforestation and climate change, we estimated the total population affected by lost safe work time due to increases in heat exposure by overlaying estimates of lost work time with LandScan 2017 data.<sup>34</sup> We define the affected population as those living within any pixel where there have been increases in exposure to higher heat indices due to deforestation and climate change.


### Role of the funding source

The study was supported by a pilot research grant from the University of Washington Population Health Initiative. Researchers were independent from the funder. The funder had no role in study design, data collection, data analysis, data interpretation, writing of the report, or the decision to submit the Article for publication.

### Results

Between 2002 and 2018, 4375 km<sup>2</sup> of forested land in Berau was cleared (figure 1A–C), corresponding to approximately 17% of the entire regency, and more than 28% of the land below 200 m, where 98% of the population lives. The mean annual daily maximum land surface temperatures over the entire regency were relatively similar in 2002 (26.40°C [SD 1.33]) and 2018 (26.65°C [SD 1.64]; figure 1D, E). However, at the pixel level (1 km<sup>2</sup>), large temperature differences occurred between 2002 and 2018, ranging from an increase of 6.70°C to a decrease of 3.50°C (figure 1F). Most pixels (56%) showing a greater than 1.00°C increase between 2002 and 2018 corresponded to locations that experienced deforestation, and 80% of pixels that had the most significant cooling (<−3.00°C) occurred in the few locations that experienced forest gain (figure 1C, F).

Locations that lost forest cover between 2002 and 2018 showed a far greater relative proportion of temperature increases than locations that maintained forest cover (figure 2). The mean annual daily maximum temperature



**Figure 2: Histogram comparing mean annual maximum temperature differences between 2002 and 2018**

Areas that maintained forest cover shown in blue (N=18 979 pixels), and areas that lost forest cover shown in red (N=4375 pixels). Each pixel equals 1 km<sup>2</sup>.


	2002–18	>2018 warming scenarios		
		>1.0°C	>1.5°C	>2.0°C
Increase in mean daily temperature (°C)	0.41	0.9*	1.4*	1.9*
Mean increase in population-weighted mean daily temperature (°C)	0.86	0.9†	1.4†	1.9†
Heat-attributable increase in all-cause mortality (range)	7.3–8.5%	8.5–10.2%	12.8–15.3%	17.0–20.4%
Increase in heat-related mortality (deaths per year)‡	101–118	118–141§	177–212§	236–282§

\*Berau warms 0.94°C for every 1°C of global warming (appendix p 2). †Assumes 2017 population distribution remains unchanged and 2018 forest extent remains unchanged. ‡Based on heat-mortality slopes for the Philippines and Vietnam from Lee and colleagues (2019).<sup>35</sup> §At 2018 population size.


**Table: Estimates of the increase in temperature and annual heat-related mortality from 2002, Berau regency**

difference between 2002 and 2018 across pixels that lost forest cover was 1.03°C (SD 1.03) and across pixels that kept forest cover it was 0.08°C (0.67), indicating that, on average, deforestation drove 0.95°C (95% CI 0.97–0.92;  $p < 0.0001$ ) of additional warming. In addition to greater mean daily maximum temperatures, deforested areas experienced the majority of extreme warming between 2002 and 2018 (figure 2). More than 84% of the pixels with temperature increases greater than 2°C were co-located with pixels experiencing forest loss. 92.4% of the pixels with warming greater than 3°C were co-located with deforestation, as were 98.1% with warming greater than 4°C, and 100% with warming greater than 5°C.

In 2018, the population-weighted mean temperature in Berau was 0.86°C higher than in 2002, an increase that contributed to an estimated 7.3–8.5% of all-cause mortality in 2018, or 101–118 additional deaths per year as a result of 2002–18 climate warming and deforestation (table). An additional increase in global mean temperature of 1.0°C would increase the proportion of all-cause mortality attributable to heat in Berau by an additional 9% compared to 2018 estimates (equivalent to 118–141 additional annual deaths at current population



**Figure 3: Histogram comparing difference in work hours lost per day between 2002 and 2018**  
 Only pixels lower than 200 m are shown to reduce bias of cooler, higher elevation pixels. Each pixel equals 1 km<sup>2</sup>. Pixels that kept forest cover shown in blue (N=9972 pixels) and pixels that lost forest cover shown in red (N=3914 pixels).



**Figure 4: Estimates of impact of expected future global warming on work hours lost per day**  
 Panel A shows work hours lost in 2018; the same data are shown in appendix 2 (p 23) but are rescaled here for context. Panels B–D add 1.0°C, 1.5°C, and 2.0°C of global warming, to current 2018 conditions. These results assume no additional deforestation.

levels and assuming no further deforestation), while an increase in global mean temperature of 1.5°C would increase all-cause mortality by 14% (equivalent to 177–212 additional annual deaths), and an increase in global mean temperature of 2°C would increase all-cause mortality by 19% (equivalent to 236–282 additional annual deaths).

In areas that lost forest cover, the mean difference between 2002 and 2018 in safe work hours lost per day was 0.31 h (95% CI 0.30–0.32;  $p < 0.00001$ ), compared to

0.03 h (0.03–0.04;  $p < 0.00001$ ) in pixels that kept forest cover. Compared with 2002 (appendix 2 p 23), the spatial pattern in safe work hours lost per day in 2018 (mean 0.08 h [SD 0.23]; appendix 2 p 23) closely mirrored the spatial pattern of forest loss (figure 3; appendix 2 p 23). If only pixels below 200 m elevation are considered, 67.3% of pixels with safe work time lost per day of greater than 0.5 h were co-located with forest lost pixels. When pixel-average safe work loss increased to 1.0 h lost per day, the likelihood that the pixel had experienced deforestation increased to 77.2%, whereas with 1.5 h lost per day the likelihood of deforestation increased to 88.1%, and with 2.0 h lost per day it increased to 96.2%. Although increases in forest cover between 2002 and 2018 were rare (2.1% of pixels), these areas experienced an average gain in safe work hours of 0.02 h per day.

In 2018, LandScan data showed 13.2% of the total population in deforested pixels where at least some portion of the workday had hotter conditions than recommended for safe work. By contrast, LandScan data showed 4.0% of the population in forested pixels where at least some portion of the workday had hotter conditions than recommended for safe work. In areas exceeding 0.5 h of safe work lost per day, 9.5% of the total population was in deforested locations compared with 1.3% in forested locations. In areas exceeding lost safe work hours of 1.0 h per day, 4.7% of the population was in locations that lost forest cover compared with 0.7% in forested locations. For areas exceeding 1.5 h of safe work lost per day, 0.6% of the population was in deforested locations, compared with 0% in forested locations.


Figure 4 shows the spatial distribution of lost safe work hours when 1.0°C (figure 4B), 1.5°C (figure 4C), and 2.0°C (figure 4D) of global warming is applied to average present conditions (figure 4A). With 2.0°C of global warming, some locations in Berau would lose 5 h of safe work time per day (figure 4D). The spatial extent of conditions corresponding to 2 h or more of lost safe work hours per day ranged from 20 km<sup>2</sup> in 2018 (corresponding to 11.6% of Berau's total population that lives in two major urban centres, as shown in appendix 2 p 14) to 607 km<sup>2</sup> with an additional 2.0°C of global warming (corresponding to 49.5% of the population; figure 5). With 2.0°C of warming relative to present conditions, only 74 km<sup>2</sup> of forested areas (0.7% of total forested land below 200 m) will exhibit conditions leading to 2 h or more of lost safe work hours per day. By contrast, 392 km<sup>2</sup> of land that lost cover (10% of total land that was deforested since 2002) will exhibit conditions corresponding to 2 h or more of lost safe work hours per day. Thus, 65% of the pixels that show more than 2 h of lost safe work hours per day under a 2.0°C increase in global warming are associated with deforestation.

## Discussion

Forest loss in Berau is associated with significant increases in heat exposure, all-cause mortality, and

unsafe work conditions. Locally, deforestation has already caused temperature increases higher than 5°C. These observed temperature increases associated with deforestation exceed local projected end-of-century warming under high emission scenarios (approximately 2.2–5.1°C warmer than the present day under RCP8.5 projections; appendix 2 p 13), highlighting the immediate and drastic effects of deforestation on heat exposure. Importantly, temperature changes from deforestation occur at much shorter timescales than global climate change, which will progress over the course of decades to centuries. Any relative comparison of local temperature changes from deforestation will shift as the planet continues to warm. In our analysis, the increases in heat exposure between 2002 and 2018 were associated with an estimated increase in all-cause mortality that will approximately double again with an additional 2°C of global warming, which is likely to occur under high greenhouse gas emissions scenarios by 2077 (±12 years; appendix 2 p 13). Finally, we found that increases in unsafe work conditions due to heat exposure from 2002 to 2018 were associated with a ten-fold increase in the amount of lost safe work hours in deforested areas compared to forested areas, and that deforested areas might exhibit conditions corresponding to up to 5 h of safe work lost during an average workday with an additional 2°C of global warming relative to present conditions. Taken together, these findings highlight an urgent need for action, as threats to health caused by increasing heat exposure and mortality risks, particularly among older people, very young children, and those with chronic diseases, are compounded by the effects on household and community wellbeing resulting from reduced productivity among otherwise healthy workers.

The associated increases in all-cause mortality from increases in heat exposure are particularly noteworthy, and provide much needed estimates of the impacts of heat-related mortality in the tropics, where data are scarce.<sup>44,45</sup> Our estimates of the proportion of all-cause mortality associated with temperature increases between 2002 and 2018 (7.3–8.5%) indicate the substantial impact of heat-related mortality when compared with other major health challenges in the region. For example, the GBD 2017 data for East Kalimantan reported that maternal neonatal disorders accounted for 3.6%, neglected tropical diseases and malaria accounted for 1.2%, respiratory infections accounted for 6.8%, and transportation injuries accounted for 3.8% of total deaths. Our estimates of all-cause mortality rates with 2°C of additional global warming from 2018 levels indicate increases of 24–29% compared to 2002 levels, making heat-related mortality comparable to mortality due to other long-term public health challenges in Asia. For comparison, analyses from Asia indicate that tobacco smoking, which is prevalent particularly among men in Indonesia<sup>46</sup> and associated with increased risks of stroke and coronary heart disease, has been estimated to



**Figure 5: Impacts of deforestation and expected global warming on area (A) and population (B) exposed to more than 2 work hours lost per day**

Although the area exposed to more than 2 work hours lost per day is small relative to the total area below 200 m (15 225 km<sup>2</sup>), the population exposed is significant. 98% of the total population lives below 200 m, where most of the deforestation has occurred to date.

account for 29.3% of all-cause mortality in individuals born on or after 1930.<sup>47</sup>

Heat exposure can contribute to mortality in several ways. A vast amount of published literature has shown that higher air temperatures directly increase mortality and morbidity in the general population by causing heat stroke, heat exhaustion, and dehydration, as well as indirectly by exacerbating existing cardiovascular and renal diseases, leading to ischaemic stroke, ischaemic heart disease, cardiac dysrhythmia, hypotension, and acute renal failure.<sup>48,49</sup> For populations with low adaptive capacity, the relationship between temperature increases and all-cause mortality might be more pronounced, as shown by Lee and colleagues.<sup>35</sup> Recent studies in Berau have found that even under favourable work conditions, working in deforested versus forested areas for just 90 min can result in elevated core body temperatures exceeding 38.5°C<sup>50</sup> and decreased cognitive performance.<sup>51</sup> For outdoor workers engaged in heavy physical activities in the tropics, where heat and humidity are already high, productivity declines can occur with increasing heat stress, as the human body naturally responds to heat stress by triggering reductions in physical work intensity and internal heat generation.<sup>4</sup>

Our mortality projections assume that exposure to, and the mortality risk from, higher temperatures will not be mediated through additional adaptive responses beyond those already implemented.<sup>11</sup> Given the constraints on adaptive behaviours, this is a reasonable assumption. Although the slopes of heat-mortality dose–response curves have decreased in several high-income countries,<sup>52,53</sup> this decline was correlated with

increases in air conditioning use,<sup>54</sup> advances in per-capita income or health-care access and quality,<sup>55</sup> reduced prevalence of cardiovascular mortality risk factors such as smoking,<sup>56</sup> and policies specifically targeted at reducing heat vulnerability.<sup>55</sup> For rural communities that largely lack access to infrastructure and markets, there is limited evidence that the slope of the heat-mortality dose–response curve will decrease in the near future.

Outdoor work is common in areas experiencing deforestation, as human-initiated land use pressures remain the primary driver of forest loss, especially in areas experiencing high land use pressure, such as Berau.<sup>19</sup> Deforestation is driven largely by outdoor labour-intensive industries, such as mining, farming, and palm oil production.<sup>19</sup> Even under the optimistic assumption of no further deforestation, which is extremely unlikely given current trends,<sup>57</sup> deforested areas could exhibit conditions corresponding to 5 h of unsafe work during the workday by the end of the 21st century. This finding is important, as a 2019 study in the same region<sup>11</sup> found that outdoor labourers worked, on average, 6.5 h a day and are already shifting work schedules to avoid the hottest times of the day. Even an average loss of 2 work hours in deforested areas would probably be detrimental to livelihoods without some adaptive strategies, such as shifting to non-outdoor livelihood activities, as that would account for a third of the time allocated to work. In subsistence agricultural settings that are common in Berau,<sup>58</sup> there might be more flexibility in organisation of work to allow a slower pace of work and increase rest-taking behaviours compared to industrial agricultural settings.<sup>7</sup> However, a study of survey data on self-reported adaptation strategies found that outdoor workers in Berau are already having to adapt to hotter temperatures due to deforestation,<sup>11</sup> suggesting that those engaged in outdoor work might already be approaching their adaptive capacity through behavioural adaptations.

Our findings underscore the major challenge of the combined effects of deforestation and climate change for the approximately 800 million people living in the world's tropical forest nations,<sup>59</sup> a population that is expected to substantially increase by 2050.<sup>60</sup> Rural populations in these countries contribute the least to global emissions,<sup>17</sup> yet bear a disproportionate burden of the adverse effects of climate change and deforestation.<sup>61</sup> Forests have the potential to increase community resilience to temperature increases from climate change. In our analysis, the mean annual daily maximum temperature increase was 92% lower, and lost work hours were 90% lower, in areas that maintained forest cover than in deforested areas between 2002 and 2018. Furthermore, the few areas experiencing forest cover gain saw an average gain of 0.02 h of safe work per day. There are two primary mechanisms by which forests contribute cooling services. Trees can cool local areas through shade (ie, blocking direct solar radiation) and transpiration of liquid water.<sup>62,63</sup> In tropical forests in particular, hundreds of litres of water

a day are transpired by individual trees, creating an amount of cooling that is equivalent to two average household air-conditioning units.<sup>63</sup> When deforestation occurs, it removes shade from local land cover and the deep roots that contribute much of the liquid water for tree transpiration. The result is a dry surface soil layer that can no longer effectively cool through evapotranspiration.

We note several limitations of our study that future research should address. First, although our study used established methods for estimating environmental impacts on people,<sup>25</sup> similarly to past studies we did not estimate the causal effect of deforestation-associated heat on people due to data and other limitations. There are still no optimal approaches (eg, an ensemble of models) for large-scale studies that allow us to adequately address the uncertainty in climate and health projections that results from variability in population characteristics, adaptive capacity, and bioclimate model structures.<sup>64</sup> However, our approach expands on the available evidence by adapting approaches grounded in laboratory-based, controlled human studies intended for observational use that integrate consideration of factors such as clothing, metabolic rate, and work–rest cycles that are drivers of heat stress exposure, in addition to ambient conditions, among outdoor working populations.

Second, our analyses of future warming held population constant and assumed no further deforestation, and did not assume changes in urbanisation, technological change, or economic growth. Third, our estimates of changes in heat-related mortality were based on relationships of temperature and heat-related excess mortality from Vietnam and the Philippines rather than on relationships specific to Berau or East Kalimantan. Although both Vietnam and the Philippines are closely matched to Berau in terms of the climatic, social, and health characteristics strongly determinant of heat-related mortality (appendix 2 p 16), the use of data from these countries could have biased our mortality estimates. Estimation of relationships of temperature and heat-related excess mortality specific to Berau or East Kalimantan require time-series data of daily mortality and temperature, which are currently not available.<sup>65</sup> However, our approach followed best standard practice for estimating mortality impacts in data-scarce or data-poor environments.<sup>65</sup>

Fourth, we used the heat index instead of wet bulb globe temperatures (WBGT) to calculate lost safe work hours due to temperature increases. WBGT captures important elements for assessing the risk of heat stress, but at the spatiotemporal resolution needed for our study it was unfeasible to estimate WBGT without making unrealistically strong assumptions. However, recent work<sup>66</sup> supports the use of the heat index in lieu of WBGT when WBGT measurements are impractical, and we believe the approach we used<sup>39</sup> provides a valid and valuable comparison between forested and deforested areas, thus allowing us to estimate hourly heat exposure and work–recovery cycles at a more granular temporal



level than has previously been published. Importantly, the heat index is also a commonly used and understood measure, as it is used in heat advisories by the US National Weather Service and the US Occupational Safety and Health Administration. Estimates of the total number of people experiencing increasing heat exposure could be improved with more detailed spatially explicit demographic data, such as data that provide age-stratified population counts. However, these data currently do not exist at the granularity that is needed.

Fifth, we recognise that heat index is more variable across space and time than captured here. The 1 km<sup>2</sup> resolution MODIS observations dampen the temperature extremes people experience on the ground, as do our use of mean diurnal cycles of ERA5 temperature. Furthermore, MODIS observations from the forest canopy are warmer than conditions in the forest understory,<sup>30</sup> the location of interest. Last, our assumption of constant specific humidity across Berau is most certainly a simplification of the water vapour variability (and thus relative humidity) that occurs between locations and across time. Overall, it is likely that our methodological assumptions provide conservative (high) estimates of the forest understory heat indices (appendix 2 pp 1–2). Although this limitation has no practical impact on our primary results because work hours lost are minimal in forest locations, it does suggest that we could be underestimating the cooling services provided by forests. This highlights the need for more in-situ observations comparing heat exposure across tropical landscapes.<sup>11</sup>

Our study raises concerns that deforestation, which exacerbates climate change, can have a substantial adverse effect on the health and wellbeing of populations in tropical countries, and indicates important public health benefits that could be achieved from the conservation or restoration of forests or forested landscapes. The benefits to public health are likely to manifest in both the short and long term, especially given current trajectories for global warming. Policy makers should actively prepare for, and manage, the health risks associated with deforestation, as deforestation events are relatively predictable and can be influenced by land use and management plans. Development of policies and land use decisions should be based on analyses that compare different land use and climate scenarios and consider the implications for human health and wellbeing. Preventive strategies to reduce health risks from heat exposure are often beyond the control of individuals, and systemic changes at multiple levels, including within the overall policy environment, are needed.<sup>7</sup> For rural communities in tropical countries, these ecosystem services play a particularly crucial role in maintaining health and wellbeing. Natural climate solutions,<sup>67</sup> which include the protection of forests and agroforestry practices, could contribute to reductions in heat exposure, improved work productivity, and reductions in mortality risks.

Clinicians and policy makers should therefore carefully evaluate and prioritise interventions that maximise population health and wellbeing while addressing climate change mitigation.

#### Contributors

NHW, LRVZ, LAP, DSB, ETG, TK, YJM, and JTS designed the research. NHW, LRVZ, LAP, TK and YJM did the data analysis, and NHW, LRVZ, LAP, DSB, TK, YJM, and JTS interpreted data analysis outputs. NHW, LRVZ, LAP, DSB, IA, KLE, ETG, TK, YJM, and JTS wrote the Article. All authors read and approved the final manuscript. NHW, LRVZ, LAP, DSB, IA, KLE, ETG, TK, YJM, and JTS had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. NHW, LRVZ, LAP, TK, and YJM verified the underlying data. The corresponding author (YJM) attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

#### Declaration of interests

We declare no competing interests.

#### Data sharing

All data, codes, and materials used in the analysis are available from the authors upon reasonable request via email to the corresponding author.

#### Acknowledgments

This work contains modified Copernicus Climate Change Service Information. The forest cover could not have been analysed without the National Aeronautics and Space Administration (NASA) and the US Geological Survey partnership that resulted in the Landsat mission. Support for this work was provided by a pilot research grant from the University of Washington Population Health Initiative. LAP thanks the Washington Research Foundation postdoctoral fellowship for funding. NHW and YJM thank the Bezos Earth Fund for their support.

#### References

- Mora C, Dousset B, Caldwell IR, et al. Global risk of deadly heat. *Nat Clim Chang* 2017; **7**: 501–06.
- Gasparrini A, Guo Y, Sera F, et al. Projections of temperature-related excess mortality under climate change scenarios. *Lancet Planet Health* 2017; **1**: e360–67.
- Vicedo-Cabrera AM, Sera F, Guo Y, et al. A multi-country analysis on potential adaptive mechanisms to cold and heat in a changing climate. *Environ Int* 2018; **111**: 239–46.
- Spector JT, Sheffield PE. Re-evaluating occupational heat stress in a changing climate. *Ann Occup Hyg* 2014; **58**: 936–42.
- Burke M, Hsiang SM, Miguel E. Global non-linear effect of temperature on economic production. *Nature* 2015; **527**: 235–39.
- Masuda YJ, Garg T, Anggraeni I, et al. Warming from tropical deforestation reduces worker productivity in rural communities. *Nat Commun* 2021; **12**: 1601.
- Spector JT, Masuda YJ, Wolff NH, Calkins M, Seixas N. Heat exposure and occupational injuries: review of the literature and implications. *Curr Environ Heal Reports* 2019; **6**: 286–96.
- Flouris AD, Dinas PC, Ioannou LG, et al. Workers' health and productivity under occupational heat strain: a systematic review and meta-analysis. *Lancet Planet Health* 2018; **2**: e521–31.
- Andrews O, Le Quéré C, Kjellstrom T, Lemke B, Haines A. Implications for workability and survivability in populations exposed to extreme heat under climate change: a modelling study. *Lancet Planet Health* 2018; **2**: e540–47.
- Watts N, Amann M, Ayeb-Karlsson S, et al. The *Lancet* Countdown on health and climate change: from 25 years of inaction to a global transformation for public health. *Lancet* 2018; **391**: 581–630.
- Masuda YJ, Castro B, Aggraeni I, et al. How are healthy, working populations affected by increasing temperatures in the tropics? Implications for climate change adaptation policies. *Glob Environ Chang* 2019 May; **56**: 29–40.
- Vargas Zeppetello L, Parsons LA, Spector JT, et al. Large scale tropical deforestation drives extreme warming. *Environ Res Lett* 2020; **15**: 084012.
- Cohn AS, Bhattarai N, Campolo J, et al. Forest loss in Brazil increases maximum temperatures within 50 km. *Environ Res Lett* 2019; **14**: 084047.

- 14 Patz JA, Campbell-Lendrum D, Holloway T, Foley JA. Impact of regional climate change on human health. *Nature* 2005; **438**: 310–17.
- 15 Smith KR, Woodward A, Campbell-Lendrum D, et al. Human health: impacts, adaptation, and co-benefits coordinating. In: Field CB, Barros VR, Dokken DJ, et al, eds. In: *Climate change 2014: impacts, adaptation, and vulnerability. Part A: global and sectoral aspects. Contribution of working group II to the fifth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, 709–54.
- 16 Haines A, Kovats RS, Campbell-Lendrum D, Corvalan C. Climate change and human health: impacts, vulnerability and public health. *Public Health* 2006; **120**: 585–96.
- 17 King AD, Harrington LJ. The inequality of climate change from 1.5 to 2°C of global warming. *Geophys Res Lett* 2018; **45**: 5030–33.
- 18 UN Environment Programme. Global Environment Outlook – GEO-6: healthy planet, healthy people. <https://wedocs.unep.org/handle/20.500.11822/27539> (accessed Oct 11, 2021).
- 19 Griscom BW, Ellis PW, Baccini A, Marthinus D, Evans JS. Ruslandi. Synthesizing global and local datasets to estimate jurisdictional forest carbon fluxes in Berau, Indonesia. *PLoS One* 2016; **11**: e0146357.
- 20 Huang B, Thorne PW, Banzon VF, et al. Extended reconstructed sea surface temperature, version 5 (ERSSTv5): upgrades, validations, and intercomparisons. *J Clim* 2017; **30**: 8179–205.
- 21 Raftery AE, Zimmer A, Frierson DMW, Startz R, Liu P. Less than 2°C warming by 2100 unlikely. *Nat Clim Chang* 2017; **7**: 637–41.
- 22 Christensen J, Olhoff A. Lessons from a decade of emissions gap assessments. Nairobi: UN Environment Programme, 2019.
- 23 Chavaillaz Y, Roy P, Partanen AI, et al. Exposure to excessive heat and impacts on labour productivity linked to cumulative CO<sub>2</sub> emissions. *Sci Rep* 2019; **9**: 13711.
- 24 Xu C, Kohler TA, Lenton TM, Svenning JC, Scheffer M. Future of the human climate niche. *Proc Natl Acad Sci USA* 2020; **117**: 11350–55.
- 25 Dunne JP, Stouffer RJ, John JG. Reductions in labour capacity from heat stress under climate warming. *Nat Clim Chang* 2013; **3**: 563–66.
- 26 Li D, Yuan J, Kopp RE. Escalating global exposure to compound heat-humidity extremes with warming. *Environ Res Lett* 2020; **15**: 064003.
- 27 Tigchelaar M, Battisti D, Spector J. Work adaptations insufficient to address growing heat risk for U.S. agricultural workers. *Environ Res Lett* 2020; **15**: 094035.
- 28 Hansen MC, Potapov PV, Moore R, et al. High-resolution global maps of 21st-century forest cover change. *Science* 2013; **345**: 850–53.
- 29 Wan Z, Hook S, Hulley G. MOD11A1 MODIS/Terra Land Surface Temperature/Emissivity Daily L3 Global 1km SIN Grid V006. NASA EOSDIS L Process DAAC, 2015. <https://lpdaac.usgs.gov/products/mod11a1v006/> (accessed Oct 11, 2021).
- 30 Thompson OE, Pinker RT. Wind and temperature profile characteristics in a tropical evergreen forest in Thailand. *Tellus* 1975; **27**: 562–73.
- 31 Climate Change Service. ERA5: fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store, 2017. <https://cds.climate.copernicus.eu/cdsapp#!/home> (accessed Oct 11, 2021).
- 32 National Oceanic and Atmospheric Administration, US Department of Commerce. The heat index equation. National Weather Service. 2014. [https://www.wpc.ncep.noaa.gov/html/heatindex\\_equation.shtml](https://www.wpc.ncep.noaa.gov/html/heatindex_equation.shtml) (accessed Oct 11, 2021).
- 33 Taylor KE, Stouffer RJ, Meehl GA. An overview of CMIP5 and the experiment design. *Bull Am Meteorol Soc* 2012; **93**: 485–98.
- 34 Bhaduri B, Bright E, Coleman P, Urban ML. LandScan USA: a high-resolution geospatial and temporal modeling approach for population distribution and dynamics. *GeoJournal* 2007; **69**: 103–17.
- 35 Lee JY, Kim H, Gasparrini A, et al. Predicted temperature-increase-induced global health burden and its regional variability. *Environ Int* 2019; **131**: 105027.
- 36 Campbell S, Remenyi TA, White CJ, Johnston FH. Heatwave and health impact research: a global review. *Health Place* 2018; **53**: 210–18.
- 37 Green H, Bailey J, Schwarz L, Vanos J, Ebi K, Benmarhnia T. Impact of heat on mortality and morbidity in low and middle income countries: a review of the epidemiological evidence and considerations for future research. *Environ Res* 2019; **171**: 80–91.
- 38 Hajat S, Kosatky T. Heat-related mortality: a review and exploration of heterogeneity. *J Epidemiol Community Health* 2010; **64**: 753–60.
- 39 Bernard TE, Iheanacho I. Heat index and adjusted temperature as surrogates for wet bulb globe temperature to screen for occupational heat stress. *J Occup Environ Hyg* 2015; **12**: 323–33.
- 40 WHO. Health factors involved in working under conditions of heat stress: report of a WHO scientific group [meeting held in Geneva from 29 August to 4 September 1967]. Geneva: World Health Organization, 1969.
- 41 ACGIH. 2015 TLVs and BEIs: based on the documentation of the threshold limit values for chemical substances and physical agents & biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists, 2015.
- 42 Kjellstrom T, Freyberg C, Lemke B, Otto M, Briggs D. Estimating population heat exposure and impacts on working people in conjunction with climate change. *Int J Biometeorol* 2018; **62**: 291–306.
- 43 Statistics Indonesia of Berau. Berau in numbers 2018. Tanjung Redeb: Badan Pusat Statistik Kabupaten Berau, 2019. <https://beraukab.bps.go.id/> (accessed Nov 2, 2021).
- 44 Longden T, Quilty S, Haywood P, Hunter A, Gruen R. Heat-related mortality: an urgent need to recognise and record. *Lancet Planet Health* 2020; **4**: e171.
- 45 Bloomberg MR, Bishop J. Understanding death, extending life. *Lancet* 2015; **386**: e18–19.
- 46 Jha P, MacLennan M, Chaloupka FJ, et al. Global hazards of tobacco and the benefits of smoking cessation and tobacco taxes. In: *Disease control priorities, third edition (volume 3): cancer*. The World Bank; 2015: 175–93.
- 47 Yang JJ, Yu D, Wen W, et al. Tobacco smoking and mortality in Asia: a pooled meta-analysis. *JAMA Netw Open* 2019; **2**: e191474.
- 48 Basu R. High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008. *Environ Health* 2009; **8**: 40.
- 49 Hopp S, Dominici F, Bobb JF. Medical diagnoses of heat wave-related hospital admissions in older adults. *Prev Med* 2018; **110**: 81–85.
- 50 Suter MK, Miller KA, Anggraeni I, et al. Association between work in deforested, compared to forested, areas and human heat strain: an experimental study in a rural tropical environment. *Environ Res Lett* 2019; **14**: 084012.
- 51 Masuda YJ, Garg T, Anggraeni I, et al. Heat exposure from tropical deforestation decreases cognitive performance of rural workers: an experimental study. *Environ Res Lett* 2020; **15**: 124015.
- 52 Lee W, Choi HM, Kim D, Honda Y, Guo Y-LL, Kim H. Temporal changes in mortality attributed to heat extremes for 57 cities in Northeast Asia. *Sci Total Environ* 2018; **616–17**: 703–09.
- 53 Gasparrini A, Guo Y, Hashizume M, et al. Temporal variation in heat-mortality associations: a multicountry study. *Environ Health Perspect* 2015; **123**: 1200–07.
- 54 Hondula DM, Balling RC, Vanos JK, Georgescu M. Rising temperatures, human health, and the role of adaptation. *Curr Clim Chang Reports* 2015; **1**: 144–54.
- 55 Achebak H, Devolder D, Ballester J. Heat-related mortality trends under recent climate warming in Spain: a 36-year observational study. *PLoS Med* 2018; **15**: e1002617.
- 56 Bobb JF, Peng RD, Bell ML, Dominici F. Heat-related mortality and adaptation to heat in the United States. *Environ Health Perspect* 2014; **122**: 811–16.
- 57 Song XP, Hansen MC, Stehman SV, et al. Global land change from 1982 to 2016. *Nature* 2018; **560**: 639–43.
- 58 Statistics Indonesia of Berau. Berau Regency in figures 2017. Tanjung Redeb: Badan Pusat Statistik Kabupaten Berau, 2018. <https://beraukab.bps.go.id/> (accessed Nov 2, 2021).
- 59 Chao S. Forest peoples: numbers across the world. 2012. Forest Peoples Programme. [https://www.forestpeoples.org/sites/default/files/publication/2012/05/forest-peoples-numbers-across-world-final\\_0.pdf](https://www.forestpeoples.org/sites/default/files/publication/2012/05/forest-peoples-numbers-across-world-final_0.pdf) (accessed Oct 11, 2021).
- 60 James Cook University. 2019 State of the Tropics report. June 27, 2019. <https://www.jcu.edu.au/state-of-the-tropics/publications/2019-state-of-the-tropics-report> (accessed Oct 11, 2021).
- 61 Coffel ED, Horton RM, de Sherbinin A. Temperature and humidity based projections of a rapid rise in global heat stress exposure during the 21<sup>st</sup> century. *Environ Res Lett* 2018; **13**: 014001.

- 
- 62 Bright RM, Davin E, O'Halloran T, Pongratz J, Zhao K, Cescatti A. Local temperature response to land cover and management change driven by non-radiative processes. *Nat Clim Chang* 2017; **7**: 296–302.
- 63 Ellison D, Morris CE, Locatelli B, et al. Trees, forests and water: cool insights for a hot world. *Glob Environ Change* 2017; **43**: 51–61.
- 64 Vanos JK, Baldwin JW, Jay O, Ebi KL. Simplicity lacks robustness when projecting heat-health outcomes in a changing climate. *Nat Commun* 2020; **11**: 6079.
- 65 Marlier ME, Liu T, Yu K, et al. Fires, smoke exposure, and public health: an integrative framework to maximize health benefits from peatland restoration. *GeoHealth* 2019; **3**: 178–89.
- 66 Garzón-Villalba XP, Ashley CD, Bernard TE. Benchmarking heat index as an occupational exposure limit for heat stress. *J Occup Environ Hyg* 2019; **16**: 557–63.
- 67 Griscom BW, Adams J, Ellis PW, et al. Natural climate solutions. *Proc Natl Acad Sci USA* 2017; **114**: 11645–50.