

# Review of Angiotensin-converting Enzyme Inhibitory Assay: Rapid Method in Drug Discovery of Herbal Plants

*by* Islamudin Ahmad

---

**Submission date:** 02-Sep-2019 08:12AM (UTC-0700)

**Submission ID:** 1166262148

**File name:** PhcogRev\_Vol\_11\_21\_1.pdf (498.46K)

**Word count:** 6055

**Character count:** 33589

# Review of Angiotensin-converting Enzyme Inhibitory Assay: Rapid Method in Drug Discovery of Herbal Plants

Islamudin Ahmad<sup>1,2</sup>, Arry Yanuar<sup>2</sup>, Kamarza Mulia<sup>3</sup>, Abdul Mun'im<sup>2</sup>

<sup>1</sup>Department of Pharmaceutical Sciences, Faculty of Pharmacy, Mulawarman University, Samarinda, East Kalimantan 75116, <sup>2</sup>Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Indonesia, <sup>3</sup>Department of Chemical Engineering, Faculty of Engineering, Universitas Indonesia, Depok, West Java 16424, Indonesia

## ABSTRACT

The renin-angiotensin-aldosterone system is a signaling pathway which responsible in the blood pressure regulation. Angiotensin-converting enzyme (ACE) is one of the key elements responsible for the hypertensive mechanism. It converts angiotensin-I to angiotensin-II. The discovery history of the ACE inhibitory activity assay method has been through a long stage for decades and development continues until today. The ACE inhibitory activity has become an effective screening method in the search for new antihypertensive agents from herbal plants. Some of *in vitro* assay methods were used to examine the activity of ACE inhibitors based on the substrate usage, such as; Cushman and Cheung Method using a substrate hippuryl-histidyl-leucine (HHL), Holmquist method using a substrate furanacryloyl-tripeptide, Elbl and Wagner method using a substrate benzoyl-[I-14C] glycyl-L-histidine-L-leucine, Carmel and Yaron method using a substrate o-aminobenzoyl-glycyl-p-nitrophenylalanilproline, and Lam method using 3-hydroxybutyryl-glycyl-glycyl-glycine as substrate. Several different methods to measure the results of enzymatic reactions or separating substrate with products, including spectrophotometric, fluorometric, high-performance liquid chromatography, electrophoresis, and radiochemistry. Application of the test method for screening the ACE inhibitors activity and investigation of active compounds from natural products can be done easily with this method, it is very helpful in research because the results obtained are simple, accurate, and rapid.

**Key words:** Angiotensin-converting enzyme inhibitory activity, angiotensin-converting enzyme, herbal plants, *in vitro* assay method, Renin-angiotensin-aldosterone system

## INTRODUCTION

Initially, the screening for antihypertensive effect in the drugs discovery from natural products mainly used empirically been done over the years and have used several experiments on animal models.<sup>[1-3]</sup> Studies in the drug discovery, especially as an antihypertensive has developed rapidly since the discovery of the angiotensin-converting enzyme (ACE). The ACE converts angiotensin decapeptide inactive, into active octapeptide angiotensin II in the kidneys, especially in the renin-angiotensin-aldosterone system.<sup>[4-8]</sup> The activity of ACE inhibitory by *in vitro* has become an effective assay method in the drugs discovery as antihypertensive. This has been demonstrated by comparing the assay method of seven kinds medicines (captopril, enalapril, zofenopril, ramipril, fosinopril, lisinopril, and SQ 29852) as the ACE inhibitor.<sup>[9]</sup> However, in studies, the activity of ACE inhibitory for a positive control is more widely used a captopril because the drug is most widely used as antihypertensive and heart failure, and also have a free radical scavenger activity are highly relevant as an ACE inhibitor.<sup>[10]</sup>

### Correspondence:

Dr. Abdul Mun'im,  
Faculty of Pharmacy, Universitas Indonesia,  
Building A, 3<sup>rd</sup> Floor, Rumpun Ilmu Kesehatan,  
Kampus UI Depok, West Java 16424, Indonesia.  
E-mail: munimabdoel@gmail.com

### Access this article online

Quick Response Code: 	Website: www.phcogrev.com
	DOI: 10.4103/phrev.phrev_45_16

In modern medicine, the drug discovery has become more specific and focus on particular target objectives. The identification of receptor or enzyme as a molecular target which has an important role in the disease regulation and then performs searches the ligand or substrate or inhibitor of a specific target is the reason behind this approach. The discovery of a new drugs mainly from natural materials are directly aimed at the molecular target (receptor or enzyme) is more effective and efficient than conventional methods using animal model experiments performed with the treatment and observation in general, and require treatment and observation are more complicated if performed on specific targets (e.g. receptors or enzymes), as well as the type of the test sample to be used.<sup>[2,9,11-13]</sup> Considering the potential of natural resources are abundant so that the necessary a special strategy conduct research one of which is an assay method of the ACE inhibitors activity *in vitro*. The present studies review aims to highlight the discover history, assay methods, and application in natural products drugs discovery of ACE inhibitory activity assay.

## MATERIALS AND METHODS

This paper reviews about the ACE Inhibitory Activity Assay from comprehensive literature. The literature was searched between August and October 2016 from the electronic databases including PubMed, Scopus, ScienceDirect, and Google Scholar.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

**Cite this article as:** Ahmad I, Yanuar A, Mulia K, Mun'im A. Review of angiotensin-converting enzyme inhibitory assay: Rapid method in drug discovery of herbal plants. Phcog Rev 2017;11:1-7.

## RESULTS AND DISCUSSION

### The discover history of *in vitro* angiotensin-converting enzyme inhibitory assay methods

The assay method of ACE activity *in vitro* was begun in 1954–1957, when Skeggs *et al.* succeeded in isolating and purifying the enzyme which can hydrolyze decapeptide angiotensin I, then release vasopressive octapeptide angiotensin II and histidine-leucine dipeptide inactive or commonly referred to as “converting enzyme” from the horses plasma.<sup>[14]</sup> From the results of the discovery, the action mechanism of this enzyme can be determined [Figure 1].<sup>[15]</sup> However, at that time, there has been no progress on the development of *in vitro* assay methods.

About eleven years later (1968–1969) with the discovery of radiometric assay using the labeled angiotensin I substrate, wherein the release of radioactive histidine-leucine which serves as an enzymatic activity index<sup>[16]</sup> and further developed methods for chemical assay of the ACE, where the enzymatic reaction product based on the determination of histidine-leucine with fluorometric method on different substrates.<sup>[17]</sup> In 1970–1971, Cushman and Cheung managed to find a spectrophotometric assay method for measuring the amount of ACE to produce hippuric acid (HA) from hippuryl-histidyl-leucine (HHL) as substrate.<sup>[18]</sup>

Carmel and Yaron (1977–1978) developed a measurement method of the ACE inhibitory activity using an *o*-aminobenzoylglycine-*p*-nitrophenylalanylproline as a substrate and then hydrolyzed into *o*-aminobenzoylglycyl.<sup>[19]</sup> At the same time, Hayakari *et al.*, developed assay methods of the ACE inhibitory activity in a spectrophotometry manner using HHL as substrate and a colorimetric reagent of HA namely 2,4,6-trichloro-*s*-triazine (TT).<sup>[20]</sup>

In 1979–1991, some research reported the usage of a substrate other than HHL for assay method of ACE inhibitory activity such as Holmqvist *et al.* using tripeptide furanacryloyl (FA-PGG) as a substrate,<sup>[21]</sup> Baudin *et al.*, using benzoyl-[1-14C] glycyl-L-histidyl-L-leucine as a substrate,<sup>[22]</sup> and the usage a substrate of chromophore- and fluorophore-labelled tripeptide dansyltriglycine<sup>[23]</sup> by Elbl and Wagner. Since 1993, Doig and Smiley have developed method of ACE inhibitory activity assay using a




Figure 1: Scheme of angiotensin II formation by angiotensin-converting enzyme

shielded hydrophobic phase (SHP) column for high-performance liquid chromatography (HPLC) instrument and HHL substrate.<sup>[24]</sup> Nakamura *et al.* have performed purification and characterization of ACE inhibitory using HPLC instrument.<sup>[25]</sup>

Initially, all methods are constantly being developed is only used against the pure compound and has not been in use on samples containing multi-compound as in plant extracts. However, Hansen *et al.* began to apply this method on the plant extracts.<sup>[26]</sup> In 2007, Lam *et al.*, managed to synthesize and use a substrate 3-hydroxybutyrylglycyl-glycyl-glycine (3HB-GGG) in the assay method of the ACE inhibitory activity,<sup>[27]</sup> and this method is also applicable to samples of plant extracts.

### *In vitro* angiotensin converting enzyme assay methods

Several assay method of ACE inhibitory activity can be used to detect the activity of ACE inhibition from drugs or plant extract. Each method is distinguished by the use of substrates and measurement methods of enzymatic reaction products or separation of the substrate with the products. In addition, the author provides the names of each method based on the inventor names, ACE inhibitory activity test method is divided into several methods as follows:

#### Cushman and Cheung method

Cushman and Cheung (1970–1971) have developed the assay method of the activity of ACE inhibitors using a substrate hippuryl-histidyl-leucine (HHL), the ACE will hydrolyze HHL into HA [Figure 2].<sup>[28]</sup> The HA was measured at a wavelength of 228 nm to describe the ACE activity using a ultraviolet-visible (UV-Vis) spectrophotometer instrument. When there is an ACE inhibitor, the concentration of HA formed will be reduced.<sup>[18]</sup>

The success of Cushman and Cheung's method of assay the ACE inhibitors activity still depends on the ability to separate HA formed from the HHL substrate. Therefore, the another approach is required to facilitate the use of these methods optimally, among other: (a) The addition of TT in the sample mixture containing HA, a reaction between TT with HA formed should be measured at a wavelength of 382 nm.<sup>[20]</sup> (b) The usage of 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) to stop the reaction and Cyanuric chloride in 1,4-dioxane as a color reagent measured in photometry at a wavelength of 504 nm.<sup>[29]</sup> (c) The addition of *o*-Phthaldialdehyde which then react with the substrate hydrolysis histidine-leucine is measured by a fluoro-colorimeter instrument at a wavelength of 495 nm emission and 365 nm excitation.<sup>[30]</sup> (d) The use of benzene-sulfonyl-chloride (BSC) as a color reagent in the presence of




Figure 2: HHL hydrolysis by angiotensin-converting enzyme

quinoline<sup>[31]</sup> (referred to as BSC-visible spectrophotometry method) and then modified using a microtiter plate reader.<sup>[32]</sup>

Besides using the UV-Vis spectrophotometer, can also use the high HPLC, however, require more modifications such as the use of SHP HPLC to separate HHL and HA, which can be injected directly for the measurement of the HA absorbance using a spectrophotometer wavelength at 254 nm with a 8.0 mL flow-cell.<sup>[24]</sup> Furthermore, Mehanna and Dowling, were able to develop a more simple HPLC method,<sup>[33]</sup> which further developed by using reverse-phase-HPLC (RP-HPLC) and HEPES as a buffer.<sup>[34]</sup>

### Holmquist method

This method was first introduced by Holmquist *et al.* in 1979 using FA-PGG as a substrate. This method is based on the absorption spectrum blue shift that occurred on the substrate hydrolysis produces dipeptide and furanacryloyl-blocked amino acid.<sup>[21]</sup> This was confirmed by Lundberg *et al.*, to perform measurements of the ACE inhibitory activity in serum using an FA-PGG substrate.<sup>[35]</sup> This method has been conducted optimization, validation, and modifications of the FA-PGG usage as a substrate for screening bioactive peptides.<sup>[36,37]</sup> The FA-PGG substrate is hydrolyzed into a dipeptide (glycylglycine) and furanacryloyl-phenylalanine by ACE and measured at a wavelength of 328 nm and 352 nm,<sup>[37,38]</sup> the addition of ethylenediaminetetraacetic acid (EDTA) aimed to stopping the enzymatic reaction<sup>[37]</sup> and other modifications using a microplate reader.<sup>[38]</sup> Lahogue *et al.* has reported the application of HPLC-UV method for the determination of the activity of ACE inhibitors.<sup>[39]</sup>

### Elbl and Wagner method

This method was developed by Elbl and Wagner in 1991 with the usage of the chromophore- and fluorophore-labeled tripeptide dansyl triglycine as substrate, which is cleaved by the ACE into dansylglycine and diglycine [Figure 3].<sup>[23]</sup> Furthermore, this method modified by using a microtiter plate which is injected directly into the UV detection.<sup>[26,40]</sup> Duncan *et al.* have been applying this method for screening the activity of ACE inhibitors from Zulu medicinal plants.<sup>[41]</sup>

Lacaille-Dubois *et al.* modify Elbl and Wagner method in which the hippuric-glycine-glycine substrate is cleaved by ACE and react with trinitrobenzenesulfonate acid to form 2,4,6-trinitrophenyl-glycine-glycine were separated by RP-HPLC.<sup>[42,43]</sup> In addition, it also has been validated a colorimetric assay for screening ACE inhibitors from plant extracts *in vitro* by Serra *et al.*<sup>[44]</sup>




Figure 3: Dansyltriglycine hydrolysis by angiotensin-converting enzyme

### Baudin method

Baudin *et al.* have developed a method with radiometric technology to measure ACE activity in urine and using a substrate that has been labeled namely benzoyl-[1-14C] glycyl-L-histidyl-L-leucine. The substrate will be cleaved at the glycine-histidine bonding and form benzoyl-[1-14C] glycine. This test method can be used in the study of ACE in the urine as a marker if there is kidney damage.<sup>[22]</sup>

### Carmel and Yaron method

The Carmel and Yaron methods began to be discovered and developed in 1978, an assay method of the ACE inhibitory activity using *o*-amino benzoyl glycyl-*p*-nitrophenylalanilproline as a substrate then hydrolyzed into *o*-amino benzoyl glycyl as a compound fluorescence, for discontinuous enzymatic reaction can be used EDTA.<sup>[19]</sup> In 2006, Sentandreu and Toldrá using this method with slight modifications were measured using a multi-scanning microplate fluorometer. The advantage of this method is the usage capacity of large sample and in a short time.<sup>[45]</sup> However, this method is not widely used.

### Lam method

Lam *et al.*, begun to study in 2007 managed to find a new substrate is 3HB-GGG for assay method of the activity of ACE. The 3HB-GGG is cleaved into Gly-Gly-Gly amino acid and 3-hydroxybutyric acid (3HB) by the ACE. Then 3HB measured using F-kit. This method is more sensitive, rapid, accurate, and suitable for conventional methods.<sup>[27]</sup> Then, Lam *et al.* carried out development using a Water-Soluble Tetrazolium Salt (WST1) to detect 3-hydroxybutyrate formed.<sup>[46]</sup> The mechanism of the assay method of the ACE inhibitor activity using ACE kit-WST1 described in Figure 4 and this method using flow injection analysis to detect directly in a rapid, simple, and accurate.<sup>[47]</sup> This enzyme has been created in the form of a kit which has been patented with the name the ACE kit-WST1. The LAM method with the 3HB-GGG substrate using the ACE kit-WST can also be applied using microplate ELISA reader.<sup>[48-50]</sup>

### Applied of the angiotensin converting enzyme inhibitory assay method in drug discovery of natural products

The screening of antihypertensive activity on natural products, especially herbs used empirically as antihypertensive has been conducted over the years. The application of the assay method of the activity of ACE inhibitory for the activity screening in the natural product research is very helpful because the results obtained in a rapid, accurate, and simple. In general, the methods described above have been conducted optimization for the type and the materials concentration used so that the reaction mechanism of the enzyme occurs the same as in the actual state in the body,<sup>[29,36]</sup> this method also was performed standardization<sup>[30]</sup>




Figure 4: The principle of the assay method of angiotensin-converting enzyme inhibitors activity using angiotensin-converting enzyme kit-Water-Soluble Tetrazolium Salt 1

**Table 1:** The results data of angiotensin-converting enzyme inhibitors activity *in vitro* from plants

Species of plant	Family	Part of plant	ACE-I activity (%)	References
Species from India				[26]
<i>Abutilon indicum</i> G. Don	Malvaceae	Root	18	
<i>Achyranthes aspera</i> L.	Amaranthaceae	Arial parts	19	
<i>Boerhavia diffusa</i> L.	Nyctaginaceae	Root	40	
<i>Cardiospermum halicacabum</i> L.	Sapindaceae	Arial parts	8	
<i>Centella asiatica</i> L.	Apiaceae	Arial parts	73	
<i>Desmodium gangeticum</i> DC.	Fabaceae	Root	27	
<i>Entada pursaetha</i> DC.	Fabaceae	Seed	34	
<i>Merremia tridentata</i> Dennst.	Convolvulaceae	Arial parts	30	
<i>Pseudarthria viscida</i> W and Arn.	Fabaceae	Root	71	
<i>Sida acuta</i> Burm.	Malvaceae	Root	32	
<i>Sida cordifolia</i> L.	Malvaceae	Root	38	
<i>Sida retusa</i> L.	Malvaceae	Root	41	
<i>Triumfetta rhomboidea</i> Jacq.	Tiliaceae	Root	61	
Species from South Africa				
<i>Pseudarthria hookeri</i> W and Am.	Fabaceae	Root	90	
Species from China				
<i>Aristolochia manshuriensis</i> Kom	Aristolochiaceae	Seed	36	
<i>Crataegus pinnatifida</i> Bunge	Rosaceae	Fruit	7	
<i>Desmodium styracin</i> Mer.	Fabaceae	Leaf, seed	39	
<i>Houttuynia cordata</i> Thunb.	Saururaceae	Whole plant	23	
<i>Plantago asiatica</i> L.	Plantaginaceae	Whole plant	35	
<i>Pueraria lobata</i> Ohwi	Fabaceae	Root	38	
<i>Pyrrosia lingua</i> Farwell	Polypodiaceae	Whole plant	24	
<i>Sedum sarmentosum</i> Bunge	Crassulaceae	Whole plant	24	
<i>Uncaria rhynchophylla</i> Jacks.	Rubiaceae	Root	59	
Species from Chile				
<i>Aristolochia chilensis</i> DC.	Elaeocarpaceae	Bark	36	
<i>Escallonia myrtilloides</i> L.	Grossulariaceae	Arial part	15	
<i>Fuchsia magellanica</i> L.	Onagraceae	Arial part	47	
<i>Geranium core-core</i> L.	Geranaceae	Arial part	33	
<i>Gunnera tinctoria</i> L.	Gunneraceae	Leaf	57	
<i>Hexachlamys edulis</i> Berg.	Myrtaceae	Leaf	91	
<i>Quichamalium chinensis</i> Molina	Santalaceae	Arial part	43	
<i>Schinus latifolius</i> L. Schinus	Anacardiaceae	Bark	74	
<i>Adenopodia spicata</i> (E. Mey.) Pres	Fabaceae	Leaf, root	97,8	[41]
<i>Adiatum capillus-veneris</i> L.	Pteridaceae	Leaf	13	
<i>Agapanthus africanus</i> (L.)	Alliaceae	Leaf, root	63	
<i>Agave americana</i> L.	Asparagaceae	Leaf	82	
<i>Catha edulis</i> (Vahl) Forssk.	Celastraceae	Leaf	82	
<i>Cannabis sativa</i> L.	Cannabaceae	Leaf, root	18	
<i>Clausena anisata</i> (Wild.) Hook.	Rutaceae	Leaf	54	
<i>Dietes iridioides</i> (L.) Sweet	Iridaceae	Leaf, root	80	
<i>Drimia elata</i> Jacq.	Asparagaceae	Leaf, bulbs	16	
<i>Drimia robusta</i> Bak.	Asparagaceae	Leaf, bulbs	23	
<i>Dombeya rotundifolia</i> (Hochst.)	Pentapetaceae	Leaf, bark	83	
<i>Ekebergia capensis</i> Sparrm	Meliaceae	Leaf	37	
<i>Hypoxis colchicifolia</i> Bak.	Hypoxidaceae	Leaf, root	37	
<i>Mesembryanthemum</i> sp.	Aizoaceae	Leaf, stem	90	
<i>Protorhus longifolia</i> (Bernh)	Anacardiaceae	Leaf	77	
<i>Rhus shirindensis</i> Bak. F.	Anacardiaceae	Leaf, bark	85	
<i>Sclerocarya birrea</i> (A. Rich.)	Anacardiaceae	Leaf	68	
<i>Stangeria eriopus</i> (Kunze) Baill.	Stangeriaceae	Leaf	55	
<i>Tulbaghia tiolacea</i> Harv.	Amaryllidaceae	Leaf, root	72	
<i>Turraea floribunda</i> Hochst.	Meliaceae	leaf	45	
<i>Cecropia glaziovii</i> Sneathlage	Cecropiaceae	Stem	20	[42]
<i>Persea americana</i> Miller	Lauraceae	Stem	31	
<i>Croton antisiphiliticus</i> Mart.	Euphorbiaceae	Leaf	32	
<i>Combretum fruticosum</i> Stuntz	Combretaceae	Leaf	38	
<i>Buddleja stachyoides</i>	Loganiaceae	Leaf	31	
<i>Ouratea semiserrata</i> .	Ochnaceae	Stem	68	
<i>Tripogandra elata</i> D. R. Hunt	Commelinaceae	Leaf	32	
<i>Commelina diffusa</i> Burm. F	Commelinaceae	Leaf	45	
<i>Tradescantia zebrina</i> Hort.	Commelinaceae	Leaf	35	
<i>Hymenaea courbaril</i> L.	Leguminosae	Resin	10	
<i>Maytenus ilicifolia</i> Mart.	Celastraceae	Leaf	19	

Contd...

Table 1: Contd...

Species of plant	Family	Part of plant	ACE-I activity (%)	References
<i>Xylopia frutescens</i> Aubl.	Annonaceae	Seed	3	
<i>Cuphea carthagenensis</i> (Jacq.)	Lythraceae	Leaf	50	
<i>Mansoa hirsuta</i> DC.	Bignoniaceae	Leaf	54	
<i>Tabebuia serratifolia</i> (Vahl)	Bignoniaceae	Stem bark	19	
<i>Clytostoma ramentaceum</i> Bur.	Bignoniaceae	leaf	16	
<i>Musanga cecropioides</i> R.	Urticaceae	Leaf	100	[43]
<i>Crataegus oxyacantha</i>	Rosaceae	Leaf	<40	
<i>Cecropia hololeuca</i>	Urticaceae	Leaf	40	
<i>Cecropia pachystachya</i> Trec.	Urticaceae	Leaf	40-60	
<i>Cecropia glaziovii</i> Sneth.	Urticaceae	Leaf	60-100	
Species of plant	Family	Part of plant	ACE-I activity (µM)	References
<i>Olea europaea</i>	Oleaceae	Isolated	26	[40]
<i>Olea lancea</i>	Oleaceae	Isolated	0.33	
<i>Allium sativum</i> L.	Liliaceae	Isolated	3.74	[52]
Species of plant	Family	Part of plant	ACE-I activity (µg/ml)	References
<i>Panax ginseng</i>	Araliaceae	Radix	5	[53]
<i>Marrubium radiatum</i>	Lamiaceae	Whole plants	72.8	[54]
<i>Calamintha organifolia</i>	Lamiaceae	Whole plants	106.2	
<i>Hyssopus officinalis</i>	Lamiaceae	Whole plants	52.0	
<i>Asperula glomerata</i>	Rubiaceae	Whole plants	165.6	
<i>Satureja thymbra</i>	Lamiaceae	Whole plants	289.8	
<i>Salvia acetabulosa</i>	Lamiaceae	Whole plants	52.7	
<i>Periploca laevigata</i>		Root	30	[55]
Species of plant	Family	Part of plant	ACE-I activity (µg/ml)	References
<i>Erythroxylum gonocladum</i> (Mart.)	Erythroxylaceae	Arial part	4.53	[56]
<i>Scurulla artopurpurea</i>	Loranthaceae	Herbs	322	[57]
<i>Catharanthus roseus</i>	Apocynaceae	Leaf	402	
<i>Swietenia mahogany</i>	Meliaceae	Seed	774	
<i>Persea americana</i>	Lauraceae	Seed	476	
<i>Oxalis corniculata</i>	Oxalidaceae	Leaf	325	
<i>Phalleria marcocarpa</i>	Thymelaeaceae	Leaf, fruit	102	
<i>Gynura procumbens</i>	Asteraceae	Leaf	227	
<i>Melia azedarach</i>	Meliaceae	Leaf	483	
<i>Hibiscus rosasinensis</i>	Malvaceae	Leaf	271	
Species of plant	Family	Part of plant	ACE-I activity (mg/ml)	References
<i>Tamarix hohenackeri</i> Bunge	Tamaricaceae	Arial parts	100	[58]
		Isolated	20	
<i>Apium graveolens</i>	Apiaceae	Whole plants	1.7	[59]
<i>Solanum torvum</i>	Solanaceae	Fruits	1.2	[60]
Species of plant	Family	Part of plant	ACE-I activity (µg/ml)	References
<i>Mucuna pruriens</i>	Fabaceae	Seed	38.44	[61]
<i>Sesamum indicum</i> L.	Pedaliaceae	Seed	30.16	[62]
<i>Peperomia pellucida</i> (L.) Kunth	Piperaceae	Herb	7.17	
<i>Nasturtium officinale</i>	Brassicaceae	Herb	15.40	
<i>Peperomia pellucida</i> (L.) Kunth	Piperaceae	Ethyl acetate extract herb	3.34	[63]
		Isolated	7.72	

ACE-I: Angiotensin-converting enzyme inhibitory

and validation,<sup>[36]</sup> between each method as well as with conventional methods performed *in vivo* method. Selection of assay methods based on the substrate type depending on the availability of instruments that can be used to measure the inhibitory activity of ACE from the samples.<sup>[51]</sup> Some herbs that have been carried out the screening of activity using the *in vitro* assay method of ACE inhibitors activity as shown in Table 1. Based on the screening results of ACE inhibitors activity from various data sources [Table 1] show that the research of new drug discovery from natural materials is still very limited when compared with the availability of abundant natural resources around us.

The new drug discovery from natural materials are not only limited to the screening phase the activity but till the determining phase of compounds

is responsible as ACE inhibitors. The search of active compound can be performed easily with this method, in general, can be started from screening the activity of the extract, fractions, and isolates. Even for the isolates that have been characterized and determined the structure can be more optimal if combined with the *in silico* assay method<sup>[13,64]</sup> to obtain a prediction overview of the action mechanism and the group most responsible of the compounds studied. Several studies based on the empirical usage of natural products such as the celery plant (*Apium graveolens*) as antihypertensive and isolation of the junipediol A 8-OβD-glucoside (1-βD-glucosyloxy-2- (3-methoxy-4-hydroxyphenyl) -propane -1,3-diol) compound has the strongest activity as an ACE inhibitor compared to another known compound (such as, 11,21-dioxo-3

b, 15a, 24-trihydroxyurs-12-ene-24-O- $\beta$ -D-glucopyranoside, chrysoeriol-7-O- $\beta$ -D-apiosylglucoside, apiin, icaraside D<sub>2</sub>, luteolin-7-O- $\beta$ -D-glucoside, apigenin-7-O- $\beta$ -D-glucoside, roseoside, and isofraxidin- $\beta$ -D-glucoside).<sup>[59]</sup> Garlic (*Allium sativum*) has been known to dipeptide compounds that play a role as an ACE inhibitor, namely Gly-Phe, Ser-Phe, Gly-Tyr, Ser-Tyr, Asn-Phe, Asn-Tyr, and Phe-Tyr, with the IC<sub>50</sub> were 277.9, 130.2, 72.1, 66.3, 46.3, 32.6, and 3.74  $\mu$ M,<sup>[52]</sup> respectively. The (E)-2,3-dihydroxycyclopentyl-3-(3',4'-dihydroxyphenyl) acrylate compound was isolated from wild Eggplant (*Solanum torvum*) (IC<sub>50</sub> of 778 mg/mL).<sup>[60]</sup> *Suaeda physophora* Pall has isolated a quinoline alkaloids (Suaeda) and flavonoid compounds that also have a potential of ACE inhibitors activity.<sup>[65]</sup> The astibin (with IC<sub>50</sub> of 4.53  $\mu$ g/mL) were isolated from *Erythroxylum gonocladum* plant.<sup>[56]</sup> The quercetin isolated from *Peperomia pellucida* L. Kunth herbs with IC<sub>50</sub> of 7.72  $\mu$ g/mL.<sup>[63]</sup> By knowing the active compounds that play a role as an ACE inhibitor, the researchers of natural products are more focused on enrichment, dereplication, and isolation the active compound that can be applied in the *in vivo* assay method and further development.

## CONCLUSION

By the discovery of the various assay methods of ACE inhibitors activity. This method is very applicable for testing on natural products so that the researchers can perform quickly, accurately, and simple, so they can more focus on the development of extraction, fractionation, and isolation methods for the enrichment and dereplication of the active compounds. The selection of the assay methods for determining the inhibitory activity of ACE is only based on subject to availability of the substrate and the measurement instruments for example spectrophotometric, photo fluorometer, HPLC, and others.

## Acknowledgement

The authors are thankful to Ministry of Research, Technology and Higher Education, Republic of Indonesia and Directorate of Research and Community Engagement (DRPM), Universitas Indonesia for financially supporting the study via a grant "Hibah PUPT 2015" and "PITTA 2016".

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- López-Fandiño R, Otte J, Camp JV. Physiological, chemical and technological aspects of milk-protein-derived peptides with antihypertensive and ACE-inhibitory activity. *Int Dairy J* 2006;16:1277-93.
- Fujita M, Ohnishi K, Takaoka S, Ogasawara K, Fukuyama R, Nakamura H. Antihypertensive effects of continuous oral administration of nattokinase and its fragments in spontaneously hypertensive rats. *Biol Pharm Bull* 2011;34:1696-701.
- Lingbeck JM, O'Bryan CA, Martin EM, Adams JP, Crandall PG. Sweetgum: An ancient source of beneficial compounds with modern benefits. *Pharmacogn Rev* 2015;9:1-11.
- Skeggs LT Jr., Marsh WH, Kahn JR, Shumway NP. The existence of two forms of hypertension. *J Exp Med* 1954;99:275-82.
- Skeggs LT Jr., Kahn JR, Shumway NP. The preparation and function of the hypertensin-converting enzyme. *J Exp Med* 1956;103:295-9.
- Gude D. How full is our antihypertensives pipeline? *J Pharmacol Pharmacother* 2012;3:7-11.
- Tabassum N, Ahmad F. Role of natural herbs in the treatment of hypertension. *Pharmacogn Rev* 2011;5:30-40.
- Sayari N, Saidi N, Sila A, Ellouz-Chaabouni S, Bougataf A. Chemical composition, angiotensin I-converting enzyme (ACE) inhibitory, antioxidant and antimicrobial activities of *Ononis natrix* leaves extracts. *Free Radic Antioxid* 2016;6:23-33.
- Cushman DW, Wang FL, Fung WC, Grover GJ, Harvey CM, Scales RJ, *et al.* Comparisons *in vitro*, *ex vivo*, and *in vivo* of the actions of seven structurally diverse inhibitors of angiotensin converting enzyme (ACE). *Br J Clin Pharmacol* 1989;28 Suppl 2:115S-30S.
- Chopra M, Scott N, McMurray J, McLay J, Bridges A, Smith WE, *et al.* Captopril: A free radical scavenger. *Br J Clin Pharmacol* 1989;27:396-9.
- Fuglsang A, Nilsson D, Nyborg NC. Characterization of new milk-derived inhibitors of angiotensin converting enzyme *in vitro* and *in vivo*. *J Enzyme Inhib Med Chem* 2003;18:407-12.
- Dost T, Kafkas S, Gokalp F, Karul A, Birincioglu M. Effects of angiotensin converting enzyme inhibition on adiponectin levels and lipid profile in the ovariectomized-aged rats. *J Pharmacol Pharmacother* 2014;5:21-6.
- Muhammad SA, Fatima N. *In silico* analysis and molecular docking studies of potential angiotensin-converting enzyme inhibitor using quercetin glycosides. *Pharmacogn Mag* 2015;11 Suppl 1:S123-6.
- Skeggs LT Jr., Kahn JR, Lentz K, Shumway NP. The preparation, purification, and amino acid sequence of a polypeptide renin substrate. *J Exp Med* 1957;106:439-53.
- Guang C, Phillips RD, Jiang B, Milani F. Three key proteases – Angiotensin-I-converting enzyme (ACE), ACE2 and renin – Within and beyond the renin-angiotensin system. *Arch Cardiovasc Dis* 2012;105:373-85.
- Huggins CG, Thampi NS. A simple method for the determination of angiotensin I converting enzyme. *Life Sci* 1968;7:633-9.
- Piquilloud Y, Reinharz A, Roth M. Studies on the angiotensin converting enzyme with different substrates. *Biochim Biophys Acta* 1970;206:136-42.
- Cushman DW, Cheung HS. Spectrophotometric assay and properties of the angiotensin-converting enzyme of rabbit lung. *Biochem Pharmacol* 1971;20:1637-48.
- Carmel A, Yaron A. An intramolecularly quenched fluorescent tripeptide as a fluorogenic substrate of angiotensin-I-converting enzyme and of bacterial dipeptidyl carboxypeptidase. *Eur J Biochem* 1978;87:265-73.
- Hayakari M, Kondo Y, Izumi H. A rapid and simple spectrophotometric assay of angiotensin-converting enzyme. *Anal Biochem* 1978;84:361-9.
- Holmquist B, Bünning P, Riordan JF. A continuous spectrophotometric assay for angiotensin converting enzyme. *Anal Biochem* 1979;95:540-8.
- Baudin B, Bénéteau-Burnat B, Baumann FC, Giboudeau J. A reliable radiometric assay for the determination of angiotensin I-converting enzyme activity in urine. *J Clin Chem Clin Biochem* 1990;28:857-61.
- Elbl G, Wagner H. A new method for the *in vitro* screening of inhibitors of angiotensin-converting enzyme (ACE), using the chromophore- and fluorophore-labelled substrate, dansyltriglycine. *Planta Med* 1991;57:137-41.
- Doig MT, Smiley JW. Direct injection assay of angiotensin-converting enzyme by high-performance liquid chromatography using a shielded hydrophobic phase column. *J Chromatogr B Biomed Sci Appl* 1993;613:145-9.
- Nakamura Y, Yamamoto N, Sakai K, Okubo A, Yamazaki S, Takano T. Purification and characterization of angiotensin I-converting enzyme inhibitors from sour milk. *J Dairy Sci* 1995;78:777-83.
- Hansen K, Nyman U, Smitt UW, Adersen A, Gudiksen L, Rajasekharan S, *et al.* *In vitro* screening of traditional medicines for anti-hypertensive effect based on inhibition of the angiotensin converting enzyme (ACE). *J Ethnopharmacol* 1995;48:43-51.
- Lam le H, Shimamura T, Sakaguchi K, Noguchi K, Ishiyama M, Fujimura Y, *et al.* Assay of angiotensin I-converting enzyme-inhibiting activity based on the detection of 3-hydroxybutyric acid. *Anal Biochem* 2007;364:104-11.
- Thongnopnua P, Poeknapo C. High-performance liquid chromatographic determination of enalapril in human plasma by enzyme kinetic analytical method. *J Pharm Biomed Anal* 2005;37:763-9.
- Schnaith E, Beyrau R, Bückner B, Klein RM, Rick W. Optimized determination of angiotensin I-converting enzyme activity with hippuryl-L-histidyl-L-leucine as substrate. *Clin Chim Acta* 1994;227:145-58.
- Oliveira EM, Santos RA, Krieger JE. Standardization of a fluorimetric assay for the determination of tissue angiotensin-converting enzyme activity in rats. *Braz J Med Biol Res* 2000;33:755-64.
- Li GH, Liu H, Shi YH, Le GW. Direct spectrophotometric measurement of angiotensin I-converting enzyme inhibitory activity for screening bioactive peptides. *J Pharm Biomed Anal* 2005;37:219-24.
- Chen J, Wang Y, Ye R, Wu Y, Xia W. Comparison of analytical methods to assay inhibitors of angiotensin I-converting enzyme. *Food Chem* 2013;141:3329-34.

33. Mehanna AS, Dowling M. Liquid chromatographic determination of hippuric acid for the evaluation of ethacrynic acid as angiotensin converting enzyme inhibitor. *J Pharm Biomed Anal* 1999;19:967-73.
34. Sharifi N, Sourí E, Ziai SA, Amin G, Amini M, Amanlou M. Isolation, identification and molecular docking studies of a new isolated compound, from *Onopordon acanthium*: A novel angiotensin converting enzyme (ACE) inhibitor. *J Ethnopharmacol* 2013;148:934-9.
35. Lundberg PA, Lindstedt G, Andersson T, Branegård B, Lundquister G, Nyström E. Single reagent assay for angiotensin converting enzyme in serum. *J Clin Chem* 1984;30:163-8.
36. Vermeirssen V, Van Camp J, Verstraete W. Optimisation and validation of an angiotensin-converting enzyme inhibition assay for the screening of bioactive peptides. *J Biochem Biophys Methods* 2002;51:75-87.
37. Murray BA, Walsh DJ, FitzGerald RJ. Modification of the furanacryloyl-L-phenylalanyl-glycylglycine assay for determination of angiotensin-I-converting enzyme inhibitory activity. *J Biochem Biophys Methods* 2004;59:127-37.
38. Li Y, Zhou J, Huang K, Sun Y, Zeng X. Purification of a novel angiotensin I-converting enzyme (ACE) inhibitory peptide with an antihypertensive effect from loach (*Misgurnus anguillicaudatus*). *J Agric Food Chem* 2012;60:1320-5.
39. Lahogue V, Réhel K, Taupin L, Haras D, Alloume P. A HPLC-UV method for the determination of angiotensin I-converting enzyme (ACE) inhibitory activity. *Food Chem* 2010;118:870-5.
40. Hansen K, Adersen A, Christensen SB, Jensen SR, Nyman U, Smitt UW. Isolation of an angiotensin converting enzyme (ACE) inhibitor from *Olea europaea* and *Olea lancea*. *Phytomedicine* 1996;2:319-25.
41. Duncan AC, Jäger AK, van Staden J. Screening of Zulu medicinal plants for angiotensin converting enzyme (ACE) inhibitors. *J Ethnopharmacol* 1999;68:63-70.
42. Castro Braga F, Wagner H, Lombardi JA, de Oliveira AB. Screening the Brazilian flora for antihypertensive plant species for *in vitro* angiotensin-I-converting enzyme inhibiting activity. *Phytomedicine* 2000;7:245-50.
43. Lacaille-Dubois MA, Franck U, Wagner H. Search for potential angiotensin converting enzyme (ACE)-inhibitors from plants. *Phytomedicine* 2001;8:47-52.
44. Serra CP, Côrtes SF, Lombardi JA, Braga de Oliveira A, Braga FC. Validation of a colorimetric assay for the *in vitro* screening of inhibitors of angiotensin-converting enzyme (ACE) from plant extracts. *Phytomedicine* 2005;12:424-32.
45. Sentandreu MÁ, Toldrá F. A rapid, simple and sensitive fluorescence method for the assay of angiotensin-I converting enzyme. *Food Chem* 2006;97:546-54.
46. Lam le H, Shimamura T, Manabe S, Ishiyama M, Ukeda H. Assay of angiotensin I-converting enzyme-inhibiting activity based on the detection of 3-hydroxybutyrate with water-soluble tetrazolium salt. *Anal Sci* 2008;24:1057-60.
47. Lam le H, Shimamura T, Ishiyama M, Ukeda H. Flow injection analysis of angiotensin I-converting enzyme inhibitory activity with enzymatic reactors. *Talanta* 2009;79:1130-4.
48. Lau CC, Abdullah N, Shuib AS, Aminudin N. Novel angiotensin I-converting enzyme inhibitory peptides derived from edible mushroom *Agaricus bisporus* (J.E. Lange) Imbach identified by LC-MS/MS. *Food Chem* 2014;148:396-401.
49. Tran HB, Yamamoto A, Matsumoto S, Ito H, Igami K, Miyazaki T, *et al.* Hypotensive effects and angiotensin-converting enzyme inhibitory peptides of reishi (*Ganoderma lingzhi*) auto-digested extract. *Molecules* 2014;19:13473-85.
50. Li F, Yamaki K, Cheng Y, Fang Y. Assessment and separation of angiotensin I-converting enzyme inhibitory peptides in Chinese soypaste. *Int J Food Eng* 2015;11:301-5.
51. Shalaby SM, Zakora M, Otte J. Performance of two commonly used angiotensin-converting enzyme inhibition assays using FA-PGG and HHL as substrates. *J Dairy Res* 2006;73:178-86.
52. Suetsuna K. Isolation and characterization of angiotensin I-converting enzyme inhibitor dipeptides derived from *Allium sativum* L (garlic). *J Nutr Biochem* 1998;9:415-9.
53. Persson IA, Dong L, Persson K. Effect of Panax ginseng extract (G115) on angiotensin-converting enzyme (ACE) activity and nitric oxide (NO) production. *J Ethnopharmacol* 2006;105:321-5.
54. Loizzo MR, Saab AM, Tundis R, Menichini F, Bonesi M, Piccolo V, *et al.* *In vitro* inhibitory activities of plants used in Lebanon traditional medicine against angiotensin converting enzyme (ACE) and digestive enzymes related to diabetes. *J Ethnopharmacol* 2008;119:109-16.
55. Hajji M, Masmoudi O, Souissi N, Triki Y, Kammoun S, Nasri M. Chemical composition, angiotensin I-converting enzyme (ACE) inhibitory, antioxidant and antimicrobial activities of the essential oil from *Periploca laevigata* root barks. *Food Chem* 2010;121:724-31.
56. Lucas-Filho MD, Silva GC, Cortes SF, Mares-Guia TR, Perpétua Ferraz V, Serra CP, *et al.* ACE inhibition by astilbin isolated from *Erythroxylum gonocladum* (Mart.) O.E. Schulz. *Phytomedicine* 2010;17:383-7.
57. Rinayanti A, Radji M, Mun'im A, Suyatna FD. Screening angiotensin converting enzyme (ACE) inhibitor activity of antihypertensive medicinal plants from Indonesia. *Int J Pharm Teach Pract* 2013;4:527-32.
58. Xing Y, Liao J, Tang Y, Zhang P, Tan C, Ni H, *et al.* ACE and platelet aggregation inhibitors from *Tamarix hohenackeri* Bunge (host plant of Herba Cistanches) growing in Xinjiang. *Pharmacogn Mag* 2014;10:111-7.
59. Simaratanamongkol A, Umehara K, Noguchi H, Panichayupakaranant P. Identification of a new angiotensin-converting enzyme (ACE) inhibitor from Thai edible plants. *Food Chem* 2014;165:92-7.
60. Simaratanamongkol A, Umehara K, Niki H. Angiotensin-converting enzyme (ACE) inhibitory activity of *Solanum torvum* and isolation of a novel methyl salicylate glycoside. *J Funct Foods* 2014;11:557-62.
61. Chaudhary SK, De A, Bhadra S, Mukherjee PK. Angiotensin-converting enzyme (ACE) inhibitory potential of standardized *Mucuna pruriens* seed extract. *Pharm Biol* 2015;53:1614-20.
62. Saputri F, Mun'im A, Lukmanto D, Aisyah S, Rinandy J. Inhibition of angiotensin converting enzyme (ACE) activity by some Indonesia edible plants. *Int J Pharm Sci Res* 2015;6:1054-9.
63. Kurniawan A, Saputri F, Ahmad I, Mun'im A. Isolation of angiotensin converting enzyme (ACE) inhibitory activity quercetin from *Peperomia pellucida*. *Int J PharmTech Res* 2016;9:115-21.
64. Chang YW, Allii I. *In silico* assessment: Suggested homology of chickpea (*Cicer arietinum* L.) legumin and prediction of ACE-inhibitory peptides from chickpea proteins using BLAST and BIOPEP analyses. *Food Res Int* 2012;49:477-86.
65. Men R, Li N, Xing Y, Tang Y, Tan C, Meng F, *et al.* Chemical constituents and ACE inhibitory activity of desert plant *Suaeda physophora* Pall. *Acta Pharm Sin B* 2013;3:328-32.

# Review of Angiotensin-converting Enzyme Inhibitory Assay: Rapid Method in Drug Discovery of Herbal Plants

## ORIGINALITY REPORT

11%

SIMILARITY INDEX

5%

INTERNET SOURCES

9%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** Faiyaz Ahmed. "Radical scavenging and angiotensin converting enzyme inhibitory activities of standardized extracts of *Ficus racemosa* stem bark", *Phytotherapy Research*, 06/17/2010  
Publication 1%
- 2** [www.jove.com](http://www.jove.com)  
Internet Source 1%
- 3** Arunee Simaratanamongkol, Kaoru Umehara, Hiroshi Noguchi, Pharkphoom Panichayupakaranant. "Identification of a new angiotensin-converting enzyme (ACE) inhibitor from Thai edible plants", *Food Chemistry*, 2014  
Publication 1%
- 4** Submitted to Institute of Technology, Nirma University  
Student Paper <1%
- 5** [biomedpharmajournal.org](http://biomedpharmajournal.org)  
Internet Source <1%

6

Annika Lagemann, Andreas Dunkel, Thomas Hofmann. " Activity-Guided Discovery of ( )-Malic Acid 1'- - $\beta$ -Gentiobioside as an Angiotensin I-Converting Enzyme Inhibitor in Lettuce (*Lactuca sativa*) ", Journal of Agricultural and Food Chemistry, 2012

Publication

<1%

7

[www.aclso.saa.ars.usda.gov](http://www.aclso.saa.ars.usda.gov)

Internet Source

<1%

8

V. K. Jimsheena, Lalitha R. Gowda. "Colorimetric, High-Throughput Assay for Screening Angiotensin I-Converting Enzyme Inhibitors", Analytical Chemistry, 2009

Publication

<1%

9

Neena Baby, Kollencheri Puthenveetil Vinayan, Nivedita Pavithran, Arun Grace Roy. "A pragmatic study on efficacy, tolerability and long term acceptance of ketogenic diet therapy in 74 South Indian children with pharmaco-resistant epilepsy", Seizure, 2018

Publication

<1%

10

[studentsrepo.um.edu.my](http://studentsrepo.um.edu.my)

Internet Source

<1%

11

Progress in Experimental Cardiology, 1998.

Publication

<1%

12

Submitted to Tezpur University

<1%

13

[sutir.sut.ac.th:8080](http://sutir.sut.ac.th:8080)

Internet Source

<1%

14

Arunee Simaratanamongkol, Kaoru Umehara, Hiroki Niki, Hiroshi Noguchi, Pharkphoom Panichayupakaranant. "Angiotensin-converting enzyme (ACE) inhibitory activity of Solanum torvum and isolation of a novel methyl salicylate glycoside", Journal of Functional Foods, 2014

Publication

<1%

15

Chau, C.F.. "The development of regulations of Chinese herbal medicines for both medicinal and food uses", Trends in Food Science & Technology, 200606

Publication

<1%

16

Guang, Cuie, and Robert D. Phillips. "Plant Food-Derived Angiotensin I Converting Enzyme Inhibitory Peptides", Journal of Agricultural and Food Chemistry, 2009.

Publication

<1%

17

[www.nature.com](http://www.nature.com)

Internet Source

<1%

18

Ping Wu, Yiting Guo, Moses Kwaku Golly, Haile Ma, Ronghai He, Shilong Luo, Cheng Zhang, Liuping Zhang, Jinhua Zhu. " Feasibility study

<1%

on direct fermentation of soybean meal by under non-sterile conditions ", Journal of the Science of Food and Agriculture, 2018

Publication

19

Bruno Baudin, Jacqueline Giboudeau.  
"Carboxypeptidase A Hydrolyses Benzoylglycyl-Histidyl-Leucine but not Furylacryloyl-Phenylalanyl-Glycyl-Glycine, Two Usual Substrates for Angiotensin 1-Converting Enzyme", Enzyme and Protein, 1994

Publication

<1%

20

Submitted to Letterkenny Institute of Technology

Student Paper

<1%

21

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

<1%

22

Lam, L.H.. "Flow injection analysis of angiotensin I-converting enzyme inhibitory activity with enzymatic reactors", Talanta, 20090915

Publication

<1%

23

[www.scielo.br](http://www.scielo.br)

Internet Source

<1%

24

[repub.eur.nl](http://repub.eur.nl)

Internet Source

<1%

25

Ju-Chi Liu, Feng-Lin Hsu, Jen-Chen Tsai, Paul Chan, Jenny Ya-Hsin Liu, G.Neil Thomas, Brian

<1%

Tomlinson, Ming-Yu Lo, Jung-Yaw Lin.  
"Antihypertensive effects of tannins isolated  
from traditional Chinese herbs as non-specific  
inhibitors of angiotensin converting enzyme",  
Life Sciences, 2003

Publication

---

26

[repositorio.unicamp.br](http://repositorio.unicamp.br)

Internet Source

<1%

---

27

Submitted to Universiti Putra Malaysia

Student Paper

<1%

---

28

Hayakari, M.. "A rapid and simple  
spectrophotometric assay of angiotensin-  
converting enzyme", Analytical Biochemistry,  
197802

Publication

<1%

---

29

Fernanda A Ronchi. "N-domain angiotensin-  
converting enzyme isoform expression in tissues  
of Wistar and spontaneously hypertensive rats",  
Journal of Hypertension, 10/2005

Publication

<1%

---

30

Myeonghwa Cha. "Production and  
Characterization of a Soy Protein-Derived  
Angiotensin I-Converting Enzyme Inhibitory  
Hydrolysate", Journal of Medicinal Food,  
09/2005

Publication

<1%

---

32

Ramesh Gondru, Ratni Saini, Krishnaiah Vaarla, Sarika Singh, Narsimha Sirassu, Rajitha Bavantula, Anil K. Saxena. "Synthesis and Characterization of Chalcone-Pyridinium Hybrids as Potential Anti-Cancer and Anti-Microbial Agents", ChemistrySelect, 2018

Publication

&lt;1%

33

C.D. Collins, J.N.B. Bell. "Experimental studies on the deposition to crops of radioactive gases released from gas-cooled reactors —II. Hydrogen sulphide-35", Journal of Environmental Radioactivity, 1997

Publication

&lt;1%

34

Ana Silvia Suassuna Carneiro Lúcio, Jackson Roberto Guedes da Silva Almeida, Emídio Vasconcelos Leitão da-Cunha, Josean Fachine Tavares et al. "Alkaloids of the Annonaceae: Occurrence and a Compilation of Their Biological Activities", Elsevier BV, 2015

Publication

&lt;1%

35

Submitted to Monash University Sunway Campus Malaysia Sdn Bhd

Student Paper

&lt;1%

36

Yesmine Ben Henda, Anis Labidi, Ingrid Arnaudin, Nicolas Bridiau et al. "Measuring

&lt;1%

Angiotensin-I Converting Enzyme Inhibitory Activity by Micro Plate Assays: Comparison Using Marine Cryptides and Tentative Threshold Determinations with Captopril and Losartan", Journal of Agricultural and Food Chemistry, 2013

Publication

---

37

Lopez-Fandino, R.. "Physiological, chemical and technological aspects of milk-protein-derived peptides with antihypertensive and ACE-inhibitory activity", International Dairy Journal, 200611

Publication

---

38

Edible Medicinal and Non Medicinal Plants, 2015.

Publication

---

39

Le Hoang Lam, Tomoko Shimamura, Ken Sakaguchi, Katsuya Noguchi, Munetaka Ishiyama, Yume Fujimura, Hiroyuki Ukeda. "Assay of angiotensin I-converting enzyme-inhibiting activity based on the detection of 3-hydroxybutyric acid", Analytical Biochemistry, 2007

Publication

---

40

Submitted to University of Huddersfield

Student Paper

---

<1%

<1%

<1%

<1%

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On

# Review of Angiotensin-converting Enzyme Inhibitory Assay: Rapid Method in Drug Discovery of Herbal Plants

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---