

# Steroids and isoquercetin from *Lygodium microphyllum*

*by* Hadi Kuncoro

---

**Submission date:** 30-May-2019 11:21PM (UTC-0700)

**Submission ID:** 1138240549

**File name:** 2483\_pdf.pdf (511.31K)

**Word count:** 4911

**Character count:** 22241



## Steroids and isoquercetin from *Lygodium microphyllum*

Hadi Kuncoro<sup>1\*</sup>, Kindi Farabi<sup>2</sup>, Laode Rijai<sup>1</sup>

<sup>1</sup>Laboratory of Pharmaceuticals Research and Development, TROPICAL PHARMACA, Faculty of Pharmacy, Mulawarman University, Samarinda 75119, Kalimantan Timur, Indonesia. <sup>2</sup>Departement of Chemistry, Faculty of Mathematics and Natural Sciences, Padjadjaran University, Jatinangor 45363, Sumedang, Jawa Barat, Indonesia.

### ARTICLE INFO

#### Article history:

Received on: 19/01/2017

Accepted on: 06/04/2017

Available online: 30/11/2017

**Key words:** *Lygodium microphyllum*, stigma-5(6)-en-3 $\beta$ -ol, Stigmast-4-en-3-one, quercetin 3-O- $\beta$ -D-glucopyranoside, Lygodiaceae.

### ABSTRACT

Two steroids, stigma-5(6)-en-3 $\beta$ -ol or  $\beta$ -sitosterol (**1**) and Stigmast-4-en-3-one (**2**) and quercetin 3-O- $\beta$ -D-glucopyranoside or Isoquercetin (**3**) has been isolated from an extract of n-hexane and ethyl acetate from the leaves of plants *Lygodium microphyllum*. The chemical structures of compounds **1**, **2** and **3** identified based on spectroscopic data and comparison of spectral data obtained from the literature. The discovery of steroid compounds, stigma-5 (6) -en-3 $\beta$ -ol, Stigmast-4-en-3-one, and quercetin 3-O- $\beta$ -D-glucopyranoside in plants *L. microphyllum* first reported in this study

### INTRODUCTION

Fern has survived since Paleozoic era and can adapt to variety environmental changes (Wallace *et al*, 1991), thus fern has a lot of useful secondary metabolites, including flavonoids, steroids, alkaloids, phenols, triterpenoids, and various kinds of amino acids and fatty acids (Zeng-fu *et al*, 2008). One from thousands of species of ferns that have interesting pharmacological benefits is Lygodiaceae family. The only genus in this family is the genus *Lygodium* (Guo-gang *et al*, 2012). Generally, *Lygodium* genus is a group of ferns that spread and always propagate in other plants. Genus *Lygodium* are different from other kinds of ferns because it has roots that crawl on the

ground rhizomes and fleshy and can only live in the open because they like the sunlight. Some plants from genus *Lygodium* are invasive and have become a problem in a number of forest areas. There are the fast growing plant and lack of predator makes them dominate, displacing wildlife, threatens biodiversity, and enhance the human-animal conflict. One of the invasive species from genus *Lygodium* is *L. microphyllum*. Plants of this genus have a variety of properties that have been widely recognized, thus the utilization of plants from genus is quite expected. Some herbs *Lygodium* widely used by people one of them as traditional medicine as hepatitis medicine (Zheng and Xing, 2009). Back pain, rheumatism and treatment for kidney stones (Lee *et al*, 2008), expectorant, scabies, eczema and liver treatment (Upreti *et al*, 2009), diuretics, antiplasmodial, and treatment for lung and kidney (Upreti *et al*, 2009), laxative, headache and digestive disorders (Zheng dan Xing, 2009), Menstrual painkiller, contraception, bone fracture and hemorrhoid treatment (Cambie and Ash, 1994) and cough treatment (Karthik, 2011).

#### \* Corresponding Author

Hadi Kuncoro, Laboratory Of Pharmaceuticals Research and Development, TROPICAL PHARMACA, Faculty Of Pharmacy, Mulawarman University, Samarinda 75119, Kalimantan Timur, Indonesia. Email: [kuncoro\\_hadi82@yahoo.com](mailto:kuncoro_hadi82@yahoo.com)

Phytochemical Studies on the genus *Lygodium* been reported to contain a compound with a structure that is unique and diverse biological activities such as flavonoids (Zhang *et al.*, 2006), Glycoside phenolics (Ye *et al.*, 2007), naphthoquinone (Chen *et al.*, 2010), ecdysteroids (Zhu *et al.*, 2009), phenylpropanoid glycoside (Duan *et al.*, 2012).

*L. microphyllum* based on ethnobotany information and ethnopharmacological provides many benefits in the field of reproductive health as well as simple patterns make the plant is classified as herbs are easily available so that benefit of these plants is required primarily to determine the chemical constituents of this plant. In our ongoing research to find new compounds from Indonesia *Lygodium* plants, has been investigated *Lygodium* a chemical plant that grows in Borneo. In this communication, we will report the steroid compound, stigmast-5 (6) -en-3 $\beta$ -ol and stigmast-4-en-3-one from the n-hexane extract and quercetin 3-O- $\beta$ -D-glucopyranoside from the extract ethyl acetate plant *L. microphyllum*

## MATERIALS AND METHOD

Melting point measured on electrothermal melting point apparatus and not corrected. IR spectra were measured on Perkin-Elmer 1760X spectrophotometer, FT-IR on KBr. Mass spectra recorded with a mass spectrometer Water, Qtof HR-MS XEVO<sup>TM</sup>. <sup>1</sup>H and <sup>13</sup>C NMR spectra are obtained by JEOL NMR 500 MHz, JEOL NMRECZR 600 Mhz used TMS as an internal standard. Chromatographic separation is carried out on silica gel 60 (Merck), ODS (Fuji silycia). TLC Plate filled with silica gel GF<sub>254</sub> (Merck, 0.25 mm) and detection was obtained with the appearance of 10% H<sub>2</sub>SO<sub>4</sub> in ethanol followed by heating and under ultraviolet-visible light at wavelengths of 257 and 364 nm. Preparative MPLC using a Buchi Pump Controller C-610, Buchi Pump Modules C-605 with FLH-R10030B *SiliCycle* column-ISO04 *Siliasep*<sup>TM</sup>.

### Plant material

*L. microphyllum* plant material is obtained from forest areas in Samarinda, East Kalimantan in June 2014. The plant is identified by staff at the Faculty of Forestry, University of Mulawarman, Samarinda and sample specimens stored at the Faculty of Forestry, University of Mulawarman, Samarinda.

### Extraction and isolation

#### Compound isolation from n-hexane extract

Dried powder of *L. microphyllum* (3.54 kg) was extracted with methanol at room temperature. The ethanol extract obtained was concentrated at low pressure to produce methanol extract concentrated dark brown (526 g). The further concentrated methanol extract was dissolved in water (4: 1) and partitioned successively with n-hexane, ethyl acetate, and n-butanol. Evaporation of the solvent from the resulting extract each successive n-hexane extract (59 g), ethyl acetate (72 g) and n-butanol (54 g). Most of the n-hexane extract (50 g) were separated

by vacuum liquid chromatography (VLC) on silica gel G60 with n-hexane-ethyl acetate-methanol, 10% increasing polarity generated 25 fractions (H01-25). Fraction H3-11, combined (4 g) was separated by a column chromatography on silica gel (70-230 mesh) with the eluent n-hexane-ethyl acetate, 1% increased polarity generated 23 fractions (F01-23). F8 fraction separated using Medium performance liquid chromatography (MPLC) preparative with column *SiliCycle* FLH-R10030B-ISO04 *Siliasep*<sup>TM</sup> eluted by comparison isocratic solvent, chloroform: ethyl acetate (9: 1), the pump speed is 3 mL / min, accommodated every 30 seconds to 50 fractions and guided by TLC, fractions obtained 50 fractions divided into two namely D1 (1-5) and D2 (6-50), D2 subfraction separated using moderate performance liquid chromatography (MPLC) preparative with *SiliCycle* FLH-R10030B-ISO04 *Siliasep*<sup>TM</sup> column eluted by comparison isocratic solvent, toluene: chloroform (8: 2), the pump speed of 1.5 mL / min, accommodated every 30 seconds into 30 fractions and guided by TLC, subfraction D2C (15-30) are separated again using Medium Performance Liquid Chromatography (MPLC) preparative with *SiliCycle* FLH-R10030B-ISO04 *Siliasep*<sup>TM</sup> column eluted by comparison isocratic solvent, toluene: chloroform (8: 2), the pump speed of 1.2 mL / min, accommodated every 30 seconds into 30 fractions and guided by TLC, subfraction D2C2 (21-30) was purified using preparative thin layer chromatography eluted with a ratio of solvent, chloroform: ethyl acetate: acetic acid (9: 1: 0.5) and needle-shaped crystals obtained stigmast-4-en-3-one (5 mg). Fraction H10-16, combined (4.6 g) were separated by column chromatography on silica gel (70-230 mesh) with the eluent n-hexane-ethyl acetate, 5% increasing polarity generated 25 fractions (I01-25). I012-17 fraction combined (230 mg) were separated by preparative thin layer chromatography on silica gel GF<sub>254</sub> with chloroform eluent: ethyl acetate: acetic acid (9: 1: 0.5) to produce as much as 40.6 mg white solid that is subsequently crystallized with n-hexane-ethyl acetate produced white crystals colorless stigmast-5 (6) -en-3 $\beta$ -ol (38 mg).

#### Compound isolation from ethyl acetate extract

The ethyl acetate extract was separated using column chromatography open with ODS eluted with a gradient solvent ratio, water: Methanol: Ethyl Acetate, guided by TLC, produced 22 fractions (1-22). Fraction 5-7 (B) were merged and acquired 920 mg, then separated using column chromatography on silica G60 open eluted with a gradient ratio of solvent, ethyl acetate: methanol, guided by TLC and obtained 22 fractions (B1-22), fraction B2 (B2-B5) merged and acquired 420 mg, then separated using open column chromatography on silica G60 eluted with a gradient ratio of solvent, ethyl acetate: methanol, guided by TLC to obtain 22 fractions (B2.1-22), fraction B2.2 (B2B) as much as 190 mg open separated using column chromatography on silica G60 eluted with a gradient solvent ratio, chloroform: methanol, guided by TLC, and obtained 14 fractions (B2B.1-14), fractions to 9 (B2B.9) was further purified using preparative thin layer

chromatography eluted with a ratio of solvent, chloroform : methanol: acetic acid (7: 3: 0.5) to obtain yellow powder 3-O- $\beta$ -D-glucopyranoside 5.4 mg

## RESULT AND DISCUSSION

The methanol extract<sup>8</sup> of the dried powder *L. microphyllum* was concentrated and partitioned successively with n-hexane, ethyl acetate, and n-butanol. Part of n-hexane extract separated with various chromatographic techniques on the stationary phase silica gel and reverse phase to produce steroid compounds (1), (2) and the compound of Isoquercetin (3).




Fig. 1: Chemical structure of steroid (1) stigmast-5(6)-en-3 $\beta$ -ol or  $\beta$ -sitosterol.




Fig. 2: Chemical structure of steroid (2) Stigmast-4-en-3-one.




Fig. 3: Chemical structure of compound (3) 3-O- $\beta$ -D-glucopyranoside or isoquercetin.

### Stigmast-5(6)-en-3 $\beta$ -ol (1).

Crystal white needles; melting point 134-136 °C. IR (KBr)  $\nu_{\text{max}}$  3440, 2934, 2890, 1640, 1190  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (aseton-d<sub>6</sub>, 500 MHz), Table 1;  $^{13}\text{C}$  NMR (aseton-d<sub>6</sub>, 125 MHz), table 3, TOF MS (negatif ion mode) [m/z 413.0911, calculations for C<sub>28</sub>H<sub>50</sub>O m/z [414.0818].

$^1\text{H}$ NMR (aseton-d<sub>6</sub>, 500 MHz)  $\delta$  0.74 (3H, s), 0.84 (3H, d, 6.2) 0.86 (3H, d, 6.1), 0.88 (3H, t, 1.89), 0.95 (1H, m), 0.96 (3H, d, 6.7), 0.97 (1H, m), 1.02 (3H, s), 1.06 (1H, dd, 10.5; 5.5), 1.11

(1H, dd, 10.5; 9.5), 1.13 (1H, m), 1.15 (1H, m), 1.15 (1H, m), 1.19 (2H, d, 5.6), 1.22 (1H, m), 1.23 (2H, m), 1.27 (2H, m), 1.42 (1H, m), 1.55 (2H, m), 1.6 (2H, m), 1.69 (1H, dt, 9.5; 6.0), 1.77 (1H, dt, 9.0; 6.0), 1.94 (2H, dt, 5.6; 8.5), 1.95 (2H, m), 2.05 (2H, m), 2.21 (2H, d, 6.5), 3.39 (1H, m), 5.31 (1H, d, 5.6).

$^{13}\text{C}$ NMR (aseton-d<sub>6</sub>, 125 MHz),  $\delta$  37.4 (t) (C-1), 31.7 (t) (C-2), 70.9 (d) (C-3), 42.5 (t) (C-4), 141.5 (s) (C-5), 120.7 (d) (C-6), 31.8 (t) (C-7), 36.2 (d) (C-8), 50.4 (d) (C-9), 36.5 (s) (C-10), 20.9 (t) (C-11), 39.8 (t) (C-12), 42.2 (s) (C-13), 56.8 (d) (C-14), 25.9 (t) (C-15), 28.1 (t) (C-16), 56.1 (d) (C-17), 11.5 (q) (C-18), 18.4 (q) (C-19), 40.5 (d) (C-20), 19.3 (q) (C-21), 33.9 (t) (C-22), 24.1 (t) (C-23), 45.9 (d) (C-24), 31.9 (d) (C-25), 18.9 (q) (C-26), 18.5 (q) (C-27), 22.9 (t) (C-28), 11.4 (q) (C-29).

These data were found to be consistent with those reported in the literature for stigmast-5(6)-en-3 $\beta$ -ol or  $\beta$ -sitosterol (Chaturvedula and Prakash, 2012), and compound 1 identified as  $\beta$ -sitosterol.

Table 1: NMR Data compound 1.

C Position	stigmast-5(6)-en-3 $\beta$ -ol		Compound 1
	$^{13}\text{C}$ -NMR $\delta_c$ (ppm, mult.)	$^{13}\text{C}$ -NMR $\delta_c$ (ppm, mult.)	$^1\text{H}$ -NMR $\delta_H$ (ppm) ( $\Sigma\text{H}$ , mult., $J =$ Hz)
1	37.5	37.4 (t)	1.06 (1H, dd, 10.5; 5.5) 1.11 (1H, dd, 10.5; 9.5)
2	31.9	31.7 (t)	1.69 (1H, dt, 9.5; 6.0) 1.77 (1H, dt, 9.0; 6.0) 3.39 (1H, m)
3	72.0	70.9 (d)	2.21 (2H, d, 6.5)
4	42.5	42.5 (t)	-
5	140.9	141.5 (s)	-
6	121.9	120.7 (d)	5.31 (1H, d, 5.6)
7	32.1	31.8 (t)	1.94 (2H, dt, 5.6; 8.5)
8	34.2	36.2 (d)	1.42 (1H, m)
9	50.3	50.4 (d)	0.95 (1H, m)
10	36.7	36.5 (s)	-
11	21.3	20.9 (t)	1.55 (2H, m)
12	39.9	39.8 (t)	1.19 (2H, d, 5.6)
13	42.6	42.2 (s)	-
14	56.9	56.8 (d)	1.13 (1H, m)
15	26.3	25.9 (t)	1.23 (2H, m)
16	28.5	28.1 (t)	1.95 (2H, m)
17	56.3	56.1 (d)	1.15 (1H, m)
18	12.0	11.5 (q)	0.74 (3H, s)
19	19.0	18.4 (q)	1.02 (3H, s)
20	38.2	40.5 (d)	1.22 (1H, m)
21	19.2	19.3 (q)	0.86 (3H, d, 6.1)
22	34.2	33.9 (t)	2.05 (2H, m)
23	26.1	24.1 (t)	1.6 (2H, m)
24	46.1	45.9 (d)	0.97 (1H, m)
25	29.4	31.9 (d)	1.15 (1H, m)
26	20.1	18.9 (q)	0.84 (3H, d, 6.2)
27	19.6	18.5 (q)	0.96 (3H, d, 6.7)
28	23.3	22.9 (t)	1.27 (2H, m)
29	12.2	11.4 (q)	0.88 (3H, t, 1.89)

\*Measurements were taken at aseton-d<sub>6</sub> on 500 MHz for  $^1\text{H}$  and 125 MHz for  $^{13}\text{C}$ .

### Stigmast-4-en-3-one (2),

Crystal white needles; melting point 87-89°C,  $^1\text{H}$  NMR (Pyridin-D<sub>6</sub>, 600 MHz), Table 2;  $^{13}\text{C}$  NMR (Pyridin-D<sub>6</sub>, 150 Mhz), table 3.

Table 2: NMR data compound 2.

C position	Stigmast-4-en-3-one		Compound 2	
	<sup>13</sup> C-NMR δ <sub>c</sub> (ppm, mult.)	<sup>1</sup> H-NMR δ <sub>H</sub> (ppm) (ΣH, mult., J = Hz)	<sup>13</sup> C-NMR δ <sub>c</sub> (ppm, mult.)	<sup>1</sup> H-NMR δ <sub>H</sub> (ppm) (ΣH, mult., J = Hz)
1	35.9	1.48; 1.23	35.7 (t)	1.76 (1H, m) 1.44 (1H, m)
2	33.9	2.36; 2.31	34.0 (t)	2.32 (1H, m) 2.37 (1H, m)
3	199.6	-	199.3 (s)	-
4	123.7	5.70	123.9 (d)	5.80 (1H, s)
5	171.7	-	171.6 (s)	-
6	32.0	2.00; 1.90	32.6 (t)	2.03 (2H, m)
7	29.3	1.17; 1.41	32.1(t)	1.56 (1H, m) 1.61 (1H, m)
8	35.6	1.35	35.4 (d)	1.76 (1H, m)
9	53.8	1.32	53.7 (d)	0.68 (1H,m)
10	39.6	-	38.5 (s)	-
11	21.0	1.54; 1.26	20.9 (t)	1.32 (2H, m)
12	39.6	1.55; 1.31	39.7 (t)	1.87 (2H, m)
13	42.8	-	42.4 (s)	-
14	55.9	1.02	55.8 (d)	0.82 (1H, m)
15	24.1	1.61; 1.33	28.3 (t)	1.62 (1H, m) 1.75 (1H, m)
16	33.8	1.61; 1.33	24.2 (t)	1.45 (2H, d, 9.6)
17	55.8	1.15	56.0 (d)	1.02 (1H, m)
18	11.9	0.70	11.9 (q)	0.61 (3H, s)
19	17.3	1.17	16.9 (q)	0.95 (3H, s)
20	36.1	1.64	36.2 (d)	1.31 (1H, m)
21	18.6	0.97	18.8 (q)	0.79 (3H, d, 6.6)
22	35.6	1.26; 1.26	34.2 (t)	1.32 (2H, m)
23	26.0	1.25; 1.25	26.2 (t)	1.18 (2H, m)
24	45.8	1.47	45.8 (d)	1.19 (2H, m)
25	28.1	1.83	29.3 (d)	0.94 (1H, m)
26	19.8	0.83	19.8 (q)	0.81 (3H, d, 6.5)
27	19.0	0.81	19.0 (q)	0.79 (3H, d, 6.7)
28	23.0	1.54; 1.54	23.2 (t)	1.23 (2H, m)
29	14.1	0.90	12.0 (q)	0.83 (3H, t, 6.5)

\*400 MHz for <sup>1</sup>H and 100 MHz for <sup>13</sup>C, in CDCl<sub>3</sub> (Halilu *et al.*, 2013) and Measurements compound 3 were taken at 600 MHz for <sup>1</sup>H and 150 MHz for <sup>13</sup>C, in Pyridin-D<sub>6</sub>

Table 3: NMR data compound 3.

C Position	quercetin 3-O-β-D-glucopyranoside		Compound 3	
	<sup>13</sup> C NMR δ <sub>c</sub> (mult)	<sup>1</sup> H NMR δ <sub>H</sub> (Integral, mult, J=Hz)	<sup>13</sup> C NMR δ <sub>c</sub> (mult)	<sup>1</sup> H NMR δ <sub>H</sub> (Integral, mult, J=Hz)
2	158.44 (s)	-	156.97 (s)	-
3	135.64 (s)	-	134.09 (s)	-
4	179.48 (s)	-	178.96 (s)	-
5	162.99 (s)	-	161.36 (s)	-
6	99.89 (d)	6.19 (1H, d, 2)	100.38 (d)	-
7	165.97 (s)	-	163.15 (s)	6.07 (1H, d, 2.4)
8	94.73 (d)	6.38 (1H, d, 2)	94.67 (d)	-
9	158.44 (s)	-	157.60 (s)	6.23 (1H, d, 2.4)
10	105.68 (s)	-	102.68 (s)	-
1'	123.08 (s)	-	121.61 (s)	-
2'	117.59 (d)	7.71 (1H, d, 2)	116.00 (d)	-
3'	145.87 (s)	-	144.73 (s)	7.68 (1H, d, 2.4)
4'	149.83 (s)	-	148.81 (s)	-
5'	116.01 (d)	6.87 (1H, d, 8.4)	114.67 (d)	-
6'	123.20 (d)	7.58 (1H, dd, 2; 8.4)	121.75 (d)	6.82 (1H, d, 8.4)
1''	104.39 (d)	5.23 (1H, d, 7.6)	103.54 (d)	7.55 (1H, dd, 2.4; 8.4)
2''	75.73 (d)	3.48 (1H, d, 9.2)	74.35 (d)	5.11 (1H, dd, 6.8; 7.2)
3''	78.11 (d)	3.35 (1H, d, 8.8)	76.81 (d)	3.46 (1H, d, 9)
4''	71.22 (d)	3.43 (1H, d, 9.6)	69.78 (d)	3.39 (1H, d, 9.6)
5''	78.35 (d)	3.24 (1H, m)	77.00 (d)	3.32 (1H, d, 9.6)
6''	62.58 (t)	3.73 (1H, dd, 2; 11.6) 3.56 (1H, dd, 5.2; 11.6)	61.14 (t)	3.19 (1H, m) 3.68 (1H, dd, 2.4; 11.4) 3.56 (1H, dd, 6.0; 11.4)

\* 400 MHz for <sup>1</sup>H and 100 MHz for <sup>13</sup>C, in CD<sub>3</sub>OD (Islam *et al.*, 2012) and Measurements compound 3 were taken at 600 MHz for <sup>1</sup>H and 150 MHz for <sup>13</sup>C.

$^1\text{H}$  NMR (Pyridin-D<sub>6</sub>, 600 MHz)  $\delta$  0.61 (3H, s), 0.68 (1H, m), 0.79 (3H, d, 6.6), 0.79 (3H, d, 6.7), 0.81 (3H, d, 6.5), 0.82 (1H, m), 0.83 (3H, t, 6.5), 0.94 (1H, m), 0.95 (3H, s), 1.02 (1H, m), 1.18 (2H, m), 1.19 (2H, m), 1.23 (2H, m), 1.31 (1H, m), 1.32 (2H, m), 1.32 (2H, m), 1.44 (1H, m), 1.45 (2H, d, 9.6), 1.56 (1H, m), 1.61 (1H, m), 1.62 (1H, m), 1.75 (1H, m), 1.76 (1H, m), 1.76 (1H, m), 1.87 (2H, m), 2.03 (2H, m), 2.32 (1H, m), 2.37 (1H, m), 5.80 (1H, s).

$^{13}\text{C}$  NMR (Pyridin-D<sub>6</sub>, 150 Mhz)  $\delta$  35.7 (t) (C-1), 34.0 (t) (C-2), 199.3 (s) (C-3), 123.9 (d) (C-4), 171.6 (s) (C-5), 32.6 (t) (C-6), 32.1 (t) (C-7), 35.4 (d) (C-8), 53.7 (d) (C-9), 38.5 (s) (C-10), 20.9 (t) (C-11), 39.7 (t) (C-12), 42.4 (s) (C-13), 55.8 (d) (C-14), 28.3 (t) (C-15), 24.2 (t) (C-16), 56.0 (d) (C-17), 11.9 (q) (C-18), 16.9 (q) (C-19), 36.2 (d) (C-20), 18.8 (q) (C-21), 34.2 (t) (C-22), 26.2 (t) (C-23), 29.3 (d) (C-24), 45.8 (d) (C-25), 19.8 (q) (C-26), 19.0 (q) (C-27), 23.2 (t) (C-28), 12.0 (q) (C-29). These data were found to be consistent with those reported in the literature for Stigmast-4-en-3-one (Halilu *et al.*, 2013; Barla *et al.*, 2006 and Della Greca *et al.*, 1990), and compound 2 identified as Stigmast-4-en-3-one.

### 3-O- $\beta$ -D-glucopyranoside (3)

Yellow powder, melting point 872.6 °C.  $^1\text{H}$  NMR (Pyridin-D<sub>6</sub>, 600 MHz), table 3;  $^{13}\text{C}$  NMR (Pyridin-D<sub>6</sub>, 150 Mhz), table 3.

$^1\text{H}$  NMR (Pyridin-D<sub>6</sub>, 600 MHz)  $\delta$  6.07 (1H, d, 2.4), 6.23 (1H, d, 2.4), 7.68 (1H, d, 2.4), 6.82 (1H, d, 8.4), 7.55 (1H, dd, 2.4; 8.4), 5.11 (1H, dd, 6.8; 7.2), 3.46 (1H, d, 9), 3.39 (1H, d, 9.6), 3.32 (1H, d, 9.6), 3.19 (1H, m), 3.68 (1H, dd, 2.4; 11.4), 3.56 (1H, dd, 6.0; 11.4).

$^{13}\text{C}$  NMR (Pyridin-D<sub>6</sub>, 150 Mhz)  $\delta$  156.97 (s) (C-2), 134.09 (s) (C-3), 178.96 (s) (C-4), 161.36 (s) (C-5), 100.38 (d) (C-6), 163.15 (s) (C-7), 94.67 (d) (C-8), 157.60 (s) (C-9), 102.68 (s) (C-10), 121.61 (s) (C-1'), 116.00 (d) (C-2'), 144.73 (s) (C-3'), 148.81 (s) (C-4'), 114.67 (d) (C-5'), 121.75 (d) (C-6'), 103.54 (d) (C-1''), 74.35 (d) (C-2''), 76.81 (d) (C-3'') 69.78 (d) (C-4''), 77.00 (d) (C-5''), 61.14 (t) (C-6'').

These data were found to be consistent with those reported in the literature for 3-O- $\beta$ -D-glucopyranoside (Shi *et al.*, 2016; Islam *et al.*, 2012 and Li *et al.*, 2008), and compound 3 identified as 3-O- $\beta$ -D-glucopyranoside.

Compound 1 obtained in the form of colorless white crystal with a melting point of 134-136 °C and soluble in acetone. The molecular formula of compound 1 was established as C<sub>29</sub>H<sub>50</sub>O based HR-TOFMS spectrum (negative ion mode)  $m/z$  413.0911, the calculation for C<sub>29</sub>H<sub>50</sub>O  $m/z$  414.0818 with NMR data (Table 1), therefore required five double bond equivalent. Compound 1 does not fluoresce under ultraviolet light at a wavelength of 254 and 360 nm, indicating the absence of conjugated double bonds. Infrared absorption due to a hydroxyl group, a double bond, and the ether was observed in 3440, 2934, 2890, 1640, 1190 cm<sup>-1</sup>.  $^1\text{H}$  NMR spectrum compounds 1 expressed their olefinic proton signals [ $\delta_{\text{H}}$  5.31 (1H, d,  $J=4.9$  Hz), six methyl proton signal consisting of two methyl tertiary [ $\delta_{\text{H}}$  0.74 (3H, s) and 1.02 (3H, s)], four secondary methyl [ $\delta_{\text{H}}$  0.84 (3H, d,  $J=6.2$  Hz), 0.86 (3H, d,

$J=6.1$  Hz), 0.88 (3H, t,  $J=1.89$  Hz) dan 0.96 (3H, d,  $J=6.7$  Hz)] and one methyn signal sp<sup>3</sup> oxygenated on 3.39 (1H, m). Total carbon signals twenty-nine observed in  $^{13}\text{C}$  NMR spectrum. These signals are specified with DEPT and HMQC experiment as two carbon sp<sup>2</sup> ( $\delta_{\text{C}}$  120.7 and 141.5), six methyls, eleven methylene sp<sup>3</sup>, seven methyn sp<sup>3</sup>, one methyn oxygenated with  $\delta_{\text{C}}$  70.9 and two carbon sp<sup>3</sup> quartener. This functionality is counted as one of the totals of five double bond equivalent. Comparison of physical data and physicochemical compound 1 with stigmast-5(6)-en-3 $\beta$ -ol (Chaturvedula and Prakash, 2012) shows a very high suitability, thus compound 1 was identified as stigmast-5(6)-en-3 $\beta$ -ol or  $\beta$ -sitosterol.

The reported of  $\beta$ -phytosterol compounds in plants *L. microphyllum* first reported in this study thus provide clues phytochemical their steroid compounds in the genus *Lygodium*.

Compound 2 obtained in the form of a colorless white crystals with a melting point 87-89°C and soluble in chloroform.  $^1\text{H}$  NMR spectrum compounds show six methyl signal at  $\delta_{\text{H}}$  ppm 0.61 (3H), 0.79 (3H), 0.79 (3H), 0.81 (3H), 0.83 (3H), and 0.95 (3H), these suggest that there are six methyl group. The presence of down field signal at  $\delta_{\text{H}}$  ppm 5.80 (1H) shows olefinic proton.  $^{13}\text{C}$  NMR spectrum show 29 signal (Table 2) suggesting that compound consist of 29 carbon atom. Compound 2 does not fluorescent under ultraviolet light at a wavelength 254 and 360 nm, indicated the absence of conjugated double bonds. Comparison of physical data and physicochemical compound 2 Stigmast-4-en-3-one shows a very high suitability (Halilu *et al.*, 2013; Barla *et al.*, 2006 and Della Greca *et al.*, 1990), thus compound 2 was identified as Stigmast-4-en-3-one.

Compound 3 obtained in the form of yellow powder with a melting point 872.6 °C and completely soluble in methanol, Fluorescent under ultraviolet light at a wavelength 254 and 360 nm, indicated there is founded conjugated double bonds. Comparison of physical data and physicochemical compound 3 3-O- $\beta$ -D-glucopyranoside or Isoquercetin shows a very high suitability (Shi *et al.*, 2016; Islam *et al.*, 2012 and Li *et al.*, 2008), thus compound 3 was identified as 3-O- $\beta$ -D-glucopyranoside or Isoquercetin.

## CONCLUSION

Two steroids, stigma-5(6)-en-3 $\beta$ -ol or  $\beta$ -sitosterol (1) and Stigmast-4-en-3-one (2) and quercetin 3-O- $\beta$ -D-glucopyranoside or Isoquercetin (3) in plants *L. microphyllum* first reported in this study.

## ACKNOWLEDGMENT

Authors gratefully acknowledge to Directorate General of Higher Education, Ministry of Research, Technology and higher education, Indonesia on funds Competitive Research Grant and PKPI Sandwich Like 2015 Scholarship. Our thanks also to Prof. Dr. Unang Supratman in Central Laboratory, Padjadjaran University, Mr. Ahmad Darmawan, M.Sc. and Sofa Fajriah, M.Si in Chemical Research Center, Indonesian Institute of Sciences,

Serpong, Tangerang and Prof. Dr. Yoshito Shiono in Laboratory of Natural Products, Department Of Bioresources Engineering, Faculty of Agriculture, Yamagata University, Japan for NMR measurements.

## REFERENCES

- Barla A, Bırman H, KÜLTÜR Ş, and Öksüz S. Secondary metabolites from *Euphorbia helioscopia* and their vasodepressor activity. *Turkish Journal of Chemistry*. 2006; 30(3): 325-332. Available at: <http://journals.tubitak.gov.tr/chem/abstract.htm?id=8338>.
- Cambie RC & Ash J., 1994. *Fijian Medicinal Plants*. CSIRO. Melbourne.
- Chaturvedula VSR & Prakash I. Isolation of stigmasterol and  $\beta$ -sitosterol from the dichloromethane extract of *Rubus suavissimus*. *International Current Pharmaceutical Journal*. 2012; 1(9): 239-242. [ONLINE] Available at <http://imsear.hellis.org/handle/123456789/152876>
- Chen L, Zhang G, He J, Jin Guan J, Pan C, Mi W, & Wang Q. New naphthoquinone from the root of *Lygodium japonicum* (Thunb.) Sw. *Journal of Natural Medicine*. 2010; 64: 114–116. doi: 10.1007/s11418-009-0376-y
- Della Greca M, Monaco P, & Previtera L. Stigmasterols from *Typha latifolia*. *Journal of Natural Products*. 1990; 53(6): 1430-1435. DOI: 10.1021/np50072a005.
- Duan Y, Dai, He YR, Kurihara H, Li Y, & Yao X. A new phenylpropanoid glucoside from the aerial parts of *Lygodium japonicum*. *Journal of Asian Natural Products Research*. 2012; 14 (3): 286–292. doi:10.1080/10286020.2011.650690
- Guo-Gang Z, Ying-Cui H, Hong-Xia L, Lin-Xia Z, & Li-Juan C. 2012. *The Research of Lygodium*. Intechopen. Shanghai.
- Halilu ME, October N, Balogun M, Agunu A, Abubakar A. & Abubakar MS. Isolation and Characterization of Steroids from Petroleum Ether Extract of Stem Bark of *Parinari curatellifolia* Planch ex. Benth (Chrysobalanaceae). *Journal of Natural Sciences Research*. 2013; 3(6).
- Islam M, Al-Amin MD, Siddiqi MMA, Akter S, Haque MM, Sultana N, & Chowdhury AS, Isolation of quercetin-3-O-beta-D-glucopyranoside from the leaves of *Azadirachta Indica* and antimicrobial and cytotoxic screening of the crude extracts. *Dhaka University journal of science*. 2012; 60(1):11-14.
- Karthik V, Raju K, Ayyanar M, Gowrishankar K, & Sekar T. Ethnomedicinal uses of pteridophytes in Kolli Hills, Eastern Ghats of Tamil Nadu, India. *Journal of Natural Products Plant Resources*. 2011; 50-55
- Lee S, Xiao C, & Pei S. Ethnobotanical survey of medicinal plants at periodic markets of Honghe Prefecture in Yunnan Province, China. *Journal of Ethnopharmacology*. 2008; 117: 362–377. doi:10.1016/j.jep.2008.02.001
- Li YL, Li J, Wang NL, & Yao XS. Flavonoids and a new polyacetylene from *Bidens parviflora* Willd. *Molecules*. 2008; 13(8): 1931-1941. doi:10.3390/molecules13081931.
- Shi GQ, Yang J, Liu J, Liu SN, Song HX, Zhao WE, & Liu YQ. Isolation of flavonoids from onion skins and their effects on K562 cell viability. *Bangladesh Journal of Pharmacology*, 2016; 11(S1): 18-25. DOI: <http://dx.doi.org/10.3329/bjpv.v11iS1.26419>.
- Upreti K, Jalal JS, Tewari LM, Joshi GC, Pangtey YPS, & Tewari G. Ethnomedicinal uses of Pteridophytes of Kumaun Himalaya, Uttarakhand, India. *Journal of American Science*. 2009; 5: 167–170.
- Wallace RA, Sander GP, & Ferl RJ. 1991. *Biology: The Science of Life*. Harper Collins. New York.
- Ye W, Fan C, Zhang L, Yin Z, & Zhao S. A new phenolic glycoside from the roots of *Lygodium japonicum*. *Fitoterapia*. 2007; 78: 600–601. doi:10.1016/j.fitote.2007.04.003
- Zhang L, Yin Z, Ye W. Flavonoids from *Lygodium japonicum*. *Biochemical Systematics and Ecology*. 2006; 34: 885-886. doi: 10.1016/j.bse.2006.04.009
- Zeng-fu LI, Huil H, Hang-yi Z, & Jun-chen Z. Review on the extraction of flavonoids from fern. *Journal of San University*. 2008; 25: 22.
- Zheng XL, & Xing FW. Ethnobotanical study on medicinal plants around Mt. Yinggeling, Hainan Island, China. *Journal of Ethnopharmacology*. 2009; 124: 197–210. doi:10.1016/j.jep.2009.04.042
- Zhu L, Zhang G, Chen L, Wang S, Li P, Li L. A new ecdysteroside from *Lygodium japonicum* (Thunb.) Sw. *Journal of Natural Medicinal*. 2009; 63: 215–219. doi:10.1007/s11418-008-0310-8.

### How to cite this article:

Kuncoro H, Farabi K, Rijai L. Steroids and isoquercetin From *Lygodium microphyllum*. *J App Pharm Sci*, 2017; 7 (11): 136-141.

# Steroids and isoquercetin from *Lygodium microphyllum*

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

16%

PUBLICATIONS

8%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Choudhary, M.I.. "Cholinesterase inhibitory pregnane-type steroidal alkaloids from *Sarcococca hookeriana*", *Steroids*, 200504  
Publication 2%
- 2 [banglajol.info](http://banglajol.info)  
Internet Source 1%
- 3 "Pharmacology is the Backbone of Rational Prescribing", *Journal of Applied Pharmaceutical Science*, 2017  
Publication 1%
- 4 Submitted to Universiti Teknologi Malaysia  
Student Paper 1%
- 5 [journals.itb.ac.id](http://journals.itb.ac.id)  
Internet Source 1%
- 6 Ashish Kumar, Gopi Chand, Vijai K. Agnihotri. "A new oxo-sterol derivative from the rhizomes of ", *Natural Product Research*, 2017  
Publication 1%
- 7 [www.journal.uinjkt.ac.id](http://www.journal.uinjkt.ac.id)  
Internet Source 1%



8

Xiang-Rong Tian, Hai-Feng Tang, Yu-Shan Li, Hou-Wen Lin, Xiao-Li Chen, Ning Ma, Min-Na Yao, Ping-Hu Zhang. "New Cytotoxic Oxygenated Sterols from the Marine Bryozoan *Cryptosula pallasiana*", *Marine Drugs*, 2011

Publication

1%

9

Ace Tatang Hidayat, Achmad Zainuddin, Danar Dono, Wawan Hermawan, Hideo Hayashi, Unang Supratman. "Synthetic and Structure-Activity Relationship of Insecticidal Bufadienolides", *Natural Product Communications*, 2014

Publication

1%

10

Luo, Y.. "Steryl esters and phenylethanol esters from *Syringa komarowii*", *Steroids*, 200608

Publication

1%

11

[repository.hkbu.edu.hk](http://repository.hkbu.edu.hk)

Internet Source

1%

12

[journal.unpad.ac.id](http://journal.unpad.ac.id)

Internet Source

1%

13

Julinton Sianturi, Mayshah Purnamasari, Darwati, Desi Harneti, Tri Mayanti, Unang Supratman, Khalijah Awang, Hideo Hayashi. "New bisamide compounds from the bark of *Aglaia eximia* (Meliaceae)", *Phytochemistry Letters*, 2015

1%

14

[de.slideshare.net](http://de.slideshare.net)

Internet Source

1%

---

15

Arrieta-Baez, D.. "Modeling suberization with peroxidase-catalyzed polymerization of hydroxycinnamic acids: Cross-coupling and dimerization reactions", *Phytochemistry*, 200604

Publication

1%

---

16

[docplayer.net](http://docplayer.net)

Internet Source

1%

---

17

Yinggang Luo. "A new glucoceramide from the watermelon begonia, *Pellionia repens*", *Lipids*, 10/2004

Publication

1%

---

18

[japsonline.com](http://japsonline.com)

Internet Source

1%

---

Exclude quotes  On

Exclude matches  < 1%

Exclude bibliography  On

# Steroids and isoquercetin from *Lygodium microphyllum*

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---