

PhcogJ 4

by Islamudin Ahmad

Submission date: 28-Feb-2022 12:41AM (UTC+0000)

Submission ID: 1772275511

File name: PhcogJ_Vol_13_4.pdf (667.85K)

Word count: 4095

Character count: 21546

Phytochemicals and Antioxidant Activities Evaluation of *Origanum vulgare* (L.) Stem Bark Extracts

Rosmalena¹, Putu Ayu Widyastuti², Fatmawaty Yazid¹, Neneng Siti Silfi Ambarwati^{3,*}, Islamudin Ahmad⁴

Rosmalena¹, Putu Ayu Widyastuti², Fatmawaty Yazid¹, Neneng Siti Silfi Ambarwati^{3,*}, Islamudin Ahmad⁴

¹Department of Medical Chemistry, Faculty of Medicine, Universitas Indonesia, South Jakarta, 10430 Jakarta, INDONESIA

²Medical Student, Faculty of Medicine, Universitas Indonesia, South Jakarta, 10430 Jakarta, INDONESIA

³Department of Cosmetology, Faculty of Engineering, Universitas Negeri Jakarta, East Jakarta, 13220 Jakarta, INDONESIA

⁴Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Mulawarman, Samarinda, 75119 East Kalimantan, INDONESIA

Correspondence

Neneng Siti Silfi Ambarwati

Department of Cosmetology, Faculty of Engineering, Universitas Negeri Jakarta, East Jakarta, 13220 Jakarta, INDONESIA

Phone no. +62399439923

E-mail: nenengsitisilfi@unj.ac.id

History

- Submission Date: 31-03-2021;
- Review completed: 12-05-2021;
- Accepted Date: 25-05-2021.

DOI : 10.5530/pj.2021.13.124

Article Available online

<http://www.phcogj.com/v13/i4>

Copyright

© 2021 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



ABSTRACT

The present study aimed to evaluate phytochemical and antioxidant activity (*in vitro* and *in vivo*) of *Origanum vulgare* (L.) ethanolic extract. The phytochemical test was assessed using the Clute method in ethanol, ethyl acetate, and hexane. *In vitro* evaluation of antioxidant activity was determined by radical scavenging assay using DPPH (2,2-diphenyl-1-picrylhydrazyl) as an artificial free radical activity. *In vivo* test was conducted to evaluate the effect of malondialdehyde (MDA) level in blood plasma during maximum physical activity treatment. *In vivo* test was done using 25 male Sprague Dawley rats in pre and post-test control group design. The phytochemical test of *O. vulgare* ethanol extract was showed some compounds, such as a flavonoid, alkaloid, triterpenoid/steroid, essential oil, and tannin, then in ethyl acetate and hexane. *In vitro* assay showed that *O. vulgare* extract has strong antioxidant activity with an IC₅₀ value of 133.47 µg/mL. While in the *in vivo* test, the most effective dosage is 20 mg/200 gr B.W., represented by a significant decrease of MDA level (0.509 nmol/mL) before and after treatment. So, the ethanolic extract of clove has potency as an herbal antioxidant because of the low level of IC₅₀ and can decrease the MDA level.

Key Words: 2,2-diphenyl-1-picrylhydrazyl, Antioxidant activity, Malondialdehyde, *Origanum vulgare* (L.), Phytochemical.

INTRODUCTION

The most common cause of death in Indonesia is a degenerative disease.¹ The three main causes of death in Indonesia in 2016 were ischemic heart disease, cerebrovascular disease, and diabetes.² Degenerative diseases can occur due to damage to cells in the body by free radicals.³ Free radicals are reactive oxygen atoms because the electrons are unpaired, also called reactive oxygen species (ROS).⁴ These free radical compounds are unstable, very reactive, and try to stabilize themselves in attempting to take electrons from other compounds (including proteins, carbohydrates, lipids, and DNA).⁵ ROS attacks normal cells in the body and causes cells to become damaged, called oxidative stress.⁶ Oxidative stress is a condition of imbalance between prooxidants and antioxidants in our bodies.⁷

Oxidative stress conditions can be overcome by consuming antioxidants to balance prooxidants.⁸ Oxidative stress can cause more free radicals; cell damage occurs, cells become susceptible to infection, degenerative diseases, DNA cell division is disturbed, causing cell mutations.⁵ This oxidative damage can accompany most or even cancers, diabetes, heart disease, arthritis, neurodegenerative disorders, atherosclerosis, osteoporosis, pancreatitis, special health for women, and pre-eclampsia.⁵ We know that free radicals are produced continuously during normal respiration and metabolism, triggered by exposure to air pollution, radiation from X-rays, exposure to sunlight, drugs, viruses, bacteria, parasites, dietary fat, stress, and injury.⁵

Antioxidant compounds can prevent oxidation of other compounds by oxidizing themselves, thus preventing cell damage.⁵ Therefore, antioxidants play a role in preventing diseases such as cancer, heart disease, dementia, diabetes, and stroke.^{5,9} The most significant source of antioxidants in plants is *Origanum vulgare* (L.), which grows well in Indonesia. This study was conducted to test the antioxidants contained in *O. vulgare* and the ability to neutralize free radicals *in vitro* (with the DPPH test) and *in vivo* using Sprague Dawley mice. This study hypothesizes that *O. vulgare* has exogenous antioxidant effects on free radicals, both *in vitro* through IC₅₀ and *in vivo* values through the MDA level.

This study aimed to provide knowledge to Indonesian people about the potential of *O. vulgare* as one of the herbal antioxidants, which is useful for treating oxidative stress because it is widely used in daily life.

MATERIALS AND METHODS

The activity test of *O. vulgare* extract *in vivo* was carried out with the Faculty of Medicine's ethics committee's approval, the University of Indonesia. The ethical license number is 0071/UN2.F1/ETIK/2018. The making of *O. vulgare* extract and IC₅₀ determination were carried out in the Department of Chemical Medicine, Faculty of Medicine, University of Indonesia. The phytochemical analysis and measurement of Malondialdehyde (MDA) levels from the *in vivo* assay were carried out in the Department of Biochemistry, Faculty of Medicine, University of Indonesia. Handling and treating animal models were carried out and followed the Laboratory of Animal Experiments and Toxicology of the Ministry of Health of the Republic of Indonesia.

Cite this article: Rosmalena, Widyastuti PA, Yazid F, Ambarwati NSS, Ahmad I. Phytochemicals and Antioxidant Activities Evaluation of *Origanum vulgare* (L.) Stem Bark Extracts. Pharmacogn J. 2021;13(4): 965-970.

Preparation of *O. vulgare* stem bark extract

The stem bark of *O. vulgare* was taken from the Research Institute for Spice and Medicinal Plants in Bogor, West Java. The bark was first cleaned and dried at room temperature for 24 hours, then smoothed using a blender.

Manufacture of *O. vulgare* extract

Making *O. vulgare* extract using bark powder (500 grams) with maceration method and solvent used 70% ethanol (1000 ml). After three days, the solution is filtered, and the filtrate is collected. Maceration is repeated until the filtrate is colorless. The collected filtrate was evaporated using a rotary evaporator (50 rpm, 30-40°C). The concentrated filtrate was then dried using an oven at 40°C. The extract was stored at 4°C.¹⁰

Phytochemical analysis

Phytochemical analysis was carried out to determine the compounds contained in the extract of *O. vulgare*. Before phytochemical testing, the extract was dissolved with three different solvents, namely 70% ethanol (polar), ethyl acetate (semi-polar), and hexane (non-polar). Phytochemical tests include the saponin test, flavonoid test, terpenoid, steroid test, alkaloid test, tannin test, and glycoside test.¹¹

In vitro test

In vitro antioxidant test to find 50% inhibition concentration (IC₅₀) by dissolving the extracted sample in methanol added 0.5 mL of 100 µg/ml 2,2-diphenyl-1-picrylhydrazyl (DPPH) solution in a test tube which had been wrapped in aluminum foil. The mixture was vortexed for 30 seconds and incubated inside dark for 30 minutes at 37°C.¹² The absorbance measurements were carried out using a spectrophotometer at a wavelength of 515 nm.¹³ DPPH inhibition was calculated using the formula.¹²

$$\% \text{ Inhibition} = \frac{\text{absorbance of blank} - \text{absorbance of sample}}{\text{absorbance of blank}} \times 100\%$$

IC₅₀ was obtained using linear regression analysis (Microsoft Excel 2015, version 15.17) from several points between the concentration of the sample (x) to the percentage of inhibition (y) with the equation $y = ax + b$.

In vivo test

Preparation for sprague dawley rats

This study used 25 healthy male Sprague Dawley rats aged 2 to 3 months with bodyweight between 180-200 grams, which were divided into five groups, namely the positive control group (5 mg vitamin C/200 gr b.w./day), the negative control group, the 5 mg/200 gr b.w./day extract group, the 10 mg/200 gr b.w./day extract group, and the dose group extract of 20 mg/200 gr b.w./day conducted about ten days.¹⁴

Malondialdehyde (MDA) test

Blood samples (500 µL) were taken at two times, before and after the rats were given extract for ten days, followed by physical exercise (swimming for ten minutes).¹⁵ Blood is taken using a capillary tube in the canthus orbital sinus and stored in a heparin tube.¹⁶ Blood plasma was obtained by centrifuging blood samples and stored at -20°C. A mixture of 250 µL of blood plasma with 100 µL of 8.1% sulfate dodecyl sodium (SDS), 750 µL of 0.5 M HCl, 125 µL aqua bidest, and 750 µL 20 M thiobarbiturate (TBA) vortexed and heated up to 90°C for 15 minutes. After being cooled for 10 minutes, 500 µL aqua bidest and 2.5 mL n-butanol pears were added and divorced. The mixture was then centrifuged at 3000

rpm, and the filtrate was measured for absorbance at a wavelength of 520 nm.¹⁷ To calculate the MDA level, a linear equation was made between the concentration of MDA (Y) and absorbance (X).¹⁸

Statistical analysis

Analysis of the MDA test data *in vivo* using a normality test followed by a one-way T and ANOVA test with SPSS 24 program.¹⁹ The normality test was used to determine whether the data were normally distributed ($p > 0.05$) or not.²⁰ The T-test was used to compare MDA levels before and after antioxidant administration and physical exercise.²¹ A one-way ANOVA test was used to determine the effectiveness of test samples as antioxidants.²²

RESULTS

Phytochemical analysis

The phytochemical analysis was conducted to observe the antioxidant compound present in the *O. vulgare* stem bark ethanolic extract (Table 1). “+” means those antioxidant substances are present, shown by a reaction in each extract with a different type of solvent, while “-” means the antioxidant substances are absent.

In vitro antioxidant activity of *O. vulgare* extract

In vitro antioxidant activity assay was conducted to know the extract's ability to neutralize free radicals from DPPH, shown by the IC₅₀ value (Table 2 for vitamin C, Table 3 for *O. vulgare* stem bark ethanolic extract) and then compared between the IC₅₀ value between Vitamin C and *O. vulgare* stem bark ethanolic extract.

In vivo antioxidant activity of *O. vulgare*

The antioxidant activity of *O. vulgare* was also evaluated by comparing the MDA level in the plasma before and after giving the treatment (Table 4).

Statistical analysis

A paired T-Test was used to know if there is a significant difference between two reliable data groups, shown by the value of $p < 0.05$ (Table 5).

One-Way ANOVA was used to evaluate if there is a significant difference in the mean of two or more unrelated variables (Table 6). It also determines the most effective dosage of the *O. vulgare* stem bark extract, reducing oxidative stress.

After conducting One-Way ANOVA, the following test is a Post Hoc Test based on the Test homogeneity of variances (Table 7). Post Hoc Test was used to know which data has a significant value.

Table 1: Phytochemical Analysis Result of *O. Vulgare* extract.

Phytochemicals Constituent	Extract		
	Ethanol	Ethyl acetate	n-Hexane
Glycosides	+	-	-
Saponin	+	+	-
Flavonoid	+	+	-
Alkaloid	+	+	-
Terpenoid/ Steroid	+	-	-
Essential oil	+	+	+
Tannin	+	+	-

Table 2: In Vitro Antioxidant Activity of Vitamin C.

Concentration ($\mu\text{g}/\text{mL}$)	Absorbance	Inhibition (%)	Equation	IC_{50} ($\mu\text{g}/\text{mL}$)
Blank	2.4718	0		
1	2.1888	11.45	$y = 3.1434x + 25.063$ $R^2 = 0.80956$	4.53
5	1.3529	46.05		
10	0.5303	76.28		
25	0.1117	95.35		

Table 3: In Vitro Antioxidant Activity of Ethanol Extract of *O. vulgare*.

Concentration ($\mu\text{g}/\text{mL}$)	Absorbance	Inhibition (%)	Equation	IC_{50} ($\mu\text{g}/\text{mL}$)
Blank	2.3238	0		
10	2.1472	7.60	$y = 0.3476x + 3.7099$ $R^2 = 0.84258$	133.47
50	2.1080	9.29		
100	1.0366	55.38		
200	0.7506	67.70		

Table 4: Mean and differences in MDA level.

No.	Groups	Mean and Difference of MDA Level		
		Before Treatment (nmol/ mL)	After Treatment (nmol/ mL)	MDA Level Differences (nmol/ mL)
1	Positive Control	0.924	0.670	-0.254
2	Negative Control	0.866	0.986	0.120
3	Extract 5 mg	1.354	0.854	-0.500
4	Extract 10 mg	1.180	0.706	-0.474
5	Extract 20 mg	1.218	0.709	-0.509

Table 5: The result of Paired T-Test of MDA level.

MDA Level	Number of Subjects	Mean (nmol/mL) \pm Standard Deviation	Mean Difference (nmol/mL) \pm Standard Deviation	95% Confidence Interval of the Difference	P
Before Treatment	25	1.132 (0.168)			
After Treatment	25	0.761 (0.107)	0.371 (0.178)	0.298 - 0.445	<0.0001

Table 6: The result of One-Way ANOVA of MDA level.

Groups	Number of Subjects	Mean (nmol/mL) (Standard Deviation)	P
Positive Control	5	-0.500 (0.135)	0.000
Negative Control	5	-0.474 (0.042)	
Extract 5 mg	5	-0.509 (0.099)	
Extract 10 mg	5	-0.254 (0.057)	
Extract 20 mg	5	-0.119 (0.053)	

Table 7: The result of Post Hoc Test.

		Mean Differences	95% Confidence Interval		P
			Minimum	Maximum	
Positive Control	Dose 10 mg	-0.246	-0.415	-0.077	0.002
	Dose 20 mg	-0.380	-0.549	-0.211	0.000
Negative Control	Dose 10 mg	-0.220	-0.389	-0.051	0.005
	Dose 20 mg	-0.355	-0.524	-0.186	0.000
Dose 5 mg	Dose 10 mg	-0.255	-0.424	-0.086	0.001
	Dose 20 mg	-0.389	-0.558	-0.220	0.000
Dose 10 mg	Positive Control	0.246	0.077	0.415	0.002
	Negative Control	0.220	0.051	0.389	0.005
Dose 20 mg	Dose 5 mg	0.255	0.086	0.424	0.001
	Positive Control	0.380	0.211	0.549	0.000
Dose 5 mg	Negative Control	0.355	0.186	0.524	0.000
	Dose 10 mg	0.389	0.220	0.558	0.000

DISCUSSION AND CONCLUSION

Phytochemical analysis

The phytochemical tests showed that only glycosides and triterpenoids or steroids were detected in the ethanol fraction. Saponins, flavonoids, alkaloids, and tannins were detected in the ethanol fraction and ethyl acetate fraction. And only essential oils are found in all fractions. Most compounds are detected in the ethanol fraction because they are polar solvents, while ethyl acetate is a semi-polar solvent. Essential oils can be detected in all fractions.

In vitro antioxidant activity of *O. vulgare*

The *in vitro* antioxidant activity of the extract of *O. vulgare* was determined using the DPPH test because of its effectiveness and specificity. The DPPH test also requires only a few extract samples. IC₅₀ is an extract concentration that can reduce the number of free radicals by 50%. The DPPH test is based on the reaction between DPPH as a free radical with a hydrogen-donor molecule from the extract of *O. vulgare*. IC₅₀ values obtained from the ethanol extract of the bark of *O. vulgare* was 133.47 µg/mL. According to the classification of antioxidant intensity by Jun et al. (2003), includes strong antioxidant (<50 µg/mL), active antioxidant (50-100 µg/mL), moderate antioxidant (101-250 µg/mL), weak antioxidant (251-500 µg/mL), and inactive antioxidant (>500 µg/mL). *O. vulgare* extract has moderate activity²³.

The value of R² = 0.84258 indicates that with the extract concentration increasing, the antioxidant activity of the extract of *O. vulgare* will also increase.

In vivo antioxidant activity of *O. vulgare* extract

Malondialdehyde (MDA) results from a long chain peroxidation of unsaturated fatty acids found in fat membranes²⁴. This peroxidation reaction occurs when the level of oxidative stress increases²⁵. The antioxidant activity of *O. vulgare* extract was tested *in vivo* by comparing the levels of MDA of rats before and after extracts and physical exercise. The comparison was analyzed statistically using the Paired T-Test method. The p-value obtained from the Paired T-Test is 0.000, smaller than 0.05. This showed that there are at least two groups of data that have significant differences between MDA levels before and after treatment. The mean MDA level before treatment was 1.132 ± 0.168 nmol/mL, while the MDA level after treatment was 0.761 ± 0.107 nmol/mL.

One-Way ANOVA was conducted to find out comparisons in the group as a whole. Significance (p) was 0.000 (< 0.05), indicating that there is a minimum of two groups that have significant differences. According to the Post Hoc Bonferroni analysis results, differences in MDA levels between the positive control and some different concentration of *O. vulgare* extract.

O. vulgare extract has IC₅₀ value as an antioxidant 133.47 (µg/mL). This shows that antioxidant activity is moderate in reducing DPPH radical levels in *In Vitro*. The ability of the antioxidant extract is not as strong as Vitamin C (positive control). *In vivo* studies showed that *O. vulgare* extract administration at doses of 10 mg and 20 mg can provide antioxidant effects compared to other groups.

ACKNOWLEDGMENTS

We would like to express our gratitude to International Indexed Publication Grant for Students Final Projects from the University of Indonesia 2018, which have supported this research. This study was supported by the grant of Publikasi Terindeks Internasional untuk Tugas Akhir Mahasiswa (PITTA) 2018, with grant number: 2124/UN2.R3.1/HKP.05.00/2018

CONFLICTS OF INTEREST


The authors declared no conflicts of interest.

REFERENCES

- Srimpton R, Rokx C. *The Double Burden of Malnutrition in Indonesia*; 2013.
- Mboi N, Surbakti IM, Trihandini I, Elyazar I, Smith KH, Ali PB, Kosen S, Fliemos K, Ray SE, Cao J, Glenn SD, Miller-Petrie MK, Mooney MD, Riedl JL, Ningrum DNA, Idris F, Siregar KN, Harimurti P, Bernstein RS, Pangestu T, Sidharta Y, Naghavi M, Murray CJL, Hay SI. On the road to universal health care in Indonesia, 1990–2016: A systematic analysis for the global Burden of Disease Study 2016. *Lancet*. 2018;392(10147):581-591. doi:10.1016/S0140-6736(18)30595-6
- Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods : Impact on human health. *Phcog Rev*. 2010;4(8):118-127. doi:10.4103/0973-7847.70902
- Aprioku JS, Aprioku JS, Har P. Pharmacology of free radicals and the impact of reactive oxygen species on the testis. *J Reprod Infertil*. 2013;14(20):158-172.
- Talaulikar VS, Manyonda IT. Vitamin C as an antioxidant supplement in women's health: A myth in need of urgent burial. *Eur J Obstet Gynecol Reprod Biol*. 2011;157(1):10-13. doi:10.1016/j.ejogrb.2011.03.017
- Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O. Oxidative stress and antioxidant defense. *World Allergy Organ J*. 2012;5(1):9-19. doi:10.1097/WOX.0b013e3182439613
- Rahal A, Kumar A, Singh V, Tadav B, Tiwari R. Oxidative stress, prooxidants, and antioxidants : The interplay. *BioMed Res Int*. 2014;2014:1-19. doi:10.1155/2014/761264
- Pojlsak B, Šuput D, Milisav I. Achieving the balance between ROS and antioxidants : When to use the synthetic antioxidants. *Oxid Med Cell Longev*. 2013;2013:1-12.
- Sati SC, Sati N. Medicinal plants as a source of antioxidants. *Res J Phytochem*. 2014;4(4):213-224. doi:10.3923/rjphyto.2010.213.224
- Ambarwati NSS, Elya B, Malik A, Hanafi M. Evaluation of antimicrobial activities of *Garcinia lattissima* Miq. stem bark extract. *J Young Pharm*. 2017;9(1):118-121.
- Ambarwati NSS, Elya B, Malik A, Hanafi M. Phytochemical and antimicrobial studies on *Garcinia lattissima* Miq. fruit extract. *Asian J Pharm Clin Res*. 2017;10(7):230-232. doi:10.22159/ajpcr.2017.v10i7.18528
- Ambarwati NSS, Elya B, Nur A, Puspitasari N, Malik A, Hanafi M. Activity of fractions from *Garcinia lattissima* Miq. leaves ethyl acetate extract as antibacterial against *Bacillus subtilis* and antioxidant. *Adv Sci Lett*. 2018;24(9):6366-6370. doi:10.1166/asl.2018.12717
- Saha MR, Hasan SMR, Akter R, Hossain MM, Alam MS, Alam MA, Mazumder MEH. *In vitro* free radical scavenging activity of methanol extract of the leaves of *Mimusops elengi* Linn. *Bangl J Vet Med*. 2008;6(2):197-202.
- Nna VU, Abu Bakar AB, Md Lazin MRML, Mohamed M. Antioxidant, anti-inflammatory and synergistic anti-hyperglycemic effects of Malaysian propolis and metformin in streptozotocin-induced diabetic rats. *Food Chem Toxicol*. 2018;120:305-320. doi:10.1016/j.fct.2018.07.028
- Hasanvand A, Pirzadroozbahani N, Ahmadizar F, et al. Evaluation of the antioxidant effects of Zolpidem in the rat model of cisplatin-induced nephrotoxicity. *J Ren Inj Prev*. 2018;7(4):235-239. doi:10.15171/jrip.2018.54
- Parasuraman S, Raveendran R, Kesavan R. Blood sample collection in small laboratory animals. *J Pharmacol Pharmacother*. 2010;1(2):87. doi:10.4103/0976-500X.72350
- Nagaraj, D SK, Paunipagar P.V. Study of serum malondialdehyde and vitamin C in smokers. *J Sci Innov Res*. 2014;3(6):569-571.
- Zeb A, Ullah F. A simple spectrophotometric method for the determination of thiobarbituric acid reactive substances in fried fast foods. *J Anal Methods Chem*. 2016;2016(9412767):1-5.

19. Park HM. *Univariate Analysis and Normality Test Using SAS, STATA, and SPSS*. <http://www.indiana.edu/~statmath>.
20. Ghasemi A, Zahediasl S. Normality tests for statistical analysis: A guide for non-statisticians Asghar. *Int J Endocrinol Metab*. 2012;10(2):486-489. doi:10.5812/ijem.3505
21. Rasyid HN, Orth MS, Ismiarto YD, Orth MS, Prasetya R, Orth MS. The efficacy of flavonoid antioxidant from chocolate bean extract: Prevention of myocyte damage caused by reperfusion injury in predominantly anaerobic sports. *Malaysian Orthop J*. 2012;6(3):3-6.
22. Martha R, Gutierrez P, Cotera LBF, Maria A, Gonzalez N. Evaluation of the antioxidant and anti-glycation effects of the hexane extract from *Piper auritum* leaves *in vitro* and beneficial activity on oxidative stress and advanced glycation end-product-mediated renal injury in streptozotocin-treated diabetic rats. *Molecules*. 2012;17:11897-11919. doi:10.3390/molecules171011897
23. Jun M, Fu HY, Hong J, Wan X, Yang CS, Ho CT. Comparison of antioxidant activities of isoflavones from kudzu root (*Pueraria lobata* Ohwi). *J Food Sci*. 2003;68(6):2117-2122. doi:10.1111/j.1365-2621.2003.tb07029.x
24. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev*. 2014;2014:1-31.
25. Wadhwa N, Mathew BB, Jatava SK, Tiwari A. Lipid peroxidation: mechanism, models and significance. *Int J Curr Sci*. 2012;3:29-38.

GRAPHICAL ABSTRACT



SUMMARY

- The phytochemical test of *O. vulgare* ethanol extract was showed some compounds, such as a flavonoid, alkaloid, triterpenoid/steroid, essential oil, and tannin, then in ethyl acetate and hexane.
- In vitro* assay showed that *O. vulgare* extract has strong antioxidant activity with an IC₅₀ value of 133.47 µg/mL.
- In vivo* test, the most effective dosage is 20 mg/200 gr B.W., represented by a significant decrease of MDA level (0.509 nmol/mL) before and after treatment.

ABOUT AUTHORS



9
Dr. Rosmalena Rosmalena, a lecturer and researcher at Medical Chemistry, Faculty of Medicine, Universitas Indonesia. Research interest in medicinal chemistry, and natural product chemistry.



Putu Ayu Widyatuti, a medical student at Faculty of Medicine, Universitas Indonesia. Research interest in medicinal chemistry.



Dra. Fatmawaty, M.Si, a lecturer and researcher at Medical Chemistry, Faculty of Medicine, Universitas Indonesia. Research interest in medicinal chemistry, and natural product chemistry.



Dr. Neneng Siti Silfi Ambarwati; a Lecturer at Cosmetology Department, Faculty of Engineering, Universitas Negeri Jakarta, East Jakarta, Indonesia. The research focused on natural products for drug and cosmetic discovery and development, extraction technology, and cosmetic ingredients (cosmeceuticals).



7
Dr. Islamudin Ahmad, a lecturer and Researcher at Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Mulawarman, East Kalimantan, Indonesia. He has experience in the area of Pharmacognosy and Natural Product Chemistry, working in drugs discovery of natural products, green extraction engineering, isolation and identification of active compound, screening activity mainly degenerative diseases.

Cite this article: Rosmalena, Widyastuti PA, Yazid F, Ambarwati NSS, Ahmad I. Phytochemicals and Antioxidant Activities Evaluation of *Origanum vulgare* (L.) Stem Bark Extracts. Pharmacog J. 2021;13(4): 965-970.

ORIGINALITY REPORT

15%

SIMILARITY INDEX

10%

INTERNET SOURCES

12%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | Submitted to School of Business and Management ITB
Student Paper | 2% |
| 2 | www.innovareacademics.in
Internet Source | 1% |
| 3 | Rahmawati Rahmawati, Harti Widiastuti, Eka Sulistya. "In Vitro Anti-Inflammatory Assay of Bitter Melon (<i>Momordica charantia</i> L.) Ethanol Extract", <i>Jurnal Fitofarmaka Indonesia</i> , 2020
Publication | 1% |
| 4 | Talaulikar, V.S.. "Vitamin C as an antioxidant supplement in women's health: a myth in need of urgent burial", <i>European Journal of Obstetrics and Gynecology</i> , 201107
Publication | 1% |
| 5 | Submitted to University of Melbourne
Student Paper | 1% |
| 6 | hal.archives-ouvertes.fr
Internet Source | 1% |
| 7 | islamudinahmad.com
Internet Source | |

1 %

8

María Pilar de Torre, Jose Luis Vizmanos, Rita Yolanda Caverro, María Isabel Calvo.

1 %

"Improvement of antioxidant activity of oregano (*Origanum vulgare* L.) with an oral pharmaceutical form", *Biomedicine & Pharmacotherapy*, 2020

Publication

9

Elvira Lesmana, Ade Arsianti, Trivani Putri, Priscilla Aya Maheswari Subroto.

1 %

"Phytochemistry test and cytotoxic activity of macroalgae *Euclima* Sp. as a growth inhibitor of breast MCF-7 cancer cells", *AIP Publishing*, 2019

Publication

10

Primariadewi Rustamadji, Elvan Wiyarta, Tricia Dewi Anggraeni, Trifonia Pingkan Siregar.

1 %

"Adult granulosa cell tumor with minor foci of juvenile granulosa cell tumor in postmenopausal woman: A rare case report", *International Journal of Surgery Case Reports*, 2021

Publication

11

www.mdpi.com

Internet Source

1 %

12 Xiao-Li Zhang, Yu-Shan Guo, Chun-Hua Wang, Guo-Qiang Li, Jiao-Jiao Xu, Hau Yin Chung, Wen-Cai Ye, Yao-Lan Li, Guo-Cai Wang. "Phenolic compounds from *Origanum vulgare* and their antioxidant and antiviral activities", *Food Chemistry*, 2014
Publication

13 scitepress.org
Internet Source

14 www.dovepress.com
Internet Source

15 repository.cardiffmet.ac.uk
Internet Source

16 himatipanunpad.blogspot.com
Internet Source

17 Marissa Angelina, Muhammad Hanafi, Franciscus D. Suyatna, T. Mirawati S., Shirley Ratnasari, Beti Ernawati Dewi. " Antiviral Effect of Sub Fraction Leaves Extract to Dengue Virus Serotype-2 strain New Guinea C in Human Cell Line Huh-7 it-1 ", IOP Conference Series: Earth and Environmental Science, 2017
Publication

18 pt.scribd.com
Internet Source

19 Baiceanu, Elisabeta, Laurian Vlase, Andrei Baiceanu, Madalina Nanes, Dan Rusu, and Gianina Crisan. "New Polyphenols Identified in Artemisiae abrotani herba Extract", *Molecules*, 2015. <1 %
Publication

20 Submitted to Syiah Kuala University <1 %
Student Paper

21 apjmt.mums.ac.ir <1 %
Internet Source

22 jurnal.ugm.ac.id <1 %
Internet Source

23 mafiadoc.com <1 %
Internet Source

24 tjnpr.org <1 %
Internet Source

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On

PhcogJ 4

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
