

JAPTR

by Islamudin Ahmad

Submission date: 28-Feb-2022 12:45AM (UTC+0000)

Submission ID: 1772277952

File name: JAdvPharmTechRes_2022_13_1_50_336199_compressed.pdf (1.25M)

Word count: 3932

Character count: 21243



Journal of Advanced Pharmaceutical Technology & Research

An Official Publication of Society of Pharmaceutical Education & Research

Volume 13 / Issue 1 / January-March 2022

www.japtr.org



Medknow

JAPTR

New robustaflavone from *Garcinia latissima* Miq. leave and Its antibacterial activity

Neneng Siti Silfi Ambarwati,
Berna Elya¹, Amarila Malik²,
Hanita Omar³,
Muhammad Hanafi^{4,5},
Islamudin Ahmad⁶

⁹ Department of Cosmetology, Faculty of Engineering, Universitas Negeri Jakarta, East Jakarta, ⁵Department of Pharmaceutical Sciences, Faculty of Pharmacy, Faculty of Pharmacy, Pancasila University, South Jakarta, Jakarta, ¹Pharmacognosy-Phytochemistry and ²Microbiology-Biotechnology, Faculty of Pharmacy, Universitas Indonesia, Depok, ⁴Research Centre for Chemistry, Indonesian Institute of Sciences, South Tangerang, Banten, ⁶Department of Pharmaceutical Sciences, and Pharmaceutical Research and Development Laboratory of FARMAKA TROPIS, Faculty of Pharmacy, Universitas Mulawarman, Samarinda, East Kalimantan, Indonesia, ³Centre of Foundation Studies in Science, Chemistry Division, University of Malaya, Kuala Lumpur, Malaysia

J. Adv. Pharm. Technol. Res.

ABSTRACT

Isolation and determination of antibacterial compounds from plants are essential to obtain a new antibacterial as a substitute for conventional resistant antibiotics. This study aims to isolate and identify a new robustaflavone as antibacterial activity from *Garcinia latissima* Miq. leave. In this study, the isolation process was carried out using column chromatography followed by preparative thin layer chromatography (TLC) based on the TLC profile. The fraction D was tested for anti-bacterial *Bacillus subtilis* using the TLC bioautography method. The isolates obtained were then identified using ¹H-NMR, ¹³C-NMR, distortionless enhancement by polarization transfer, heteronuclear single quantum coherence, and heteronuclear multiple bond coherence. The Activity assay of the isolate was performed using the microdilution method. A pure compound obtained the result of the separation process with eluent n-hexane: Ethyl acetate (3:2) with R_f 0.6. This spot follows the spot in the contact bioautographic result of fraction D, the spot with R_f 0.6 gives an inhibition zone. After identifying and purifying the isolate were known as Robustaflavone, this compound has activity against *B. subtilis* with a (minimum inhibitory concentration) value of 2500 ppm. Robustaflavone successfully isolated and identified from *G. latissima* leave and its antibacterial activity.

Key words: Antibacterial, *Bacillus subtilis*, *Garcinia latissima* Miq, minimal inhibitory activity, Robustaflavone

INTRODUCTION

Antibiotics have an essential role in the world of health.^[1] Bacterial resistance to antibiotics makes the

disease difficult to cure.^[2] Therefore, further research needs to be done to get new antibiotics, mainly from plants. Plants are superior because they are a renewable natural resource.^[3]

Indonesia is a country that has a very high diversity of plants. The Mangosteen or Clusiaceae family (old name: Guttiferae) is one of the plant members growing in Indonesia.^[4] *Garcinia latissima* Miq. is one of the plants of the Clusiaceae tribe. However, scientific data related to this plant is still minimal and has been studied by writers, especially those from Indonesia.

Address for correspondence:

Dr. Islamudin Ahmad,
Building of Pharmaceutical Research and Development
Laboratory of FARMAKA TROPIS, 2nd Floor, Jalan Kesehatan,
Faculty of Pharmacy, Universitas Mulawarman, Samarinda,
75119 East Kalimantan, Indonesia.
E-mail: islamudinahmad@farmasi.unmul.ac.id

Submitted: 15-May-2021

Revised: 26-Jul-2021

Accepted: 09-Nov-2021

Published: 21-Jan-2022

22

Access this article online

Quick Response Code:	Website: www.japtr.org
	DOI: 10.4103/japtr.japtr_132_21

5

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

²⁴ For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Ambarwati NS, Elya B, Malik A, Omar H, Hanafi M, Ahmad I. New robustaflavone from *Garcinia latissima* Miq. leave and Its antibacterial activity. J Adv Pharm Technol Res 2022;13:50-5.

Isolation from active fractions of natural ingredients is essential to get pure active compounds as natural drugs.^[5] Ambarwati *et al.* have reported the activity assay results as antibacterial of the fractions of the ethyl acetate extract of *G. latissima* leave against *Bacillus subtilis*.^[6] *B. subtilis* is a gram-positive bacteria that can cause infection.^[7] Large amounts of *B. subtilis* infection in the intestine can cause diarrhea.^[8] Therefore, this study is needed to get active compounds as antibacterial mainly for *B. subtilis*. Moreover, there has been much antibiotic resistance increasing.

The robustaflavone biflavonoid compound has been isolated from the fruit of the *Nandina domestica* family Berberidaceae,^[9] a plant commonly used as a cough medicine and is spread in China, Japan, India, and Korea, *Rhus succedanea* and *Garcinia multiflora*.^[10] Meanwhile, the isolation, identification, and antibacterial activity of robustaflavone from *G. latissima* leave has not been reported. Robustaflavone has also been isolated from *Rhus succedanea* and *Garcinia multiflora* and shows activity against HIV-1 reverse transcriptase (RT), with an IC₅₀ value of 65 μ M.^[10]

This study aims to isolate and identify a new robustaflavone as antibacterial activity from *G. latissima* leave.

MATERIALS AND METHODS

Materials

The leaves of *G. latissima* Miq. were obtained and identified by The Center for Plant Conservation Bogor Botanic Gardens, Indonesian Institute of Sciences, Bogor, West Java, Indonesia. The voucher specimen (LFF-09-2015/FF-UI/V/2020) was stored at Laboratory of Pharmacognosy-Phytochemistry, Faculty of Pharmacy, Universitas Indonesia. The bacterial *B. subtilis* American Type Culture Collection 6633 was the collection of Microbiology Laboratory, Faculty of Pharmacy, Universitas Indonesia.

The chemical materials for extraction, fractionation, and isolation used n-hexane, ethyl acetate, methanol from PT Duta Pratama Chemika Bogor, Indonesia, and were distilled before used, aqua demineralization (PT Brataco Chemika, Indonesia), pro analysis solvents (n-hexane, ethyl acetate, methanol, chloroform, dichloromethane, acetone) from PT Smart Lab Indonesia, formic acid pro analysis (Merck, Germany), thin layer chromatography (TLC)-silica gel 60 GF₂₅₄ (Merck, Germany) for thin-layer chromatography, silica gel G60 70–230 mesh (Merck 7734.1000, Germany) and Sephadex LH-20 (Merck, Germany) for the stationary phase of column chromatography, silica gel GF₂₅₄₊₃₆₆ (Merck, Germany) for preparative-TLC. The anti-bacterial assay materials used nutrient agar (Merck, Germany), aqua bidestilata, sterilized sodium chloride 0.9%, ethanol 70%, methanol, thiazolyl blue tetrazolium bromide (BBI Life Sciences).

Instrumentation

Column chromatography equipment, a chamber for TLC, vials, and bottles for column chromatography yield collected, Ultra Violet-Visible Spectrometer (Camag, Japan), analytical balance, glasses equipment, the ultraviolet light of 254 and 366 nm, oven, refrigerator, nuclear magnetic resonance spectrophotometry (BRUKER Ascend™ 600 MHz), microplate 96-well.

Extraction and isolation process

The fraction of the ethyl acetate extract of *G. latissima* leaves, which has inhibitory activity against *B. subtilis*, was isolated using conventional column chromatography based on a chromatogram (TLC) pattern on ultraviolet light at 254 nm and 366 nm.^[11] If many spots were still detected on the chromatogram pattern, then purification was performed using column chromatography. Silica gel G60 was used as a stationary phase in column chromatography if the fraction is nonpolar, and Sephadex LH-20 was used if the fraction is semi-polar or polar.^[12] The mobile phase was determined based on the mobile phase in TLC data using a combination of n-hexane and ethyl acetate to obtain the separated spots with R_f value from 0.25 to 0.35.^[13] The LH-20 Sephadex column chromatography using the mobile phase chloroform: Methanol.^[14] The results of column chromatography were further refined using preparative-TLC (TLC-P with silica gel GF₂₅₄₊₃₆₆).^[15,16]

Bioautography assay of fraction D

TLC bioautography assay was performed using bilayer media.^[17] The agar media inoculated with *B. subtilis* ATCC 6633 poured into the agar media, which is already stable in the petri dish. The TLC plate containing the fraction spot is affixed to the bilayer layer, solidified for 1 h so that the compound contained on the TLC plate diffuses into the media. Then the TLC plate was removed, then the media was incubated at 37°C for 24 h. Inhibition zones on media showed fraction spot activity against *B. subtilis*.^[18]

Identification and structure determination of isolate

The isolate was identified with measure chemical shift signals (δ)¹ H-NMR and ¹³C-NMR using NMR Spectroscopy, (distortionless enhancement by polarization transfer [DEPT]), (heteronuclear single quantum coherence [HSQC]), and (heteronuclear multiple bond coherence [HMBC]).

Antibacterial activity assay

Antibacterial activity assay of the isolate was performed using the method of microdilution 96 well and thiazolyl blue tetrazolium bromide as an indicator on *B. subtilis* ATCC 6633, and inoculated on nutrient broth media according to some literature (1920),^[19-21] with slight modification. Briefly, a 50 μ L of isolate (20,000 ppm) were put into a well and each diluted using dimethyl sulfoxide to obtain some different concentration of 10,000 ppm, 5,000 ppm, 2,500 ppm,

1,250 ppm, 625 ppm, 312.5 ppm, 156.25 ppm, 78.13 ppm, 39.06 ppm, 19.53 ppm, and 9.77 ppm. Each well added 10 μ L of *B. subtilis* 106 colony forming unit/mL suspension and 40 μ L of nutrient broth media. Furthermore, it was incubated at 37°C for 24 h.^[1] After that, 10 μ L thiazolyl blue tetrazolium bromide 0.6 mg/mL solution and re-incubation at 37°C for 20 min.

RESULTS

Extraction, isolation, and thin layer chromatography-bioautography process

According to our previous study,^[6] The fraction D of *G. latissima* leaves ethyl acetate extract has the highest activity against *B. subtilis* compared to other fractions with its minimum inhibitory concentration (MIC) value of 312.5 ppm and the TLC profile as shown in Figure 1a. The TLC result of the isolate with mobile phase n-hexane: ethyl acetate (3:2) demonstrated in Figure 1b. Meanwhile, the Figure 1c showed that contact bioautography profile of fraction D against *B. subtilis*.

Identification of isolates

The ¹H-NMR chemical shift signals (δ) of the isolate

Table 1: ¹H-NMR dan ¹³C-NMR spectrum of isolate

Serial number	¹³ C-NMR*	δ ¹ H-NMR (m, J in Hz)
2	163.6	-
3	102.8	6.63 (s)
4	182.2	-
5-OH	161.8	13.05 (s)
6	93.9	6.45 (d, J=1.6)
7	166.4	-
8	93.90	6.23 (d, J=1.9)
9	158.8	-
10	104.2	-
1'	121.1	-
2'	131.6	8.21 (d, J=1.8)
3'	105.0	-
4'	160.5	-
5'	128.2	7.70 (d, J=8.5)
6'	127.2	7.96 (dd, J=2.3 and 8.6)
2''	163.6	-
3''	102.6	6.36 (s)
4''	182.1	-
5''	161.5	13.18 (s)
6''	104.6	-
7''	163.5	-
8''	102.8	6.71 (s)
9''	157.9	-
10''	105.0	-
1'''	123.0	-
2''' and 6'''	118.2	6.78 (2H, d, J=8.8)
3''' and 5'''	128.2	7.16 (2H, d, J=8.6)
4'''	155.2	-

CNMR: 150 MHz, H-NMR: 600 MHz

were shown in Figure 2 with tabulation in Table 1. The measurement results of chemical shift signals (δ) ¹³C-NMR and expansion of isolates showed in Figure 3a (blue) and DEPT in Figure 3b (red).

The HSQC of [Figure 4a] shows a direct correlation between δ_c 102.8 with δ_H 6.63 (s) and between δ_c 93.9 with δ_H 6.45 (d). There is a direct correlation between δ_c 131.6 with δ_H 8.21 (d) and the correlation between δ_c 128.2 with δ_H 7.70 (d). Also, there is a direct correlation between δ_c 127.2 with δ_H 7.96 (dd) and a correlation between δ_c 102.6 with δ_H 6.36 (s). The complete position and chemical shift demonstrated in Table 1. Based on the data results, the isolate was estimated as robustaflavone [in Figure 4b].




Figure 1: Thin layer chromatography profile of (a) Fraction D with mobile phase chloroform: Acetone: Formic acid (3:1:0.2); (b) Isolate (GLED2) using mobile phase n-hexane: Ethyl acetate (3:2); and (c) Contact bioautography profile of fraction D against *B. subtilis* (a) the inoculated agar medium after the thin layer chromatography plate was removed (b) inspected thin layer chromatography under UV 254 nm (c) inspected thin layer chromatography under UV 366 nm (d) Inspected thin layer chromatography under visible light. (b) shows the pure compound using eluent n-hexane and ethyl acetate (3:2), having an Rf value of 0.6. This Rf value follows the bioautography result of the fraction D extract of ethyl acetate leaves (c). In the thin layer chromatography contact bioautographic assay, the eluted and dried thin layer chromatography plate transferred to Petri plate containing the agar medium has been inoculated with *B. subtilis* for approximately one hour and then incubated for 24 hours. Inhibition zones were detected in the spot with an Rf value of 0.6 on agar medium in a petri dish ((a) of c) after incubation, and the thin layer chromatography plate was removed. This showed that the isolate was active against *B. subtilis*.




Figure 2: The proton magnetic resonance spectrum (a) and expansion (b) of isolate (Acetone- d_6 , 600 MHz)




Figure 3: Carbon magnetic resonance spectrum (a) and Carbon magnetic resonance spectrum (blue) and distortionless enhancement by polarization transfer (red) (b) of isolate




Figure 4: Heteronuclear single quantum coherence spectrum (a) and the structure design based on heteronuclear single quantum coherence spectrum (b) of isolate

Besides, it is also confirmed by HMBC spectrum data as in Figure 5a, mainly the correlation between δ_H 8.21 (d) with δ_C at δ_C 104.6 [Figure 5b], which proves the existence of a bond between the two flavonoids.




Figure 5: The heteronuclear multiple bond coherence spectrum (a) and the structure design based on the heteronuclear multiple bond coherence spectrum (b)

Antibacterial activity assay on *Bacillus subtilis*

The antibacterial activity assay against *B. subtilis* by microdilution method from this isolate (robustaflavone), with the MIC value, was 2500 ppm [as shown in Figure 6].

DISCUSSION

A total of 0.81 g of this D fraction was further purified using column chromatography with the stationary phase Sephadex LH-20 with the mobile phase of chloroform: Methanol (8: 2) and collected every 10 mL.^[22] The stationary phase of Sephadex LH-20 was used because it is lipophilic so that it was expected to be able to separate isolates from chlorophyll. the fraction component will move based on the mobile phase's movement through the stationary phase, and small molecules will be trapped in the gel matrix while the larger molecules will be outside the gel matrix and go faster.^[23] This method (Sephadex




Figure 6: Test results of minimum inhibitory concentration isolates (robustaflavones) using the microdilution method with the tetrazolium salt indicator (3-(4,5-dimethyliazol-2-yl)-2,5-diphenyltetrazolium bromide), where: minimum inhibitory concentration is minimum inhibitory concentration, MC is media control, AC is antibiotic control, GC is germ control, (-), not bacterial growth; (+) there is bacterial growth

LH-20 gel) also involves the mechanism of adsorption, partitioning, the possibility of ion exchange, and sometimes the most abundant molecule will be eluted first, and the smallest molecule will be eluted last and vice versa.^[24] This chromatography has been widely used to remove confounding pigments such as chlorophyll which tend to be larger and more lipophilic than the secondary metabolite of the plant.^[25] From this process, isolates were produced, which after further purification using TLC-partition (TLC-P) and carried out TLC obtained the chromatogram pattern.

Based on the results of ¹H-NMR measurements showed that some aromatic functional groups indicate a biflavonoid, the presence of chemical shift at 6.3H 6.63 (s) aromatic rings with 1 H, and at δ_H 6.45 (d; 1.6 Hz) and 6.23 (d, 1.9 Hz) are aromatic rings with 2H in the meta position. In chemical shift at δ_H 7.96 (dd, $J = 8.6$ and 2.3 Hz), 7.70 (d, 8.5 Hz) and 8.21 (d; 1.8 Hz) was 3H aromatic rings with ABX system. Based on this spectrum, this isolate is thought to be a flavonoid derivative. Other chemical shifts indicate the presence of 3 benzene rings at δ_H 6.71 (s); 6.36 (s) and aromatics with A2B2 systems appear at δ_H 6.78 (2H, d, 8.8 Hz) and 7.16 (2H, d, 8.6 Hz). Besides that, there is a particular group, which is in the more downfield areas at δ_H 13.05 (s) and 13.18 (s), indicating the presence of two phenol groups (-OH) that form hydrogen bonds with carbonyl groups (= C = O).

Robustaflavone [Figure 7] has been previously isolated from the bark of *Ochna schweinfurthiana* F. Hoffm,^[26] *Rhus succedanea*, and *Garcinia multiflora*.^[10] Meanwhile, this compound has also isolated from *Selaginella labordei*^[27] and *Selaginella tamariscina*.^[28]

Lin *et al.* have reported that Robustaflavone (from *Rhus succedanea*) has activity inhibiting the growth of hepatitis B virus with EC_{50} 0.25 μ M.^[29] Robustaflavone has also been reported to inhibit the action of the HIV-1 RT enzyme.^[30]

CONCLUSION

In conclusion, a biflavonoid compound (robustaflavone)




Figure 7: Robustaflavone

has been isolated from the most active fraction of ethyl acetate extract from *G. latissima*. The robustaflavone has an antibacterial activity for *B. subtilis* with a MIC value of 2500 ppm.

Acknowledgment

The authors acknowledge to gratitude "Hibah Penelitian Kolaboratif Nasional 2021 Universitas Negeri Jakarta" (Grant Number: 3/PKM/LPPM/IV/2021).

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Keawchai K, Chumkaew P, Permpoonpattana P, Srisawat T. Synergistic effect of ampicillin and dihydrobenzofuran neolignans (myticaganal C) identified from the seeds of *Myristica fragrans* Houtt. against *Escherichia coli*. *J Adv Pharm Technol Res* 2021;12:79-83.
- Ventola CL. The antibiotic resistance crisis: Part 1: Causes and threats. *P T* 2015;40:277-83.
- Perea-Moreno MA, Samerón-Manzano E, Perea-Moreno AJ. Biomass as renewable energy: Worldwide research trends. *Sustainability* 2019;11:1-19.
- Xi-Wen L, Jie L, Robson NK, Stevens PF. Clusiaceae (Guttiferae). In: *Flora of China*. Vol. 13. ST. Louis: Science Press and Missouri Botanical Garden Press; 2009. p. 1-47.
- Sasidharan S, Chen Y, Saravanan D, Sundram KM, Yoga Latha L. Extraction, isolation and characterization of bioactive compounds from plants' extracts. *Afr J Tradit Complement Altern Med* 2011;8:1-10.
- Ambarwati NS, Elya B, Nur A, Puspitasari N, Malik A, Hanafi M. Activity of fractions from *Garcinia latissima* Miq. leaves ethyl acetate extract as antibacterial against *Bacillus subtilis* and antioxidant. *Adv Sci Lett* 2018;24:6366-70.
- Mai-prochnow A, Clauson M, Hong J, Murphy AB. Gram positive and gram negative bacteria differ in their sensitivity to cold plasma. *Nat Publ Group* 2016;11:1-11.
- Kim K, He Y, Xiong X, Ehrlich A, Li X, Raybould H, et al. Dietary supplementation of *Bacillus subtilis* influenced intestinal health of weaned pigs experimentally infected with a pathogenic *E. coli*. *J Anim Sci Biotechnol* 2019;10:1-12.
- Jo A, Yoo HJ, Lee M. Robustaflavone isolated from *Nandina domestica* using bioactivity-guided fractionation downregulates inflammatory mediators. *Molecules* 2019;24:E1789.
- Lin YM, Anderson H, Flavin MT, Pai YH, Mata-Greenwood E, Pengsuparp T, et al. *In vitro* anti-HIV activity of biflavonoids isolated from *Rhus succedanea* and *Garcinia multiflora*. *J Nat Prod* 1997;60:884-8.
- Kumar S, Jyotirmayee K, Sarangi M. Thin layer chromatography: A tool of biotechnology for isolation of bioactive compound from medicinal plants. *Int J Pharm Sci Rev Res* 2013;18:126-32.
- Datta S, Zhou YD, Nagle DG. Comparative study of chromatographic medium-associated mass and potential antitumor activity loss with bioactive extracts. *J Nat Prod* 2013;76:642-7.
- Costa Fd, Garrard I, da Silva AJ, Leitão GG. Changes in the mobile phase composition on a stepwise counter-current chromatography elution for the isolation of flavonoids from *Siparuna glycyarpa*. *J Sep Sci* 2013;36:2253-9.
- Schroeder DR, Colson KL, Klohr SE, Lee MS, Matson JA, Brinen LS, et al. Pyrrolosporin A, a new antitumor antibiotic from *Micromonospora* sp. C39217-R109-7. II. Isolation, physico-chemical properties, spectroscopic study and X-ray analysis. *J Antibiot (Tokyo)* 1996;49:865-72.
- Kozyra M, Główniak K, Zabta A, Mroczek T, Cierpicki T. Column chromatography and preparative TLC for isolation and purification of coumarins from *Peucedanum verticillare* L. Koch ex DC. *J Planar Chromatogr* 2005;18:224-7.
- Gocan S. Stationary phases for thin-layer chromatography. *J Chromatogr Sci* 2002;40:538-49.
- Dewanjee S, Gangopadhyay M, Bhattacharya N, Khanra R, Dua TK. Bioautography and its scope in the field of natural product chemistry. *J Pharm Anal* 2015;5:75-84.
- Ambarwati NS, Malik A, Elya B, Hanafi M. Profile of antibacterial activity of fractions from methanol extracts of *Garcinia latissima* Miq. fruit rind. *Asian J Pharm Clin Res* 2017;10:66-8.
- Ambarwati NS, Elya B, Malik A, Hanafi M, Omar H. Isolation, characterization, and antibacterial assay of friedelin from *Garcinia latissima* Miq. leaves. *J Phys Conf Ser* 2019;1402:1-12.
- Caamal-Herrera IO, Carrillo-Cocom LM, Escalante-Réndiz DY, Aráiz-Hernández D, Azamar-Barrios JA. Antimicrobial and antiproliferative activity of essential oil, aqueous and ethanolic extracts of *Ocimum micranthum* willd leaves. *BMC Complement Altern Med* 2018;18:55.
- Biloa Messi B, Ho R, Meli Lannang A, Cressend D, Perron K, Nkengfack AE, et al. Isolation and biological activity of compounds from *Garcinia prussii*. *Pharm Biol* 2014;52:706-11.
- Kusuma SA, Parwati I, Subroto T, Rukayadi Y, Fadhilillah M, Rizaludin A. Comparison of simple and rapid extracting methods of free-tags *Mycobacterium tuberculosis* protein 64 recombinant protein from polyacrylamide gel: Electroelution and the optimized passive elution. *J Adv Pharm Technol Res* 2021;12:180-4.
- Fágáin C, Cummins PM, Connor BF. Gel-filtration chromatography. In: *Protein Chromatography: Methods and Protocols, Methods in Molecular Biology*. Berlin: Springer Science+Business Media; 2011. p. 25-33.
- Yang Y, Yuan X, Xu Y, Yu Z. Purification of anthocyanins from extracts of red raspberry using macroporous resin. *Int J Food Prop* 2015;18:1046-58.
- Agostini-Costa T da S, Vieira RF, Bizzo HR, Silveira D, Gimenes MA. Secondary metabolites. In: Dhanarasa S, editor. *Chromatography and Its Applications*. London: InTech; 2012. p. 131-64.
- Ndongo JT, Issa ME, Messi AN, Mbing JN, Cuendet M, Pegnyemb DE, et al. Cytotoxic flavonoids and other constituents from the stem bark of *Ochna schweinfurthiana*. *Nat Prod Res* 2015;29:1684-7.
- Tan WJ, Xu JC, Li L, Chen KL. Bioactive compounds of inhibiting xanthine oxidase from *Selaginella labordei*. *Nat Prod Res* 2009;23:393-8.
- Dai Y, But PP, Chu LM, Chan YP. Inhibitory effects of *Selaginella tamariscina* on immediate allergic reactions. *Am J Chin Med* 2005;33:957-66.
- Lin Y, Zembower DE, Flavin MT, Schure RM, Anderson HM, Korba BE, et al. Robustaflavone, a naturally occurring biflavanoid, is a potent non-nucleoside inhibitor of hepatitis B virus replication *in vitro*. *Bioorg Med Chem Lett* 1997;7:2325-8.
- Kurapati KR, Atluri VS, Samikkannu T, Garcia G, Nair MP. Natural products as Anti-HIV agents and role in HIV-associated neurocognitive disorders (HAND): A brief overview. *Front Microbiol* 2016;6:1-14.

ORIGINALITY REPORT

15%

SIMILARITY INDEX

10%

INTERNET SOURCES

11%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	www.ezgiakar.com Internet Source	1%
2	www.ingentaconnect.com Internet Source	1%
3	jurnal.ugm.ac.id Internet Source	1%
4	M Nur-Izzati, M Arifullah, A A Nazahatul, S Klaiklay, P Chumkaew, M Z Norhazlini, F Abdulhafiz, H Zulhazman. "Elucidation of total phenolic content and antioxidant activity in medicinal Aroid, Alocasia longiloba Miq.", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1%
5	tsukuba.repo.nii.ac.jp Internet Source	1%
6	123dok.com Internet Source	1%
7	Islamudin Ahmad, Ayun Erwina Arifianti, Aditya Sindu Sakti, Fadlina Chany Saputri,	1%

Abdul Mun'im. "Simultaneous Natural Deep Eutectic Solvent-Based Ultrasonic-Assisted Extraction of Bioactive Compounds of Cinnamon Bark and Sappan Wood as a Dipeptidyl Peptidase IV Inhibitor", *Molecules*, 2020

Publication

8

Jun Li, Lin Du, Michelle Kelly, Yu-Dong Zhou, Dale G. Nagle. "Structures and Potential Antitumor Activity of Sesterterpenes from the Marine Sponge *Hyrtios communis*", *Journal of Natural Products*, 2013

Publication

9

N S S Ambarwati, B Elya, Y Desmiaty. "Anti-elastase activity of methanolic and ethyl acetate extract from *Miq.*", *Journal of Physics: Conference Series*, 2019

Publication

10

GANAPATI SHANKARLING, Mahesh Jachak, Rupali Bhise, Ankur Chaturvedi, Vidula Kamble. "Pyrroloquinoline based Styryl Dyes Doped PMMA, PS, and PS/TiO₂ Polymer for Fluorescent Applications", *Research Square Platform LLC*, 2021

Publication

11

Michael S. Eberhart, Degao Wang, Renato N. Sampaio, Seth L. Marquard et al. "Water Photo-oxidation Initiated by Surface-Bound

1 %

1 %

1 %

1 %

Organic Chromophores", Journal of the
American Chemical Society, 2017

Publication

12

Yinggang Luo. "Novel ceramides and a new
glucoceramide from the roots of *Incarvillea
arguta*", *Lipids*, 09/2004

Publication

1 %

13

jmscr.igmpublication.org

Internet Source

1 %

14

www.freepatentsonline.com

Internet Source

1 %

15

smujo.id

Internet Source

1 %

16

Chun Feng, Qiuyan Wang, Cuifen Lu, Guichun
Yang, Zuxing Chen. "Green Synthesis of
Tetrahydrobenzo[b]Pyrans by Microwave
Assisted Multi-Component One-Pot Reactions
in PEG-400", *Combinatorial Chemistry & High
Throughput Screening*, 2012

Publication

<1 %

17

José C.J.M.D.S. Menezes, Vinícius R. Campos.
"Natural biflavonoids as potential therapeutic
agents against microbial diseases", *Science of
The Total Environment*, 2021

Publication

<1 %

18

Submitted to Macau University of Science and
Technology

<1 %

19

Motoko Watanabe, Yutaka Aoyagi, Akihiro Ohta, David E. Minnikin. "Structures of Phenolic Glycolipids from Mycobacterium Kansasii", European Journal of Biochemistry, 1997

Publication

<1 %

20

worldwidescience.org

Internet Source

<1 %

21

academicjournals.org

Internet Source

<1 %

22

docksci.com

Internet Source

<1 %

23

media.neliti.com

Internet Source

<1 %

24

Submitted to Brigham Young University

Student Paper

<1 %

25

jabonline.a2hosted.com

Internet Source

<1 %

26

pasca.unila.ac.id

Internet Source

<1 %

27

patentscope.wipo.int

Internet Source

<1 %

28

rio.tamui.edu

Internet Source

<1 %

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On

FINAL GRADE

GENERAL COMMENTS

/0

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
