

Flavonoid Compounds from Krokot Herb (*Lygodium microphyllum*) and their Antioxidant Activity against DPPH

by Hadi Kuncoro

Submission date: 30-May-2019 11:20PM (UTC-0700)

Submission ID: 1138240019

File name: 5598-21202-4-PB.pdf (281.27K)

Word count: 4623

Character count: 21684



Flavonoid Compounds from Krokot Herb (*Lygodium microphyllum*) and their Antioxidant Activity against DPPH

Hadi Kuncoro^{1,2}, Kindi Farabi¹, Laode Rijai¹, Euis Julaeha², Unang Supratman^{2,3}
& Yoshihito Shiono⁴

¹Faculty of Pharmacy, Mulawarwan University, Samarinda, 75119,
Kalimantan Timur, Indonesia

²Department of Chemistry, Faculty of Mathematics and Natural Sciences,
Universitas Padjadjaran, Jatinangor 45363, Sumedang, Indonesia

³Central Laboratory of Universitas Padjadjaran, Jatinangor 45363, Sumedang, Indonesia

⁴Department of Bioresources Engineering, Faculty of Agriculture
Yamagata University, Tsuruoka-shi, Yamagata 997-8555, Japan

E-mail: unang.supratman@unpad.ac.id

Abstract. Six flavonoid compounds, kaempferol (1), quercetin (2), acacetin (3), quercetin-3-*O*- β -D-glucopyranoside (4), kaempferol-3-*O*- β -D-glucopyranoside (5), and isorhamnetin-3-*O*- β -D-glucopyranoside (6), were isolated from the methanol extract of the *Lygodium microphyllum* herb. The chemical structures of compounds 1-6 were identified on the basis of spectroscopic evidence and comparison with previously reported spectral data. Compounds 1-6 were isolated for first time from this plant. Their antioxidant activity against DPPH (2,2-Diphenyl-1-picrylhydrazyl) was evaluated. Among the compounds, quercetin (2) showed high antioxidant activity with IC₅₀ values of 6.94 ± 0.03 μ g/mL.

Keywords: Antioxidant; DPPH; flavonoids; *Lygodium microphyllum*; *Lygodiaceae*.

1 Introduction

The fern contains many useful secondary metabolites, including flavonoids, steroids, alkaloids, phenols and triterpenoids [1]. One of the thousands of fern species is the *Lygodiaceae* family, which has only one genus *Lygodium* [2]. The *Lygodium* genus is a group of ferns that propagate around other plants and are different from other kinds of ferns in that they have fleshy rhizomes that crawl over the ground. They can only live in the open because they like sunlight. Some *Lygodium* species are invasive and can become problematic in forest areas. Their fast growth and lack of predators makes these plants dominant, displacing wildlife, threatening biodiversity, and enhancing the human-animal conflict [3]. One of the invasive species from the *Lygodium* genus is *L. microphyllum*. Phytochemical studies on the *Lygodium* genus have shown it to contain a compound with an unique structure and diverse biological activities,

such as flavonoids [4], phenolic glycoside [5], naphthoquinone [6], steroids [7], and phenylpropanoid glycoside [8].

In our continuing search for novel biologically active compounds from Indonesia *Lygodium* plants, we isolated two flavonol, quercetin and quercetin-3-*O*- β -D-glucopyranoside from the ethyl acetate extract of the *L. microphyllum* herb [9]. In further screening for antioxidant compounds from the *L. microphyllum* herb we found that the methanol extract showed a significant antioxidant activity against DPPH radical-scavenging. In this paper we report the isolation and structure elucidation of flavonoid compounds from the *L. microphyllum* herb together with their antioxidant activity against DPPH radical-scavenging.

2 Materials and Methods

2.1 General

Melting points were measured on an electrothermal melting point apparatus and were not corrected. IR spectra were measured on a Perkin-Elmer 1760X spectrophotometer, FT-IR on KBr, mass spectra were recorded with a Waters XEVO Qtof HR-TOFMS mass spectrometer and a Waters UPLC MS/MS H-Class chromatographic system. ^1H and ^{13}C NMR spectra were obtained by JEOL NMR 500 MHz, JEOL NMR ECZR 600 MHz using TMS as an internal standard. Chromatographic separation was carried out on silica gel 60 (Merck), octa desyl silane (ODS, Fuji silysia). Preparative MPLC was used with a Buchi Pump Controller C-610, Buchi Pump Modules C-605, and a SiliCycle FLH-R10030B-ISO04 Siliasep column (Butchi, Switzerland). A TLC plate filled with silica gel GF254 (Merck, 0.25 mm) and detection was obtained with the appearance of 10% H_2SO_4 in ethanol followed by heating and irradiation under ultraviolet-visible light at a wavelength of 257 and 364 nm.

2.2 Plant Material

L. microphyllum herb was collected from forest areas in Samarinda, East Kalimantan in June 2014. The plant was identified by staff at the Faculty of Forestry, University of Mulawarman, Samarinda and a sample specimen (No. 09/1117.4.1.8/LL/2011) was stored at the Faculty of Forestry, University of Mulawarman, Samarinda.

2.3 Extraction and Isolation

The *L. microphyllum* herb (3.5 kg) was extracted with methanol at room temperature for 4 days. The methanol extract was evaporated using a rotary evaporator to give a dark brown residue (133.5 g). The dark brown residue was

dissolved in 520 mL of water (4:1) and partitioned successively with *n*-hexane, ethyl acetate, and *n*-butanol. Evaporation of the solvents resulted in *n*-hexane (59 g) ethyl acetate (72 g) and *n*-butanol (54 g) extracts. All of the extracts were tested for antioxidant activity against DPPH radical scavenging and showed antioxidant activity for *n*-hexane, ethyl acetate, and *n*-butanol with IC₅₀ values of 20.36 ± 0.05, 12.36 ± 0.03 and 10.42 ± 0.03 µg/mL, respectively. The EtOAc and *n*-BuOH showed the strongest antioxidant activity and showed rich contents of flavonoid compounds using TLC analysis under UV light at wavelength 254 and 367 nm as well as with AlCl₃ as a sprayer reagent. Subsequent phytochemical analysis was therefore focused on the EtOAc and *n*-BuOH extract.

The EtOAc soluble fraction (15.50 g) was fractionated by column chromatography on silica gel 60 using a gradient of *n*-hexane-EtOAc-MeOH as eluent to give eight fractions (A–H). Fraction A (85 mg) was chromatographed on a MPLC on silica gel, eluted with a gradient of CHCl₃-MeOH (10:0–1:1), to give seven subfractions (A01–A07). Subfraction A02 (45 mg) was separated on preparative TLC silica gel GF₂₅₄, eluted with CHCl₃-EtOAc-HOAc (9:1:0.5), to give **1** (2.1 mg) and **2** (2.6 mg). Fraction B (75 mg) was chromatographed on a ODS column, eluted with H₂O-MeOH (1:1), to give five subfractions (B01–B05). Subfraction B01 (110 mg) was chromatographed on a MPLC of silica gel, eluted with a gradient of CHCl₃-MeOH (10:0–6:4), to give five subfractions (B01A–B01E). Subfraction B01B (32 mg) was separated on preparative TLC silica gel GF₂₅₄, eluted with CHCl₃-EtOAc-HOAc (9:1:0.5), to give **4** (4 mg). Subfraction B02 (25 mg) was separated on preparative TLC silica gel GF₂₅₄, eluted with CHCl₃-EtOAc-HOAc (9:0.5:0.5), to give **3** (1.8 mg). The *n*-BuOH soluble fraction (30 g) was fractionated by column chromatography on silica gel 60 using a gradient CHCl₃-MeOH to give six fractions (I–N). Fraction K (2 g) was chromatographed on a Sphadex LH-20 column, eluted with MeOH, to give five subfractions (K01–K05). Subfraction K05 (42 mg) was chromatographed on a ODS column, eluted with H₂O-MeOH (1:1), to give **5** (7.5 mg) and **6** (5.5 mg).

2.3.1 Kaempferol (1)

Yellow powder, m.p. 276–278 °C, UV (MeOH) λ_{max} nm (log ε) 353 (4.08); 266 (4.20); IR (KBr) ν_{max} 3343, 1640, 1478, 1051, 827 cm⁻¹. ¹H-NMR (acetone-*d*₆, 500 MHz): δ_H 8.17 (2H, d, *J* = 9.0 Hz, H-2' and H-6'), 7.03 (1H, d, *J* = 9.0 Hz, H-3' and H-5'), 6.54 (1H, d, *J* = 1.9 Hz, H-8), 6.26 (1H, d, *J* = 1.9 Hz, H-6); ¹³C-NMR (acetone-*d*₆, 125 MHz), see Table 1; HR- TOFMS (negative ion mode) *m/z* 285.2503 [M-H]⁻, (calcd. C₁₅H₁₀O₆, *m/z* 286.2487).

Table 1 ^{13}C -NMR Data for Compounds 1-6.

Position	1*	2*	3*	4**	5**	6**
C	δ_{C}	δ_{C}	δ_{C}	δ_{C}	δ_{C}	δ_{C}
2	157.8 (s)	157.8 (s)	160.5 (s)	157.0 (s)	159.9 (s)	158.3 (s)
3	136.7 (s)	136.8 (s)	107.6 (d)	134.1 (s)	133.2 (s)	135.3 (s)
4	176.6 (s)	176.6 (s)	175.8 (s)	178.9 (s)	177.5 (s)	180.2 (s)
5	162.3 (s)	162.4 (s)	159.6 (s)	161.3 (s)	161.2 (s)	163.0 (s)
6	99.2 (d)	99.2 (d)	96.0 (d)	100.4 (d)	98.7 (d)	99.4 (d)
7	165.1 (s)	165.0 (s)	162.3 (s)	163.2 (s)	164.2 (s)	166.9 (s)
8	94.5 (d)	94.5 (d)	95.9 (d)	94.7 (d)	93.8 (d)	94.7 (d)
9	157.8 (s)	148.4 (s)	159.4 (s)	157.6 (s)	156.2 (s)	158.0 (s)
10	104.1 (s)	104.2 (s)	106.0 (s)	102.7 (s)	103.9 (s)	105.4 (s)
1'	123.3 (s)	121.5 (s)	121.2 (s)	121.6 (s)	120.9 (s)	122.8 (s)
2'	130.5 (s)	116.2 (d)	127.6 (d)	116.0 (d)	130.9 (d)	113.9 (d)
3'	116.4 (d)	147.0 (s)	115.8 (d)	144.7 (s)	115.1 (d)	148.1 (s)
4'	162.3 (s)	145.8 (s)	160.6 (s)	148.8 (s)	156.4 (s)	150.6 (s)
5'	116.4 (d)	116.0 (d)	115.8 (d)	114.6 (d)	115.1 (d)	116.0 (d)
6'	130.5 (d)	124.5 (d)	127.6 (d)	121.7 (d)	130.9 (d)	123.8 (d)
1''	-	-	-	103.5 (d)	100.9 (d)	158.3 (s)
2''	-	-	-	74.3 (d)	74.2 (d)	135.3 (s)
3''	-	-	-	76.8 (d)	76.4 (d)	180.2 (s)
4''	-	-	-	69.8 (d)	69.9 (d)	163.0 (s)
5''	-	-	-	77.0 (d)	77.5 (d)	99.4 (d)
6''	-	-	-	61.1 (t)	60.8 (t)	166.9 (s)
4'-OCH ₃	-	-	55.8 (q)	-	-	-
5'-OCH ₃	-	-	-	-	-	56.8 (q)

*) measured in acetone-*d*₆, 125 MHz., **) measured in pyridine-*d*₆, 150 MHz.**2.3.2 Quercetin (2)**

Yellow powder, m.p. 315-317 °C, UV (MeOH) λ_{max} nm (log ϵ) 370 (3.86); 255 (3.79); IR (KBr) ν_{max} 3410, 1602, 1693 cm^{-1} . ^1H -NMR (acetone-*d*₆, 500 MHz): δ_{H} 7.73 (1H, d, $J = 2.5$ Hz, H-2'), 7.62 (1H, dd, $J = 8.5, 2.5$ Hz, H-6'), 6.87 (1H, d, $J = 8.5$ Hz, H-5'), 6.38 (1H, d, $J = 2.5$ Hz, H-8), 6.17 (1H, d, $J = 2.5$ Hz, H-6); ^{13}C -NMR (acetone-*d*₆, 125 MHz), see Table 1; ESI- TOFMS (negative ion mode) m/z 301.40 [M-H]⁻, (calcd. C₁₅H₁₀O₇, m/z 302.04).

2.3.3 Acacetin (3)

Yellow needle-like crystal. ^1H -NMR (acetone-*d*₆, 500 MHz): δ_{H} 7.80 (2H, d, $J = 7.8$ Hz, H-2', H-6'), 6.70 (2H, d, $J = 7.8$ Hz, H-3', H-5'), 6.41 (1H, d, $J = 2.5$ Hz, H-8), 6.40 (1H, d, $J = 2.5$ Hz, H-6), 6.50 (1H, s, H-3), 4.10 (3H, s, 4'-OCH₃); ^{13}C -NMR (DMSO-*d*₆, 125 MHz), see Table 1; HR-TOFMS (negative ion mode) m/z 285.0783 [M+H]⁻, (calcd. C₁₆H₁₂O₅, m/z 284.0685).

2.3.4 Quercetin-3-O- β -D-glucopyranoside (4)

Yellow powder, UV (MeOH): λ_{max} (log ϵ) 264 (3.85), 350 (3.58) nm; IR (KBr) ν_{max} 3310, 1662, 1602, 1040 cm^{-1} . $^1\text{H-NMR}$ (pyridine- d_5 , 600 MHz): δ_{H} 7.68 (1H, d, $J = 2.4$ Hz, H-2'), 7.55 (1H, dd, $J = 8.4, 2.4$ Hz, H-6'), 6.82 (1H, d, $J = 8.4$ Hz, H-5'), 6.23 (1H, d, $J = 2.4$ Hz, H-8), 6.07 (1H, d, $J = 2.4$ Hz, H-6), 5.11 (1H, d, $J = 7.2$ Hz, H-1''), 3.68 (2H, d, $J = 5.0$ Hz, H-6''), 3.46 (1H, d, $J = 9.0$ Hz, H-2''), 3.39 (1H, d, $J = 9.6$ Hz, H-3''), 3.32 (1H, d, $J = 9.6$ Hz, H-4''), 3.19 (1H, m, H-5''); $^{13}\text{C-NMR}$ (pyridine- d_5 , 150 MHz), see Table 2; HR-TOFMS (negative ion mode) m/z 463.0883 [M-H]⁻, (calcd. $\text{C}_{21}\text{H}_{20}\text{O}_{12}$, m/z 464.0955).

2.3.5 Kaempferol-3-O- β -D-glucopyranoside (5)

Yellow powder, IR (KBr) ν_{max} 3433, 2896, 1654, 1266, 839 cm^{-1} . $^1\text{H-NMR}$ (pyridine- d_5 , 600 MHz): δ_{H} 8.08 (2H, d, $J = 9.1$ Hz, H-2', H-6'), 6.88 (2H, d, $J = 9.1$ Hz, H-3', H-5'), 6.43 (1H, d, $J = 2.0$ Hz, H-8), 6.21 (1H, d, $J = 2.0$ Hz, H-6), 5.45 (1H, d, $J = 7.7$ Hz, H-1''), 4.26 (1H, dd, $J = 9.0, 7.7$ Hz, H-2''), 3.56 (1H, t, $J = 9.0$ Hz, H-3''), 3.23 (1H, t, $J = 9.0$ Hz, H-4''), 3.17 (1H, dt, $J = 4.9, 2.0$ Hz, H-5''), 3.08 (2H, d, $J = 4.9$ Hz, H-6''); $^{13}\text{C-NMR}$ (pyridine- d_5 , 150 MHz), see Table 2; HR-TOFMS (negative ion mode) m/z 447.3783 [M-H]⁻, (calcd. $\text{C}_{21}\text{H}_{20}\text{O}_{11}$, m/z 448.3769).

2.3.6 Isorhamnethin-3-O- β -D-glucopyranoside (6)

Yellow powder, UV (MeOH): λ_{max} (log ϵ) 256 (3.81), 354 (3.52) nm. $^1\text{H-NMR}$ (pyridine- d_5 , 600 MHz): δ_{H} 7.95 (1H, d, $J = 2.0$ Hz, H-2'), 7.62 (1H, dd, $J = 8.0, 2.0$ Hz, H-6'), 6.93 (1H, d, $J = 8.0$ Hz, H-5'), 6.41 (1H, d, $J = 2.0, 8.0$ Hz, H-8), 6.22 (1H, d, $J = 2.0$ Hz, H-6), 5.43 (1H, d, $J = 7.8$ Hz, H-1''), 3.97 (3H, s, 5'-OCH₃), 3.60 (2H, d, $J = 5.2$ Hz, H-6''), 3.56 (1H, t, $J = 9.0$ Hz, H-3''), 3.28 (1H, dd, $J = 9.0, 7.8$ Hz, H-2''), 3.23 (1H, t, $J = 9.0$ Hz, H-4''), 3.17 (1H, dt, $J = 9.0, 5.2$ Hz, H-5''); $^{13}\text{C-NMR}$ (pyridine- d_5 , 150 MHz), see Table 2; ESI-TOFMS (positive ion mode) m/z 479.13 [M+H]⁺, (calcd. $\text{C}_{22}\text{H}_{22}\text{O}_{12}$, m/z 478.1160).

2.4 Determination of DPPH Radical-scavenging Activity

Antioxidant activity determination was conducted according to the method described in [10] using the 2,2 diphenyl-1-picrylhydrazyl-hydrate (DPPH) radical scavenging method with L-ascorbic acid as positive control. The test samples were dissolved in DMSO and mixed with 20 mg/L of (DPPH) methanol solution to give final concentrations of 10, 50, 100, 200, 400 and 800 $\mu\text{g/mL}$. After 30 minutes at room temperature, the absorbance values were measured at 517 nm and converted into a percentage of antioxidant activity as follows:

$$\% \text{ inhibition} = \frac{\text{absorbance of control} - \text{absorbance of sample}}{\text{absorbance of control}} \quad (1)$$

The IC₅₀ value was determined by probit values and plotted against the logarithmic values of concentrations of the test samples and a linear regression curve. This is the amount of sample necessary to decrease the absorbance of DPPH by 50%. All analyses were carried out in triplicate and the results were expressed as the mean ± standard deviation (SD) and compared using the Waller-Duncan test. A value of $p < 0.05$ was considered statistically significant.

3 Results and Discussions

The phytochemical test for the EtOAc and *n*-BuOH extract showed the presence of flavonoids. By using DPPH radical scavenging assay to guide separations, the EtOAc and *n*-BuOH fractions were separated by vacuum liquid chromatography over silica gel by gradient elution. The fractions were repeatedly subjected to MPLC, reverse-phase column chromatography, Sephadex LH-20 and preparative TLC on silica gel GF₂₅₄ and yielded six antioxidant flavonoid compounds **1-6** (Figure 1).




Figure 1 Structures of compounds 1-6.

3.1 Compound 1

This was obtained as a yellow powder. The HR-TOFMS spectrum showed [M-H]⁻ m/z 285.2503 (calcd. m/z 286.2487), which corresponds to the molecular formula of C₁₅H₁₀O₆ with eleven degrees of unsaturation. The UV (MeOH) λ_{max}

nm ($\log \epsilon$) 353 (4.08), 266 (4.20), suggests the presence of flavonoid compound. The IR spectra showed absorption peaks at 3343 cm^{-1} (OH), 1640 cm^{-1} (conjugated carbonyl), 1478 cm^{-1} (C = C aromatic), 1051 cm^{-1} (asymmetric C-O-C stretch) and 827 cm^{-1} (substituted benzene). The $^1\text{H-NMR}$ spectrum of compound **1** shows a characteristic AA'BB' pattern of ring B at δ_{H} 8.17 (2H, d, $J = 9.0 \text{ Hz}$) and 7.03 (2H, d, $J = 9.0 \text{ Hz}$), corresponding to a *p*-hydroxyphenyl group. The resonances of two *meta*-coupling protons of ring A were observed at δ_{H} 6.26 (1H, d, $J = 1.9 \text{ Hz}$) and δ_{H} 6.54 (1H, d, $J = 1.9 \text{ Hz}$). The $^{13}\text{C-NMR}$ and DEPT 135° spectra showed the presence of six sp^2 methines, nine sp^2 quaternary carbons and a ketone group, C-4, resonating at δ_{C} 176.6 ppm. These functionalities account for eight of the total 11 degrees of unsaturation, and the remaining three degrees of unsaturation are consistent with the flavonoid structure. The full assignments of $^{13}\text{C-NMR}$ data of compound **1** are shown in Table 1. Comparison of the NMR data of **1** to those of kaempferol [11], shows high similarity, consequently, compound **1** was identified as a kaempferol.

3.2 Compound 2

This was obtained as a yellow amorphous powder. The molecular formula was established to be $\text{C}_{15}\text{H}_{10}\text{O}_7$ from its ESI-TOFMS spectral data (m/z 301.40 [M-H] $^-$), and $^{13}\text{C-NMR}$ data (Table 1), thus requiring eleven degrees of unsaturation. The UV spectrum of **2** shows λ_{max} at 255 and 340 nm, which is typical for a flavonol structure [11]. The IR spectrum of **2** shows absorption bands corresponding to hydroxyl (3410 cm^{-1}), carbonyl (1693 cm^{-1}) and double bond aromatic (1602 cm^{-1}) groups. The $^1\text{H-NMR}$ spectrum of **1** shows the presence of *meta*-coupled aromatic protons at δ_{H} 6.17 (1H, d, $J = 2.5 \text{ Hz}$) and 6.38 (1H, d, $J = 2.5 \text{ Hz}$) corresponding to H-6 and H-8, respectively. The $^1\text{H-NMR}$ spectrum of **2** also shows the presence of an aromatic region exhibiting an ABX system at δ_{H} 7.73 (1H, d, $J = 2.5 \text{ Hz}$, H-2'), 7.62 (1H, dd, $J = 2.5, 8.5 \text{ Hz}$, H-6'), and 6.87 (1H, d, $J = 8.5 \text{ Hz}$, H-5') due to a 3', 4' disubstitution of ring B. A total of fifteen carbon signals were observed in the $^{13}\text{C-NMR}$ spectrum. These were assigned by DEPT and HMQC experiments to fourteen sp^2 carbons and a carbonyl signal at δ_{C} 176.6. The degree of unsaturation accounted for eight out of the total of eleven double bond equivalents. The remaining three degrees of unsaturation were consistent with a flavonol structure. A detailed comparison of the NMR data of **2** with those of quercetin [12] revealed that the structures of the two compounds were very similar, therefore, compound **2** was identified as a quercetin.

3.3 Compound 3

This was obtained as a yellow powder. The HRTOFMS spectrum showed $[\text{M}+\text{H}]^+ m/z$ 285.0783 (calcd. m/z 284.0685), which corresponds to the molecular formula of $\text{C}_{16}\text{H}_{12}\text{O}_5$ with eleven degrees of unsaturation. The $^1\text{H-}$

NMR spectrum of compound **3** shows a characteristic AA'BB' pattern of ring B at δ_{H} 7.80 (2H, d, $J = 8.0$ Hz) and δ_{H} 6.70 (2H, d, $J = 8.0$ Hz), corresponding to H-2'/H-6' and H-3'/H-5', respectively. The resonances of two *meta*-coupling protons of ring A were observed at δ_{H} 6.40 (1H, d, $J = 2.5$ Hz) and δ_{H} 6.41 (1H, d, $J = 2.5$ Hz). In $^1\text{H-NMR}$ also a singlet olefinic signal was observed at δ_{H} 6.50 (1H, H-3) and a methoxy group at δ_{H} 4.10 (3H, 4'-OCH₃). The $^{13}\text{C-NMR}$ and DEPT spectra show the presence of fifteen sp^2 carbons consisting of seven sp^2 methines, eight sp^2 quaternary carbons and a ketone at δ_{C} 175.8 ppm. These functionalities account for eight of the total of eleven degrees of unsaturation, and the remaining three degrees of unsaturation were consistent with a flavonoid structure. The full assignments of the $^{13}\text{C-NMR}$ data of compound **3** are shown in Table 1. Comparison of the NMR data of **3** with acacetin [13] shows high similarity, consequently, compound **3** was identified as an acacetin.

3.4 Compound 4

This was obtained as a yellow powder. The HR-TOFMS spectrum shows m/z 463.0883 [M-H]⁻ (calcd. C₂₁H₂₀O₁₂, m/z 464.0955) and thus required twelve degrees of unsaturation. The ^1H and $^{13}\text{C-NMR}$ spectrum of **4** resembles that of **2** except for the presence of oxygenated carbons. A methylene signal was present, which occurred as a doublet at δ_{H} 3.68 (1H, d, $J = 5$ Hz) together with an anomeric signal proton at δ_{H} 5.11 (1H, d, $J = 7.2$ Hz) as well as four oxygenated methines, resonating at δ_{H} 3.46 (1H, d, $J = 9.0$ Hz, H-2''), 3.39 (1H, d, $J = 9.0$ Hz, H-3''), 3.32 (1H, d, $J = 9.6$ Hz, H-4''), and 3.19 (1H, m, H-5''), which is typical for a glucose moiety. In $^{13}\text{C-NMR}$ also a typical anomeric carbon resonating at δ_{C} 103.5 was observed, which can be assigned to the one sugar units of β -D-glucopyranoside. The HMBC spectrum shows correlations between H-1'' at δ_{H} 5.11 (1H, d, $J = 7.2$ Hz) with C-3 (δ_{C} 134.1 ppm), which indicate that the sugar group is attached at C-3 of the flavonoid structure. Comparing the NMR data of **4** (Table 2) with the literature data [14], compound **4** was identified as a quercetin-3-*O*- β -D-glucopyranoside.

3.5 Compound 5

This was obtained as a yellow powder. The HR-TOFMS spectrum shows m/z 447.3783 [M-H]⁻ (calcd. C₂₁H₂₀O₁₁, m/z 448.3769) and thus required twelve degrees of unsaturation. The ^1H and $^{13}\text{C-NMR}$ spectrum of **5** resembles that of **1** except for the presence of oxygenated carbons. A methylene signal was present, which occurred as a doublet at δ_{H} 3.08 (1H, d, $J = 5.2$ Hz) together with an anomeric signal proton at δ_{H} 5.45 (1H, d, $J = 7.75$ Hz) as well as four oxygenated methines, resonating at δ_{H} 4.26 (1H, dd, $J = 7.75, 9.0$ Hz, H-2''), 3.56 (1H, t, $J = 9.0$ Hz, H-3''), 3.23 (1H, t, $J = 9.0$ Hz, H-4''), and 3.17 (1H, dt, $J = 5.4, 9.0$ Hz, H-5''), which is typical for a glucose moiety.

In ^{13}C -NMR a typical anomeric carbon resonating at δ_{C} 100.9 was also observed, which can be assigned to the one sugar units of a β -D-glucopyranoside. The HMBC spectrum shows correlations between H-1'' at δ_{H} 5.45 (1H, d, $J=7.75$ Hz) with C-3 (δ_{C} 133.2 ppm), which indicate that the sugar group was attached at C-3 of flavonoid structure. Comparing the NMR data of **5** (Table 2) with the literature data [15], compound **5** was identified as kaempferol-3-*O*- β -D-glucopyranoside.

3.6 Compound 6

This was obtained as a yellow powder. The ESI-TOFMS spectrum showed m/z 479.13 $[\text{M}+\text{H}]^+$ (calcd. $\text{C}_{22}\text{H}_{22}\text{O}_{12}$, m/z 478.11) and thus required twelve degrees of unsaturation. The ^1H and ^{13}C -NMR spectrum of **6** had high similarity with **4**. Both were attached to a flavonoid skeleton as quercetin derivatives and sugar moieties in C-3, which is a β -D-glucopyranoside group. The main difference between **6** and **4** is the presence of a methoxy group in **6** attached at C-5' of the flavonoid skeleton. Comparing the NMR data of **6** (Table 2) with the literature data [16], compound **6** was identified as an isorhamnetin-3-*O*- β -D-glucopyranoside.

Table 2 Antioxidant Activity of Compounds 1-6 Against DPPH Radical-scavenging.

Compounds	IC ₅₀ ($\mu\text{g}/\text{mL}$)
1	14.78 \pm 0.07
2	6.94 \pm 0.03
3	16.18 \pm 0.025
4	82.55 \pm 0.18
5	70.48 \pm 0.025
6	65.28 \pm 0.022
L-ascorbic acid*	0.15 \pm 0.01

*) positive control

Antioxidant activity determination of compounds **1-6** was conducted according to the method described in [9] with L-ascorbic acid as positive control. The antioxidant activities of compounds **1-6** are shown in Table 2. Compounds **1-6** showed weaker activity compared with standard compound. Among the isolated compounds, compound **2** showed the strongest activity with an IC₅₀ value of 6.94 \pm 0.03 $\mu\text{g}/\text{mL}$. Previous structure-activity studies of flavonoids have pointed to the importance of the number and location of OH groups, which could be more important for antiradical efficacy [15]. The glycoside derivatives of compound **4-6** showed weaker antioxidant activity than the nonglycoside derivatives, which suggests that sugar moiety can decrease antioxidant activity only in flavonoid derivatives.

4 Conclusions

Six flavonoid compounds, kaempferol (**1**), quercetin (**2**), acacetin (**3**), quercetin-3-*O*- β -D-glucopyranoside (**4**), kaempferol-3-*O*- β -D-glucopyranoside (**5**), and isorhamnetin-3-*O*- β -D-glucopyranoside (**6**), were isolated from the methanolic extract of *Lygodium microphyllum* herb. Compound **2** showed the strongest antioxidant activity, with IC₅₀ values of 6.94 \pm 0.03 μ g/mL. Glycosilation will affect antioxidant activity in flavonoid derivatives.

Acknowledgements

The authors gratefully acknowledge the Directorate General of Higher Education, Ministry of Research, Technology and Higher Education, Indonesia for its funding through Competitive Research Grant and PKPI Sandwich Like 2015 Scholarship. Our thanks also go to Dr. Ahmad Darmawan, M.Sc and Sofa Fajriah, M.Si, Research Center for Chemistry, Indonesian Institute of Sciences, Serpong, Tangerang for the NMR measurements.

References

- [1] Zeng-Fu, L., Huil, H., Hang-Yi, Z. & Jun-Chen, Z., *Review on the Extraction of Flavonoids from Fern*, Journal of Sanming University, **25**(2), pp. 200-203, 2008.
- [2] Guo-Gang, Z., Ying-Cui, H., Hong-Xia, L., Lin-Xia, Z. & Li-Juan, C., *The Research of Lygodium*, Intechopen, Shanghai, 2012.
- [3] Zheng, X.L. & Xing, F.W., *Ethnobotanical Study on Medicinal Plants around Mt. Yinggeling, Hainan Island, China*, Journal of Ethnopharmacology, **124**(2), pp. 197-210, 2009.
- [4] Zhang, L., Yin, Z. & Ye, W., *Flavonoids from Lygodium Japonicum*, Biochemical Systematics and Ecology, **34**, pp. 885-886, 2006.
- [5] Ye, W., Fan, C., Zhang, L., Yin, Z. & Zhao, S., *A New Phenolic Glycoside from the Roots of Lygodium Japonicum*, Fitoterapia, **78**(7-8), pp. 600-601, 2007.
- [6] Chen, L., Zhang, G., He, J., Jin Guan, J., Pan, C., Mi, W. & Wang, Q., *New Naphthoquinone from the Root of Lygodium Japonicum (Thunb.) Sw.* Journal of Natural Medicine, **64**, pp. 114-116, 2010.
- [7] Zhu, L., Zhang, G., Chen, L., Wang, S., Li, P. & Li, L., *A New Ecdysteroside from Lygodium japonicum (Thunb.) Sw.* Journal of Natural Medicinal, **63**, pp. 215-219, 2009.
- [8] Katja, D.G., Sonda, A.A., Harneti, D.P., Mayanti, T. & Supratman, U., *7-hydroxy-6-metoxy cumarin (schopoletin) from the Bark of Chisocheton celebicus (Meliaceae)*, Jurnal Kimia Indonesia, **1**, pp. 10-14, 2017. (Text in Indonesian)

- [9] Duan, Y., Dai, He, Y.R., Kurihara, H., Li, Y. & Yao, X., *A New Phenylpropanoid Glucoside from the Aerial Parts of Lygodium Japonicum*, Journal of Asian Natural Products Research, **14**(3), pp. 286-292, 2012.
- [10] Lei, J., Xiao, Y., Huang, J., Tang, M. & Deng, P., *Two New Kaempferol Glycosides from Androsace Umbellata*, Helvetica Chimica Acta, **92**(7), pp. 1439-1444, 2009.
- [11] Chaturvedula, V.S.R. & Prakash, I., *Isolation of Stigmasterol and β -sitosterol from the Dichloromethane Extract of Rubus Suavissimus*, International Current Pharmaceutical Journal, **1**(9), pp. 239-242, 2012.
- [12] Mouffok, S., Haba, H., Lavaud, C., Long, C. & Benkhalel, M., *Chemical Constituents of Centaurea omphalotricha Coss. & Durieu ex Batt. & Trab.*, Records of Natural Products, **6**(3), pp. 292-295, 2012.
- [13] Gomes, R.A., Ramirez, R.R.A., Maciel, J.K.S., Agra, M.F., Souza, M.F.V. & Falcao-silva, V.S., *Phenolic Compounds from Sidastrum micranthum (A. St.-Hil.) fryxell and Evaluation of Acacetin and 7,4'-Di-O-methylisoscuteallarein as Modulator of Bacterial Drug Resistance*, Quimica Nova, **34**(8), pp. 283-286, 2011.
- [14] Islam, M., Al-Amin, M.D., Siddiqi, M.M.A., Akter, S., Haque, M.M., Sultana, N. & Chowdhury, A.S., *Isolation of Quercetin-3-O-beta-D-Glucopyranoside from the Leaves of Azadirachta Indica and Antimicrobial and Cytotoxic Screening of the Crude Extracts*, Dhaka University Journal of Science, **60**(1), pp. 11-14, 2012.
- [15] Sianturi, J., Farabi, K., Mayanti, T., Harneti, D., Darwati, Supratman, U., Awang, K. & Hayashi, H., *Antioxidant Constituents from the Bark of Aglaia eximia (Meliaceae)*, Makara Journal of Science, **20**(1), pp. 1-6, 2016.
- [16] Liu, H., Mou, Y., Zhao, J., Wang, J., Zhou, L., Wang, M., Wang, D., Han, J., Yu, Z. & Yang, F., *Flavonoids from Halostachys caspica and their Antimicrobial Antioxidant Activities*, Molecules, **15**(11), pp. 7933-7945, 2010.

Flavonoid Compounds from Krokot Herb (Lygodium microphyllum) and their Antioxidant Activity against DPPH

ORIGINALITY REPORT

0%

SIMILARITY INDEX

0%

INTERNET SOURCES

0%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%

Flavonoid Compounds from Krokot Herb (*Lygodium microphyllum*) and their Antioxidant Activity against DPPH

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
