

JURNAL

by Jurnal_pa Rudi Agung N Buat Gb_5

Submission date: 27-Jul-2019 08:39AM (UTC+0700)

Submission ID: 1155290090

File name: 2. Scopus 2013.pdf (420.12K)

Word count: 6954

Character count: 37118



Dietary organic selenium improves growth, survival and resistance to *Vibrio mimicus* in cultured marron, *Cherax cainii* (Austin, 2002)



Rudy Agung Nugroho*, Ravi Fotedar

Sustainable Aquatic Resources and Biotechnology, Department of Environment and Agriculture, School of Science, Curtin University, GPO Box U 1987, Perth, Western Australia 6845, Australia

ARTICLE INFO

Article history:

Received 4 December 2012
Received in revised form
12 March 2013
Accepted 4 April 2013
Available online 16 April 2013

Keywords:

Organic selenium
Marron
Growth and health
Vibrio mimicus

ABSTRACT

To determine the effects of dietary organic selenium (OS) supplementation on the growth performance and immune competence of marron, *Cherax cainii* (Austin, 2002), a group of marron were fed 0.2 g kg⁻¹ of Sel-Plex® supplemented basal diet and then compared with another group (control) of marron fed basal diet without any supplementation. After 90 days of feeding, final weight, average weekly gains (AWG), relative gain ratio (RGR), specific growth rate (SGR), survival, total and differential haemocyte counts (THC and DHC), were compared between the two groups. Surviving marron from each group were then divided into three sub-groups (three tanks per sub-group with seven marron per tank); (1) first sub-group was injected with 20 µL of 3.24 × 10⁶ cfu *Vibrio mimicus*; (2) the second sub-group was injected with 20 µL normal saline and (3) the third sub-group was not subjected to injection and became the control group. THC, DHC, neutral red retention time (NRRT) and *Vibrio* ranks of post-injected marron were evaluated for 96 h, at every 24-h interval. The results showed that after 90 days of feeding, final weight, AWG, RGR, SGR, survival, THC, proportion of hyaline cells of OS-fed marron were significantly higher ($P < 0.05$) than the control group, whereas proportion of granular and semigranular cells were not affected by dietary OS. After challenging with *V. mimicus*, survival rate of marron without dietary OS significantly decreased ($P < 0.05$) as compared to the control group of marron. THC of marron in all sub-groups were significantly reduced ($P < 0.05$) after the challenge. However, THC and granular cells of sub-groups fed OS were higher than other sub-groups. *Vibrio* ranks and NRRT of marron fed OS were significantly lower and slower, respectively, than marron fed without OS. These findings demonstrated the benefits of OS inclusion in the marron diet in terms of growth, health and disease resistance.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Vibrio species have been identified as serious pathogens to various aquatic animals [1–6] resulting in mass mortalities to many cultured vertebrates [2,7,8]. Among various *Vibrio* species, *Vibrio mimicus* is known to cause disease outbreaks in black tiger shrimp (*Penaeus monodon*) and red claw crayfish (*Cherax quadricarinatus*) [3,4,9,10]. Intramuscular injection of *V. mimicus* can produce virulent reaction and high mortalities in yabbies (*Cherax albidus*) [9]. To overcome high mortalities of cultured animals, some aquaculturists use antibiotics to prevent the virulent reaction of *V. mimicus* infection [11–13].

However, the increasing global demand for safe seafood and the need to preserve an eco-friendly environment, the application of

antibiotics, notorious for creating antibiotic-resistant pathogens and environmental deterioration has been questioned [14,15]. Thus, various dietary trace elements, such as organic selenium (OS) have been tested and used as an alternative to antibiotics [16–19]. Recently, there has been a surge in the use of organic forms of various trace elements to enhance the productivity of cultured aquatic animals [20–22]. OS has been tested to improve growth and resistance to Tera syndrome virus (TSV) in white shrimp (*Penaeus vannamei*). Five weeks of feeding 0.3 mg kg⁻¹ of OS to shrimp resulted in higher survival [23]. The dietary OS in channel catfish (*Ictalurus punctatus*) [17], tilapia (*Oreochromis niloticus*) [24] and hybrid striped bass (*Morone chrysops* × *Morone saxatilis*) [25] has also improved their growth and immunity. Further, OS is reported to be better absorbed, has higher bioavailability and is less toxic than inorganic selenium [17,26–28]. The dietary inclusion of OS also increases the number of total haemocytes and granular haemocytes in white shrimp [17,23,28–30]. In addition, OS as an integral part of selenomethionine and selenoprotein [23,31,32], is

* Corresponding author. Tel.: +61 413631155.

E-mail addresses: rudysatriana@yahoo.com, rudysatriana@gmail.com (R.A. Nugroho).

recognized as a constituent of an antioxidant enzyme responsible for preventing cellular damage and improving immune competence in grouper (*Epinephelus malabaricus*) [33].

The immunity related physiological responses measured by alterations in total haemocytes counts (THC), differential haemocyte counts (DHC) and *Vibrio* ranks can be used as indicators of immune competence and health status of several crustaceans [34–37], including marron, *Cherax cainii* [38]. The neutral red dye retention time (NRRT) technique has also been successfully used as a tool to evaluate the lipid membrane integrity of marron during infection [36]. However, the effect of dietary OS on the growth performance and immune competence of marron when challenged with *V. mimicus* is unknown. The aim of this experiment was to evaluate the effects of dietary OS on the growth performance, survival, various immune responses and *Vibrio* ranks in OS-fed marron when challenged with *V. mimicus*.

2. Materials and methods

2.1. Preparation of basal diet and test diet

All ingredients of basal diet and test diet, except OS were supplied by Specialty Feeds Pty. Ltd, Western Australia. The source of OS was from Sel-Plex®, which was supplied by Alltech Inc. USA. The basal diet was formulated using Feed LIVE software version 1.52 from Live Informatics Company Limited, Thailand (Table 1). Basal diet pellets (0.5 mm diameter, 1 mm length) were prepared by mixing all ingredients with approximately 1000 mL kg⁻¹ distilled water and pelletized using a mincer and then dried under direct sunlight. Dried pellets were then allowed to cool at room temperature, packed and stored in a dark room before being used as a control-basal diet. To prepare a test diet, 0.2 g kg⁻¹ of Sel-Plex® was added and mixed with the basal diet ingredients and then constituted into the pelleted form similar to the basal diet.

2.2. Culture system

The present experiment was carried out in the Curtin Aquatic Research Laboratory (CARL), Technology Park, Curtin University,

Table 1
Ingredient of basal diet (g kg⁻¹) used for the marron culture.

Ingredient	Content (g kg ⁻¹)
Fish oil ^a	32
Wheat bran	545.59
Soybean meal	101.5
Fish meal ^b	257.14
Calcium carbonate	0.2
Ascorbic acid	0.5
Betaine ^c	12
Premix ^d	1.5
Cholesterol	2.5
Wheat starch	47.07
Proximate composition	
Crude protein	27.05
Crude fat	8.02
Crude fibre	6.39
Moisture content (%)	9.01
Ash (%)	6.56
Dry matter (%)	90.98
Energy (Cal/g)	1833.249

All ingredients supplied by Specialty Feeds Pty Ltd, WA, Australia.

^a Cod liver oil.

^b Peruvian fishmeal, 56% CP.

^c Betaine anhydrous 97%.

^d Commercial vitamin and mineral premix for trout.

Western Australia. Eighteen plastic cylindrical culture tanks (800 mm diameter, 500 mm high, 250 L capacity, 70 L of freshwater in each tank) were used. Freshwater in each tank was continuously filtered using fluvial 205 filters (Hagen, USA) at a rate of approximately 2 L min⁻¹. Each tank was provided with aeration and ten PVC pipes (55 mm diameter, 150 mm length) were placed in each tank to provide shelter for marron. To maintain a constant temperature of 20 °C in the culture tanks, individual automatic heaters (Sonpar®, Model: HA-100, China) were used.

2.3. Animals

A total of 180 marron (average initial weight 3.29 ± 0.08 g), purchased from Aquatic Resource Management Pty. Ltd., Western Australia were used for 90 days feeding trial followed by a challenge test. All marron, after transportation were placed in the cylindrical experimental tanks for 1 week for acclimation to the culture conditions. During the acclimation period, the marron were fed the basal diet at a rate of 3% of body weight every two days. The marron after acclimation were randomly distributed into two groups (nine tanks per group with ten marron per tank). First group were fed the basal diet and the second group were fed 0.2 g kg⁻¹ of Sel-Plex® supplemented test diet. The marron in every tank were fed the diets at a rate of 3% of their body weight every second day. Before every feeding, uneaten food and faeces were siphoned out and sufficient freshwater was added to maintain a constant water level of 70 L in each tank. Water quality parameters, such as temperature, pH and dissolve oxygen were monitored weekly using Cyberscan pH 300, Eutech Instruments, Singapore. Nitrate, nitrite and ammonium were measured and recorded weekly using chemical test kits (Aquarium Pharmaceuticals™, Inc., USA).

2.4. Challenge test

At the end of the trial, both groups of marron were further divided into three sub-groups each (three tanks per sub-groups, seven marron per tank). Two sub-group, one from each group were injected with 20 µL of 3.24 × 10⁶ cfu *V. mimicus* stock suspension that was obtained from the Department of Agriculture, Western Australia; two sub-groups from each group were injected with 20 µL normal saline solution and; the third and final two sub-groups from each group were not subjected to injections (controlled sub-group). All injections were performed through the base of the fifth thoracic leg. All marron were then monitored for survival, THC and DHC, *Vibrio* ranks and NRRT at 0, 24, 48, 76 and 96 h post-injection time.

2.5. Data collection

2.5.1. Growth indices, survival and immune responses

Marron were measured for total weight using electronic balance (GX-4000, A&D Company, Ltd., Japan) immediately after acclimation and after 90 days of the feeding trial. The marron weights were used to measure final weight, average weekly gain (AWG) [39], relative gain rate (RGR) and specific growth rate (SGR) [40,41]. The marron survival was recorded every day and at 0, 24, 48, 72, 96 h post-challenge time and the surviving marron in each sub-group were also analyzed for THC and DHC.

To measure THC and DHC, at day 0 and 90 of feeding trial and at 0, 24, 48, 72, 96 h post-challenge, 0.2 mL of haemolymph was collected from each marron represented by each replicate from all treatments. Haemolymph from individual marron was withdrawn from the base of the fifth thoracic leg into a 23-gauge needle containing 0.2 mL solution of 1% glutaraldehyde in 0.2 M sodium cacodylate and dispensed into an Eppendorf tube [42]. Total

haemocytes were counted using a haemocytometer (Neubauer, Germany) under 100-fold magnification [44]. The haemocytes were counted in both grids and the resulting mean was used as mean THC.

$$\text{THC} = (\text{cells counted} \times \text{dilution factor} \times 1000) / \text{volume of grid} (0.1 \text{ mm}^3)$$

To calculate the DHC, one drop of the mixture of glutaraldehyde in sodium cacodylate and haemolymph was smeared onto a glass slide. After smearing and air-drying, it was fixed in 70% methanol for 10 min. The fixed smear was stained in May-Grünwald and Giemsa stains for 10 min each [43] and then mounted with a coverslip. The number and percentages of three major marron haemocyte types for each individual were counted using a minimum number of 200 cells from each slide. The DHC were then calculated by using the following equation:

$$\text{DHC} = (\text{Number of different haemocyte cell type} / \text{Total haemocyte cells counted}) \times 100$$

2.5.2. *Vibrio* ranks

Vibrio rank assessment was done using the procedure used by Sang et al. and Hauton et al. [36,44]. 0.1 mL of haemolymph was withdrawn into sterile syringe and then smeared onto a nutrient agar plate. The plate was then inverted and placed in an incubator at 25 °C for 24 h. Each plate was examined for colony forming units (cfu) and cfu mL⁻¹ were counted based on the total volume of 0.1 mL plate⁻¹. The cfu mL⁻¹ was ranked 1 (1–399 cfu mL⁻¹) to 10 (3600–3999 cfu mL⁻¹). A final rank of 11 was assigned as too numerous for an accurate count.

2.5.3. Neutral red retention time assay

Neutral red dye retention time was evaluated using assay based on previous protocol [45]. To prepare a stock solution, 10 mg of neutral red dye powder was dissolved in 1 mL of dimethyl sulphoxide. A working solution (dye concentration 0.02 mg mL⁻¹) was prepared by mixing 10 mL of stock solution and then diluted with 5 mL of saline water. 0.2 mL of marron haemolymph sample was transferred into an Eppendorf tube containing 0.2 mL saline water and gently mixed. The mixture of haemolymph sample was placed onto a microscope slide treated with a poly-L-lysine solution to enhance cell adhesion. The slide was immediately placed in a 10 °C incubator for 15 min to allow the haemocytes to attach to the slide. The slide was removed from the incubator and the excess haemolymph was removed. A 40 mL of neutral red working solution was added to the slide and then covered with a coverslip. The slide was then returned to the incubator. Every 15 min the slide was taken out and the sample was examined using a microscope. The time at which 50% of the haemocytes had started to lose dye from their lysosomes was recorded as the neutral red retention time of the marron lysosomal membrane.

2.6. Statistical analysis

All data were represented as mean ± standard error (SE). A student *t*-test was performed to compare the growth indices, survival and immune responses of marron between two treatment groups. Percent data of survival were normalized using an arcsine transformation before performing significant differences analysis. Multiple comparison and post hoc test (Tukey's) were performed to determine significant differences of survival, immune responses including THC, DHC, *Vibrio* ranks and NRRT after being challenge with *V. mimicus*. All statistical analysis were made using SPSS for

Microsoft software version 18 (SPSS, Inc., USA). Significance at *P* < 0.05 was used.

3. Results

3.1. Growth indices, survival and immune responses

Growth indices, survival and immune response parameters of the marron fed two different diets are presented in Table 2. After 90 days of feeding, final weight, AWG, RGR, SGR and survival were significantly higher (*T*-test, *P* < 0.05) in marron fed dietary OS than marron fed control diet. THC, percentage hyaline cells of marron fed dietary OS were significantly higher (*P* < 0.05) than control group, whereas the proportion of granular and semigranular haemocytes of marron were not affected by the dietary OS.

After being challenged with *V. mimicus*, survival rate of marron fed only the basal diet was significantly lower (*P* < 0.05) than the other marron. THC in 24 h post-challenged marron were significantly reduced (*P* < 0.05) compared to the THC before the challenge. However, after 48 h post-challenge, THC of marron fed OS supplementation were higher than any other sub-group of marron (Table 3). After 72 h of post-injection the percentage of granular cells of sub-groups with dietary OS was also higher (*P* < 0.05) than sub-groups of marron fed only basal diets, whereas the percentage of semigranular and hyaline cells, of marron fed the control diet was significantly reduced (*P* < 0.05) compared to the sub-group of marron fed the dietary OS (Fig. 1).

3.2. *Vibrio* ranks

Vibrio ranks of marron fed OS supplementation were significantly lower than marron fed without OS after being challenged with *V. mimicus* (Fig. 2). After 76-h post-injection, there was a significant decrease (*P* < 0.05) in *Vibrio* ranks in all sub-groups of marron fed OS supplementation. However, any sub-group of marron with OS in their diets showed no significant differences (*P* > 0.05) amongst each other.

3.3. Neutral red retention time (NRRT)

NRRT of all marron was significantly reduced 24 h post-challenge, respectively of dietary OS. However, the NRRT of marron fed OS was significantly longer (*P* < 0.05) than marron fed the

Table 2

Growth indices, survival and immune responses of marron after 90 days of feeding.

Parameters	Groups	
	Control	0.2 g kg ⁻¹ Sel-Plex®
Growth indices		
Final weight (g)	3.92 ± 0.05 ^a	4.20 ± 0.05 ^b
AWG (g/week)	0.049 ± 0.002 ^a	0.072 ± 0.007 ^b
RGR (%)	20.255 ± 2.26 ^a	29.69 ± 2.85 ^b
SGR (%)	0.19 ± 0.01 ^a	0.27 ± 0.02 ^b
Survival (%)	77.77 ± 3.64 ^a	94.44 ± 2.42 ^b
Immune competence		
THC (× 10 ⁶ cells/mL)	2.47 ± 0.30 ^a	3.75 ± 0.15 ^b
Granular (%)	32.66 ± 1.45 ^a	36.54 ± 1.32 ^a
Semigranular (%)	29.44 ± 1.29 ^a	29.85 ± 1.51 ^a
Hyaline (%)	37.77 ± 1.07 ^a	33.49 ± 1.79 ^b

Different alphabets (a, b) indicate significantly different means for different treatments at *P* < 0.05. AWG = average weekly gain; RGR = relative gain rate; SGR = specific growth rate. Sel-Plex® was added to basal diet as a source of organic selenium (OS).

Table 3
Total haemocyte counts (THC) and neutral red retention time (NRRT) of marron challenge with *V. mimicus*.

Parameters	Hour	Groups					
		Control diet			0.2 g kg ⁻¹ Sel-Plex®		
		BNil	BNS	BVm	OSNil	OSNs	OSVm
THC ($\times 10^6$ mL ⁻¹)	0	2.47 \pm 0.03 ^a	2.44 \pm 0.06 ^a	2.51 \pm 0.06 ^b	3.87 \pm 0.07 ^c	4.06 \pm 0.24 ^c	4.29 \pm 0.33 ^d
	24	2.61 \pm 0.14 ^a	2.04 \pm 0.09 ^b	2.21 \pm 0.03 ^b	3.81 \pm 0.04 ^c	3.69 \pm 0.04 ^c	3.52 \pm 0.13 ^d
	48	2.54 \pm 0.10 ^a	2.11 \pm 0.04 ^a	2.25 \pm 0.00 ^b	3.57 \pm 0.32 ^c	3.77 \pm 0.09 ^c	3.66 \pm 0.10 ^d
	72	2.48 \pm 0.09 ^a	2.41 \pm 0.12 ^a	2.17 \pm 0.04 ^b	3.67 \pm 0.09 ^c	3.40 \pm 0.17 ^c	4.35 \pm 0.26 ^d
	96	2.43 \pm 0.11 ^a	2.48 \pm 0.18 ^a	2.104 \pm 0.03 ^b	3.55 \pm 0.15 ^c	3.12 \pm 0.11 ^c	4.45 \pm 0.15 ^d
NRRT (min)	0	185.00 \pm 3.16 ^a	192.50 \pm 4.60 ^a	182.00 \pm 6.78 ^b	1130.50 \pm 11.40 ^c	1145.50 \pm 8.36 ^d	1125.50 \pm 5.00 ^e
	24	295.00 \pm 5.00 ^a	285.00 \pm 3.16 ^a	222.50 \pm 3.35 ^b	2122.50 \pm 7.15 ^c	279.00 \pm 2.13 ^d	245.00 \pm 3.87 ^e
	48	297.50 \pm 3.35 ^a	292.5 \pm 2.49 ^a	230.00 \pm 4.74 ^b	2122.50 \pm 6.02 ^c	2115.00 \pm 3.16 ^d	252.50 \pm 3.35 ^e
	72	287.50 \pm 2.49 ^a	295.00 \pm 3.16 ^a	233.75 \pm 3.75 ^b	2115.00 \pm 3.16 ^c	2100.00 \pm 3.16 ^d	252.50 \pm 3.35 ^e
	96	295.00 \pm 3.16 ^a	285.00 \pm 3.16 ^a	222.50 \pm 7.49 ^b	2110.00 \pm 5.00 ^c	292.50 \pm 2.49 ^d	257.50 \pm 7.15 ^e

Different alphabets (a, b, c, d, e) indicate significantly different means for different treatments at $P < 0.05$. Different numerals (1, 2) indicate significantly different means at different times at $P < 0.05$. Note: BNil = control with no injection; BNS = control with 20 μ L normal saline injection; BVm = control with 20 μ L *V. mimicus*; OSNil = 0.2 g kg⁻¹ Sel-Plex® supplementation with no challenge; OSNS = 0.2 g kg⁻¹ Sel-Plex® supplementation with 20 μ L normal saline injection; OSVm = 0.2 g kg⁻¹ Sel-Plex® supplementation with 20 μ L *V. mimicus* injection.




Fig. 1. The comparison of differential haemocyte counts (DHC) in the haemolymph of marron. Note: BNil = control with no injection; BNS = control with 20 μ L normal saline injection; BVm = control with 20 μ L *V. mimicus*; OSNil = 0.2 g kg⁻¹ Sel-Plex® supplementation with no challenge; OSNS = 0.2 g kg⁻¹ Sel-Plex® supplementation with 20 μ L normal saline injection; OSVm = 0.2 g kg⁻¹ Sel-Plex® supplementation with 20 μ L *V. mimicus* injection. * = Significantly difference at $P < 0.05$.




Fig. 2. Mean \pm SE *Vibrio* ranks of marron after being challenge with *V. mimicus*. Different alphabets (a, b, c) over bars indicate significantly different means for different treatments at $P < 0.05$. Different numerals (1, 2, 3) over bars indicate significantly different means at different times at $P < 0.05$. Note: BNil = control with no injection; BNS = control with 20 μ L normal saline injection; BVm = control with 20 μ L *V. mimicus*; OSNIl = 0.2 g kg^{-1} Sel-Plex[®] supplementation with no challenge; OSNS = 0.2 g kg^{-1} Sel-Plex[®] supplementation with 20 μ L normal saline injection; OSVm = 0.2 g kg^{-1} Sel-Plex[®] supplementation with 20 μ L *V. mimicus* injection.

control diet and continued to remain significantly longer even at 96 h post-challenge (Table 3).

4. Discussion

Micronutrient, such as selenium (Se), plays a pivotal role in improving aquaculture productivity [16,19,46] and in its organic form has proven to enhance the growth and survival [47–49] of aquatic animals. The weight gains of rainbow trout (*Oncorhynchus mykiss*) [27], hybrid striped bass (*M. chrysops* \times *M. saxatilis*) [50] and juvenile grouper (*E. malabaricus*) [47] respond positively to dietary OS supplementations. Current study showed that the application of dietary supplementation of Sel-Plex[®] as a source of OS, can significantly improve the growth and survival of marron. Sel-Plex[®] is also known as seleno-yeast that contains selenoprotein. It is a baker's yeast dried product, derived from *Saccharomyces cerevisiae* strain CNCM I-3060, cultivated in a Se-enriched fermentation medium to provide a high level of selenomethionine [51]. Selenomethionine may be incorporated into proteins in place of methionine or be metabolized to selenocysteine [30,52,53]. The present results also showed that 0.2 g kg^{-1} of dietary Sel-Plex[®] can significantly improve marron's survival irrespective of being challenged with *V. mimicus*. Similarly, improved survival of Taura Syndrome Virus (TSV)-infected shrimp (*P. vannamei*) fed Sel-Plex[®] as a source of OS has also been reported [23].

One kg of Sel-Plex[®] approximately contains 2 g of OS mainly represented by selenomethionine and has high bioavailability and appears to be 90% absorbed [50,54]. Following absorption, selenomethionine is metabolized to other forms of selenium, such as hydrogen selenide, which is the key metabolite derived from the inorganic form of selenium, selenite or selenate, and/or is diverted into pathways of methionine metabolism and finally stored as selenoprotein. Active selenoprotein as a type 1 iodothyronine 5'-deiodinase interacts with iodine and prevents abnormal hormone metabolism [55,56], which can be reflected in higher growth. In addition, OS can be deposited in muscle tissues longer than inorganic selenium and retained in muscles and hepatopancreas as selenoprotein for about three years. OS is extensively utilized and re-utilized to maintain *status quo* of selenium in animals to sustain growth performance and boost immune competence [29,57].

The number of THC decreases due to various stressors including pathogen infections [36,58]. The decrease in THC is related to defence activities of haemolymph and haemolymph lysis [36,59,60]. The present study indicated that THC of both controlled sub-groups and infected-sub-groups were significantly reduced after getting infected with *V. mimicus*. However, the marron fed OS supplementation were healthier as shown by their higher number of THC following 24 h post-challenge. OS supplementation in the diet stabilizes the proportion of circulating granular cells which play an important role in defence against bacterial infection [61] of marron through their phagocytic activities. Past research has shown that animals with better phagocytic activity and clearance efficiency have higher disease resistance [36,62–64]. In this study, the capability to reduce invasive pathogen, *V. mimicus*, were significantly increased following OS supplementation in the diet, which in turn led to increased resistance against *V. mimicus* [21,65].

The underlying mechanism(s) whereby dietary OS boosts the resistance of marron against *V. mimicus* is not properly understood. However, Alina et al. [66] stated that selenium enriched diet is assimilated into enzymes, such as antioxidant and protein which are important in improving immunity. Selenium, as an active agent plays a role in protecting cell compartments and cell membranes against lipid peroxidation due to pathogen infection [67] and promote antioxidant activity in the body via glutathione peroxides (GPX), a selenium-dependent enzyme which is a primary antioxidant enzyme for cellular defence against oxidative stress [46]. The inclusion of OS in the diet can increase the level of glutathione peroxides (GSH-Px), a main antioxidant enzyme that prevents cellular damage from free radicals [23]. GSH-Px is also associated with increasing cellular membrane stability and is linked with phospholipids hydroperoxide (PHGSH-Px), associated with the plasma membrane. In Addition, GSH-Px plays a main role in the protection of biological membrane integrity, especially during bacterial infection [68]. During bacterial or viral infection in the haemolymph, the level of lipid peroxidation is increased, due to the increased oxidative stress and induced peroxidation of membrane lipids. The increased level of lipid peroxidation can lead to a decreased membrane fluidity and membrane disorganization [69]. A study of amphipods (*Gammarus locusta*) has showed that the high level of lipid peroxidation is triggered by decreasing antioxidant enzyme activity [70]. Thus, it is possible that adding selenium in the diet may induce the antioxidant activity, in order to reduce the lipid peroxidation and enhance the lysosomal membrane stability during the bacterial invasion.

It is widely accepted that haemolymph of crustacean is the main internal defence against pathogens [71–73]. Thus, the number of bacteria in the haemolymph can be used as an indicator to evaluate the health of the animal. A low number of bacteraemia levels in the haemolymph indicates an improvement in the immune system, health status and possibly decreased susceptibility to infections [42]. Current results showed that the marron fed OS in the diet pre and post-24 h challenge with *V. mimicus*, had lower *Vibrio* ranks than marron fed without OS. Similarly, low levels of bacteraemia in the haemolymph were also reported in Bio-Mos[®]-fed infected marron [36] and western king prawn (*Penaeus latisulcatus*) fed a combination of two probiotics, *Pseudomonas synxantha* and *Pseudomonas aeruginosa* [74,75].

Bacterial infections may alter the stability of the lysosomal membrane of marron and Chinese shrimp (*Fenneropenaeus chinensis*) and can be evaluated by using the neutral red retention time (NRRT) [36,76]. The unhealthy cells, caused by decreasing lipid membrane integrity due to bacterial infection lose neutral red dye at a faster rate than healthy cells. In this study, lysosomal membrane integrity was affected by the injection of *V. mimicus*, as indicated by longer NRRT on 72-h post-challenged marron fed OS. It

is possible that dietary selenium can induce the lysosomal membrane stability and reduce the lipid peroxidation [77]. A similar finding was found in marron wherein lysosomal membrane stability increased in Bio-Mos[®]-fed marron [36].

In conclusion, supplementing 0.2 g kg⁻¹ of Sel-Plex[®], which equates to approximately 0.4 mg kg⁻¹ OS in the diet of marron is recommended to enhance growth performance, survival and disease resistance against *V. mimicus*. Further research needs to be conducted to validate the effects of OS supplementation on antioxidant enzymes activity, such as glutathione peroxidase, superoxide dismutase and catalase as well as levels of lipid peroxidase [78,79] that are related to the health and immunity of marron.

References

- [1] Sudheesh PS, Xu H-S. Pathogenicity of *Vibrio parahaemolyticus* in tiger prawn *Penaeus monodon* Fabricius: possible role of extracellular proteases. *Aquaculture* 2001;196:37–46.
- [2] Won KM, Park SL. Pathogenicity of *Vibrio harveyi* to cultured marine fishes in Korea. *Aquaculture* 2008;285:8–13.
- [3] Austin B. Vibrios as causal agents of zoonoses. *Veterinary Microbiology* 2010;140:310–7.
- [4] Austin B, Austin DA. Bacterial fish pathogens: disease of farmed and wild fish. Berlin: Springer; 2007. p. 552.
- [5] Pengsuk C, Longyant S, Rukpratanporn S, Chaivithangkura P, Sridulyakul P, Sithigorngul P. Development of monoclonal antibodies for simple detection and differentiation of *Vibrio mimicus* from *V. cholerae* and *Vibrio* spp. by dot blotting. *Aquaculture* 2010;300:17–24.
- [6] Chrisolite B, Thiagarajan S, Alavandi SV, Abhilash EC, Kalaimani N, Vijayan KK, et al. Distribution of luminescent *Vibrio harveyi* and their bacteriophages in a commercial shrimp hatchery in South India. *Aquaculture* 2008;275:13–9.
- [7] Karunasagar I, Pai R, Malathi GR, Karunasagar I. Mass mortality of *Penaeus monodon* larvae due to antibiotic-resistant *Vibrio harveyi* infection. *Aquaculture* 1994;128:203–9.
- [8] Prayitno SB, Latchford JW. Experimental infections of crustaceans with luminous bacteria related to Photobacterium and *Vibrio*. Effect of salinity and pH on infectivity. *Aquaculture* 1995;132:105–12.
- [9] Eaves LE, Ketterer PJ. Mortalities in red claw crayfish *Cherax quadricarinatus* associated with systemic *Vibrio mimicus* infection. *Diseases of Aquatic Organisms* 1994;19:233–7.
- [10] Payne M, Oakey J, Owens L. The ability of two different *Vibrio* spp. bacteriophages to infect *Vibrio harveyi*, *Vibrio cholerae* and *Vibrio mimicus*. *Journal of Applied Microbiology* 2004;97:663–72.
- [11] Supriyadi H, Rukyani A. Use of chemicals in aquaculture in Asia. In: Arthur JR, Lavilla-Pitogo CR, Subasinghe RP, editors. Proceedings of the meeting on the use of chemicals in aquaculture in Asia. Tigbauan, Iloilo, Philippines: Southeast Asian Fisheries Development Center, Aquaculture Department; 1996. p. 113.
- [12] Pathak SC, Ghosh SK, Palanisamy K. Use of chemicals in aquaculture in Asia. In: Arthur JR, Lavilla-Pitogo CR, Subasinghe RP, editors. Proceedings of the meeting on the use of chemicals in aquaculture in Asia. Tigbauan, Iloilo, Philippines: Southeast Asian Fisheries Development Center, Aquaculture Department; 1996. p. 87.
- [13] Liao CI. Use of chemicals in aquaculture in Asia. In: Arthur JR, Lavilla-Pitogo CR, Subasinghe RP, editors. Proceedings of the meeting on the use of chemicals in aquaculture in Asia. Tigbauan, Iloilo, Philippines: Southeast Asian Fisheries Development Center, Aquaculture Department; 1996. p. 193.
- [14] Capone DG, Weston DP, Miller V, Shoemaker C. Antibacterial residues in marine sediments and invertebrates following chemotherapy in aquaculture. *Aquaculture* 1996;145:55–75.
- [15] Kesarodi-Watson A, Kaspar H, Lategan MJ, Gibson L. Probiotics in aquaculture: the need, principles and mechanisms of action and screening processes. *Aquaculture* 2008;274:1–14.
- [16] Chiu ST, Hsieh SL, Yeh SP, Jian SJ, Cheng W, Liu CH. The increase of immunity and disease resistance of the giant freshwater prawn, *Macrobrachium rosenbergii* by feeding with selenium enriched-diet. *Fish and Shellfish Immunology* 2010;29:623–9.
- [17] Wang C, Lovell RT. Organic selenium sources, selenomethionine and seleno-cysteine, have higher bioavailability than an inorganic selenium source, sodium selenite, in diets for channel catfish (*Ictalurus punctatus*). *Aquaculture* 1997;152:223–34.
- [18] Wang Y. Effect of probiotics on growth performance and digestive enzyme activity of the shrimp *Penaeus vannamei*. *Aquaculture* 2007;269:259–64.
- [19] Dörr AJM, Pacini N, Abete MC, Prearo M, Elia AC. Effects of a selenium-enriched diet on antioxidant response in adult crayfish (*Procambarus clarkii*). *Chemosphere* 2008;73:1090–5.
- [20] Campa-Córdova AI, Hernández-Saavedra NY, De Philippis R, Ascencio F. Generation of superoxide anion and SOD activity in haemocytes and muscle of American white shrimp (*Litopenaeus vannamei*) as a response to [beta]-glucan and sulphated polysaccharide. *Fish and Shellfish Immunology* 2002;12:353–66.
- [21] Bachère E. Anti-infectious immune effectors in marine invertebrates: potential tools for disease control in larviculture. *Aquaculture* 2003;227:427–38.
- [22] Soltanian S, Thai TQ, Dhont J, Sorgeloos P, Bossier P. The protective effect against *Vibrio campbellii* in *Artemia nauplii* by pure [beta]-glucan and isogenic yeast cells differing in [beta]-glucan and chitin content operated with a source-dependent time lag. *Fish and Shellfish Immunology* 2007;23:1003–14.
- [23] Sritunyaluksana K, Intaraprasong A, Sa-nguanrut P, Filer K, Fegan DF. Organic selenium supplementation promotes shrimp growth and disease resistance to Taura syndrome virus. *ScienceAsia* 2011;37:24–30.
- [24] Abdel-Tawwab M, Wafeek M. Response of Nile tilapia, *Oreochromis niloticus* (L.) to environmental cadmium toxicity during organic selenium supplementation. *Journal of the World Aquaculture Society* 2010;41:106–14.
- [25] Cotter PA, Craig SR, McLean E. Hyperaccumulation of selenium in hybrid striped bass: a functional food for aquaculture? *Aquaculture Nutrition* 2008;14:215–22.
- [26] Gajčević Z, Kralik G, Has-Schön E, Pavić E. Effects of organic selenium supplemented to layer diet on table egg freshness and selenium content. *Italian Journal of Animal Science* 2009;8.
- [27] KÜÇÜKBAY FZ, Yazlak H, Karaca I, Sahin N, Tuzcu M, Cakmak MN, et al. The effects of dietary organic or inorganic selenium in rainbow trout (*Oncorhynchus mykiss*) under crowding conditions. *Aquaculture Nutrition* 2009;15:569–76.
- [28] Taylor JB, Finley JW, Caton JS. Effect of the chemical form of supranutritional selenium on selenium load and selenoprotein activities in virgin, pregnant, and lactating rats. *Journal of Animal Science* 2005;83:422.
- [29] Bell JG, Cowey CB, Adron JW, Pirie BJS. Some effects of selenium deficiency on enzyme activities and indices of tissue peroxidation in Atlantic salmon parr (*Salmo salar*). *Aquaculture* 1987;65:43–54.
- [30] Schrauzer GN. The nutritional significance, metabolism and toxicology of selenomethionine. In: Steve LT, editor. *Advances in food and nutrition research*. Academic Press; 2003. p. 73–112.
- [31] Barceloux DG. Selenium. *Journal of Toxicology Clinical* 1999;37:145–72.
- [32] Zhan X, Qie Y, Wang M, Li X, Zhao R. Selenomethionine: an effective selenium source for sow to improve Se distribution, antioxidant status, and growth performance of pig offspring. *Biological Trace Element Research* 2010;1–11.
- [33] Lin YH, Shiau SY. The effects of dietary selenium on the oxidative stress of grouper, *Epinephelus malabaricus*, fed high copper. *Aquaculture* 2007;267:38–43.
- [34] Jussila J, Mannonen A. Marron (*Cherax tenuimanus*) and noble crayfish (*Astacus astacus*) hepatopancreas energy and its relationship to moisture content. *Aquaculture* 1997;149:157–61.
- [35] Sang HM, Fotedar R. Growth, survival, haemolymph osmolality and organosomatic indices of the western king prawn (*Penaeus latisulcatus* Kishinouye, 1896) reared at different salinities. *Aquaculture* 2004;234:601–14.
- [36] Sang HM, Ky IT, Fotedar R. Dietary supplementation of mannan oligosaccharide improves the immune responses and survival of marron, *Cherax tenuimanus* (Smith, 1912) when challenged with different stressors. *Fish and Shellfish Immunology* 2009;27:341–8.
- [37] Morrissey NM. Culturing the marvellous marron in Western Australia. *Freshwater Crayfish* 2002;13.
- [38] Sang HM, Fotedar R. Effects of dietary beta-1,3-glucan on the growth, survival, physiological and immune response of marron, *Cherax tenuimanus* (Smith, 1912). *Fish and Shellfish Immunology* 2010;28:957–60.
- [39] Sang HM, Fotedar R. Effects of mannan oligosaccharide dietary supplementation on performances of the tropical spiny lobsters juvenile (*Panulirus ornatus*, Fabricius 1798). *Fish and Shellfish Immunology* 2010;28:483–9.
- [40] Wang Y, Jianzhong H, Weifen L, Zirong X. Effect of different selenium source on growth performances, glutathione peroxidase activities, muscle composition and selenium concentration of allogynogenetic crucian carp (*Carassius auratus gibelio*). *Animal Feed Science and Technology* 2007;134:243–51.
- [41] Sang HM, Fotedar R, Filer K. Effects of dietary mannan oligosaccharide on survival, growth, physiological condition, and immunological responses of marron, *Cherax tenuimanus* (Smith 1912). *Journal of the World Aquaculture Society* 2011;42:230–41.
- [42] Fotedar S, Tsvetnenko E, Evans L. Effect of air exposure on the immune system of the rock lobster *Panulirus cygnus*. *Marine Freshwater Research* 2001;52.
- [43] Bancroft JD, Stevens A. Theory and practice of histological techniques. New York: Churchill Livingstone; 1977.
- [44] Hauton C, Hawkins LE, Hutchinson S. The use of the neutral red retention assay to examine the effects of temperature and salinity on haemocytes of the European flat oyster *Ostrea edulis* (L.). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 1998;119:619–23.
- [45] Hauton C, Smith VJ. In vitro cytotoxicity of crustacean immunostimulants for lobster (*Homarus gammarus*) granulocytes demonstrated using the neutral red uptake assay. *Fish and Shellfish Immunology* 2004;17:65–73.
- [46] Han D, Xie S, Liu M, Xiao X, Liu H, Zhu X, et al. The effects of dietary selenium on growth performances, oxidative stress and tissue selenium concentration of gibel carp (*Carassius auratus gibelio*). *Aquaculture Nutrition* 2011;17:741–9.
- [47] Lin YH, Shiau SY. Dietary selenium requirements of juvenile grouper, *Epinephelus malabaricus*. *Aquaculture* 2005;250:356–63.
- [48] Lorentzen M, Maaga A, Julshamn K. Effects of dietary selenite or selenomethionine on tissue selenium levels of Atlantic salmon (*Salmo salar*). *Aquaculture* 1994;121:359–67.

- [49] Wang WN, Wang AL, Zhang YJ. Effect of dietary higher level of selenium and nitrite concentration on the cellular defense response of *Penaeus vannamei*. *Aquaculture* 2006;256:558–63.
- [50] Jaramillo Jr F, Peng LJ, Gatlin Iii DM. Selenium nutrition of hybrid striped bass (*Morone chrysops* × *M. saxatilis*) bioavailability, toxicity and interaction with vitamin E. *Aquaculture Nutrition* 2009;15:160–5.
- [51] Burdock GA, Cousins RJ. Amendment to the dossier in support of the generally recognized as safe (GRAS) status of Sel-Plex as a food ingredient. In *Fusing science and compliance*. Orlando, Florida: Burdock Group Consultants; 2010:1–173.
- [52] Fox T, Heuvel EVd, Atherton C, Dainty J, Lewis D, Langford N, et al. Bioavailability of selenium from fish, yeast and selenate: a comparative study in humans using stable isotopes. *European Journal of Clinical Nutrition* 2004;58:343–9.
- [53] Esaki N, Nakamura T, Tanaka H, Soda K. Selenocysteine lyase, a novel enzyme that specifically acts on selenocysteine. *Journal of Biological Chemistry* 1982;257:4386–91.
- [54] Robouch P, Molteni R, Mitic D, Holst CV. EURL evaluation report on the analytical methods submitted in connection with the application for authorisation of a feed additive according to Regulation (EC) No 1831/2003. Geel, Belgium: European Commission Joint Research Centre; 2011.
- [55] Foster LH, Sumar S. Selenium in health and disease: a review. *Critical Reviews in Food Science and Nutrition* 1997;37:211–28.
- [56] Burk RF, Hill KE. Regulation of selenoprotein. *Annual Review of Nutrition* 1993;13:65–81.
- [57] Aguilar F, Charrondiere UR, Dusemund B, Galtier P, Gilbert J, Gott DM, et al. SCIENTIFIC OPINION: L-selenomethionine as a source of selenium added for nutritional purposes to food supplements. Scientific Opinion of the Panel on Food Additives and Nutrient Sources Added to Food. The EFSA Journal 2009;1082:1–39.
- [58] Sahul Hameed AS, Sarathi M, Sudhakaran R, Balasubramanian G, Syed Musthaq S. Quantitative assessment of apoptotic hemocytes in white spot syndrome virus (WSSV)-infected penaeid shrimp, *Penaeus monodon* and *Penaeus indicus*, by flow cytometric analysis. *Aquaculture* 2006;256:111–20.
- [59] van de Braak CBT, Botterblom MHA, Liu W, Taverne N, van der Knaap WPW, Rombout JHWM. The role of the haematopoietic tissue in haemocyte production and maturation in the black tiger shrimp (*Penaeus monodon*). *Fish and Shellfish Immunology* 2002;12:253–72.
- [60] Omori SA, Martin GG, Hose JE. Morphology of haemocyte lysis and clotting in the ridgeback prawn, *Sicyonia ingentis*. *Cell and Tissue Research* 1989;255:117–23.
- [61] Hryniewiecka-Szyfter Z, Babula A. Total haemocyte counts and Haematopoiesis in *Saduria Entomon* (Linnaeus, 1758) (Isopoda, Valvifera) from the Baltic sea infected with the yeast *Cryptococcus La Urentii* (Kufferath Skinner). *Crustaceana* 1996;69:485–93.
- [62] Ratcliffe NA, Rowley AF, Fitzgerald SW, Rhodes CP. Invertebrate immunity: basic concepts and recent advances. *International Review of Cytology* 1985;97:183–350.
- [63] Alday-Sanz V, Roque A, Turnbull JF. Clearing mechanisms of *Vibrio vulnificus* biotype I in the black tiger shrimp *Penaeus monodon*. *Diseases of Aquatic Organisms* 2002;48:91–9.
- [64] van de Braak CBT, Botterblom MHA, Huisman EA, Rombout JHWM, van der Knaap WPW. Preliminary study on haemocyte response to white spot syndrome virus infection in black tiger shrimp *Penaeus monodon*. *Diseases of Aquatic Organisms* 2002;51:49–55.
- [65] Soderhall K. Crustacean immunity. *Developmental and Comparative Immunology* 1997;21:205–9.
- [66] Alina RA, Sara A, Barbu A, Molnar F. The influence of organic selenium on the growth and survival performances of the common carp (*Cyprinus carpio* L.), Galitian and Lausitz Variety, juveniles. *Bulletin UASVM Animal Science and Biotechnologies* 2009;66:1–2.
- [67] Rameshthangam P, Ramasamy P. Antioxidant and membrane bound enzymes activity in WSSV-infected *Penaeus monodon* Fabricius. *Aquaculture* 2006;254:32–9.
- [68] Winston GW, Di Giulio RT. Prooxidant and antioxidant mechanisms in aquatic organisms. *Aquatic Toxicology* 1991;19:137–61.
- [69] Mohankumar K, Ramasamy P. White spot syndrome virus infection decreases the activity of antioxidant enzymes in *Fenneropenaeus indicus*. *Virus Research* 2006;115:69–75.
- [70] Correia AD, Costa MH, Luis OJ, Livingstone DR. Age-related changes in antioxidant enzyme activities, fatty acid composition and lipid peroxidation in whole body *Gammarus locusta* (Crustacea: Amphipoda). *Journal of Experimental Marine Biology and Ecology* 2003;289:83–101.
- [71] Itami T, Takahashi Y, Nakamura Y. Efficacy of vaccination against vibriosis in cultured kuruma prawns *Penaeus japonicus*. *Journal of Aquatic Animal Health* 1989;1:238–42.
- [72] Shiu-Nan C. Coping with diseases in shrimp farming. *Malaysia: Infosh*; 1992. p. 113–7.
- [73] Bachère E, Mialhe E, Noël D, Boulo V, Morvan A, Rodriguez J. Knowledge and research prospects in marine mollusc and crustacean immunology. *Aquaculture* 1995;132:17–32.
- [74] Hai NV, Fotedar R. Comparison of the effects of the prebiotics (Bio-Mos® and [beta]-1,3-D-glucan) and the customised probiotics (*Pseudomonas synxantha* and *P. aeruginosa*) on the culture of juvenile western king prawns (*Penaeus latisulcatus* Kishinouye, 1896). *Aquaculture* 2009;289:310–6.
- [75] Hai NV, Fotedar R, Buller N. Selection of probiotics by various inhibition test methods for use in the culture of western king prawns, *Penaeus latisulcatus* (Kishinouye). *Aquaculture* 2007;272:231–9.
- [76] Yao CL, Wu CG, Jian-Hai X, Fuhua L, Zhi-Yong W, Xuezhe H. The lysosome and lysozyme response in Chinese shrimp *Fenneropenaeus chinensis* to *Vibrio anguillarum* and laminarin stimulation. *Journal of Experimental Marine Biology and Ecology* 2008;363:124–9.
- [77] Ursini F, Bindoli A. The role of selenium peroxidases in the protection against oxidative damage of membranes. *Chemistry and Physics of Lipids* 1987;44:255–76.
- [78] Watanabe T, Kiron V, Satoh S. Trace minerals in fish nutrition. *Aquaculture* 1997;151:185–207.
- [79] Marnett LJ. Lipid peroxidation—DNA damage by malondialdehyde. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 1999;424:83–95.

JURNAL

ORIGINALITY REPORT

13%

SIMILARITY INDEX

0%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

Sang, H.M.. "Dietary supplementation of mannan oligosaccharide improves the immune responses and survival of marron, *Cherax tenuimanus* (Smith, 1912) when challenged with different stressors", *Fish and Shellfish Immunology*, 200908

Publication

7%

2

Rudy Agung Nugroho, Hetty Manurung, Firman M. Nur, Widha Prahastika. "Terminalia catappa L. extract improves survival, hematological profile and resistance to *Aeromonas hydrophila* in *Betta sp.*", *Archives of Polish Fisheries*, 2017

Publication

6%

Exclude quotes Off

Exclude bibliography On

Exclude matches < 3%