

Evaluation of forest productivity and governance on the preservation of tropical rain forests in Kalimantan using the NGWR-TS nonparametric geospatial method

by Dr. Sifriyani M.si


Submission date: 19-Dec-2019 02:27PM (UTC+0700)

Submission ID: 1236914859

File name: Kalimantan_using_the_NGWR-TS_nonparametric_geospatial_method.pdf (1,015.03K)

Word count: 3881

Character count: 22500



Evaluation of forest productivity and governance on the preservation of tropical rain forests in Kalimantan using the NGWR-TS nonparametric geospatial method

Sifriyani ^{1*}, Ruslan ², F. H. Susanty ³

¹ Department of Mathematics, Faculty of Mathematics and Natural Sciences, Mulawarman University, Samarinda INDONESIA

² Study program of physical and health education, Faculty of Teacher Training and Education, Mulawarman University, Samarinda INDONESIA

³ Dipterocarps Forest Ecosystem Research and Development Center - Research and Development Agency, Ministry of Environment and Forestry, INDONESIA

*Corresponding author: sifriyani@fmipa.unmul.ac.id

Abstract

Tropical rainforests in Kalimantan are experiencing very rapid degradation, until 2019 the area of forest in Kalimantan has continued to shrink, so it is necessary to evaluate the handling of tropical rainforests by increasing forest conservation in Kalimantan. The purpose of this study is to carry out an experimental investigation about factors that support the preservation of tropical rainforests in 56 Districts/ Cities in Kalimantan. Data that used in this study were secondary data from Forestry Agency Indonesia. The data were 56 Districts/Cities from 5 provinces in Kalimantan. Forest Data in Districts/City were influenced by spatial dependence and spatial heterogeneity. Spatial dependence and spatial heterogeneity could be overcome by using one of the spatial statistical methods, namely the NGWR-TS Nonparametric Geospatial method. The method is the development of a nonparametric spline truncated regression that considers geographical or spatial factors. The results of the study obtained the best NGWR-TS modelling using the Gaussian kernel geographical weighting function with a minimum CV was 25.15. Using NGWR-TS with order $m = 2$ and point knot 2 with a minimum value of GCV was 1.57. The model had a value of $R^2 = 94.12$, so variables that significantly affected the preservation of tropical rainforests amounting to 94.12% were the area of Protected Forests, nature reserves and conservation, Production Forests, Area of each Districts/City, Population Density, Rate Economic Growth and Regional Development Index.

Keywords: geospatial, tropical rainforests, kernel gaussian, NGWR-TS, nonparametric

Sifriyani, Ruslan, Susanty FH (2019) Evaluation of forest productivity and governance on the preservation of tropical rain forests in Kalimantan using the NGWR-TS nonparametric geospatial method. *Eurasia J Biosci* 13: 1-7.

© 2019 Sifriyani et al

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

Forest on the island of Kalimantan, Indonesia face serious problems, namely deforestation, forest degradation and widespread critical land. Based on Forest Data from the Forestry Agency it is believed that Kalimantan Forests would shrink by 75 percent by 2020 if the deforestation rate is not stopped. Of the approximately 74 million hectares of forests owned by Kalimantan, only 71% were left in 2005. While the number in 2015 shrank to 55%. If the rate of deforestation does not change, Kalimantan is believed to be losing 6 million hectares of forest by 2020, meaning only less than one third of the remaining forest area (BPS Kalimantan Timur 2019).

According to the 2012 East Kalimantan forest area data, around 14.981.978 hectares were then depreciated so that in 2013 the forest area was around 8.563.508 hectares (BPS Indonesia 2017, Dinas

Kehutanan 2013). East Kalimantan has lost a forest area of 6.418.470 hectares in one year and this value was not small and could have a negative impact if it continues. Until 2019 the area of forest in East Kalimantan has continued to shrink, so it is necessary to evaluate the handling of tropical rainforests in East Kalimantan.

Indonesia is a country that has the third largest tropical forest after Brazil and Zaire and the functions as the world's lungs. Forest functions include regulating the water system, preventing and limiting flood, erosion, and maintaining soil fertility. Forest products provide the needs of the community in general and specifically for the needs of industrial development and exports so as to support economic development. Protecting the climate atmosphere and giving a good influence,

Received: May 2019

Accepted: October 2019

Printed: December 2019




Fig. 1. Map of Distribution of Districts/City Tropical Rain Forests in Kalimantan

providing natural beauty in general and especially in the form of nature reserves, wildlife reserves, hunting parks, and tourist parks, as well as laboratories for science, education and tourism and is one of the elements of the development strategy national.

The purpose of this study is an experimental investigation of what factors support the preservation of tropical rainforests. It also aims to investigate the influence of productivity and forest management on the Preservation of Tropical Rain Forests in 56 Districts/Cities in Kalimantan based on a spatiotemporal perspective. The analysis used NGWR-TS Nonparametric Geospatial.

Based on the description above, this research works on the development of applied spatial regression and statistical methods namely the **Evaluation of Productivity and Forest Management on the Preservation of Tropical Rain Forests in Kalimantan Using the NGWR-TS Nonparametric Geospatial Method.**

DATA AND METHODOLOGY

The data used were secondary data originated from the Forestry Agency (Dinas Kehutanan 2013) in 5 provinces in Kalimantan, namely East Kalimantan, West Kalimantan, Central Kalimantan, North Kalimantan and South Kalimantan. Some other statistical data in this study were obtained from Indonesian Central Bureau of Statistics (BPS Indonesia 2017, BPS Kalimantan Barat 2019, BPS Kalimantan Selatan 2019, BPS Kalimantan Tengah 2019, BPS Kalimantan Timur 2019, BPS Kalimantan Utara 2019).

The number of data used was 56 Districts/Cities in all regions of Kalimantan and forest data in 2019 with the

response variable y is the area of tropical rain forests. Predictor variables for forest governance consisted of protected forest area (x_1), nature reserve and conservation (x_2) and total area of each District/City (x_4). Predictor variables for forest productivity factors were Production Forest (x_3) Economic Growth Rate (x_6), Regional Development Index (x_7) and Population Density (x_5).

The statistical method is a development of nonparametric spline truncated regression that considers geographical or spatial factors. Truncated Spline is a function that is built on the basis of polynomial components and truncated components, namely pieces of polynomials that have knots, which could overcome patterns of changes in data behavior. The spline truncated approach is used as a solution to solve the problem of modelling spatial data analysis where the relationship between the response variable and predictor variables does not follow a certain pattern and there are patterns that change at certain sub-intervals. The steps for Nonparametric Geospatial NGWR-TS modelling are described in additional data.

The NGWR-TS method consists of estimation of the NGWR-TS model (Sifriyani 2018b), hypothesis testing of Goodness of Fit (Sifriyani et al. 2018a), simultaneous parameter significance test and partial significance test (Sifriyani 2018). Explanation of the method is described in supplementary data.

RESULT AND DISCUSSION

Exploration of Tropical Rainforest Data in Kalimantan

Distribution maps of Districts/City tropical rainforests in Kalimantan can be seen in Fig. 1. Districts/Cities that

Table 1. Results of Analysis NGWR-TS Nonparametric Geospatial Method

Category	Methods	Number of knot point	Coefficient of Determination (R ²)	AIC	RMSE
NGWR-TS Model Criteria	NGWR-TS Orde m=1	1	89.115	-36.44	2.99
		2	84.679	-37.25	3.67
		3	87.719	-36.74	3.23
	NGWR-TS Orde m=2	1	91.982	-35.56	2.41
		2	94.122*	-34.78	1.97
		3	92.315	-35.45	2.34
Spatial Heterogeneity Test	Breusch-Pagan Test	df	p-Value	Decision	
	108.52		2.0181E-11	There is effect of spatial heterogeneity	
CV values for weigthed function	Weighted Function	Optimal Bandwidth	Cross Validation (CV)	Generalized CV (GCV)	
	Kernel Gaussian	3.124	25.153	8.63	
	Kernel Bisquare	0.105	56.996	10.05	

had the largest area of Tropical Rain Forests is Malinau District with 3.650.818 ha of tropical forest. Distribution maps based on area of Tropical rainforests show areas with a forest area of 607.489 hectare with -3.650.818 hectare having the most area.

The NGWR-TS Nonparametric Geospatial Method

The NGWR-TS Nonparametric Geospatial Method was a truncated nonparametric spline regression model in Geographically Weighted Regression. The method was the development of nonparametric regression for spatial data with locally parameter estimators for each observation location (Sifriyani 2018). In the regression model, the assumption used was error with normal distribution mean 0 and variance $\sigma^2(u_i, v_i)$ on every location (u_i, v_i) . Coordinat (u_i, v_i) is an important factor in determining the weight used to estimate the parameters of the model. Given data $(x_{1i}, x_{2i}, \dots, x_{li}, y_i)$ and correlation between $(x_{1i}, x_{2i}, \dots, x_{li})$ and y_i are assumed to follow the multivariable nonparametric regression model as follows:

$$y_i = f(x_{1i}, x_{2i}, \dots, x_{li}) + \varepsilon_i, \quad i = 1, 2, \dots, n \quad (1)$$

with y_i is response variable and $f(x_{1i}, x_{2i}, \dots, x_{li})$ is curve with unknown form of as \sum med additive.

Mathematically the form of the relationship between the response variables y_i and predictor variables $(x_{1i}, x_{2i}, \dots, x_{li})$ at the location i for the NGWR-TS Nonparametric Geospatial model could be stated as follows (Sifriyani et al. 2017):

$$y_i = \beta_0(u_i, v_i) + \sum_{p=1}^l \sum_{k=1}^m \beta_{pk}(u_i, v_i)x_{pi}^k + \sum_{p=1}^l \sum_{h=1}^r \delta_{p,m+h}(u_i, v_i)(x_{pi} - K_{ph})_+^m + \varepsilon_i \quad (2)$$

Equation (2) is a Nonparametric Geospatial Model NGWR-TS degree m with n area.

Prior to the NGWR-TS nonparametric Geospatial method stage, the first step is to check whether tropical rainforest data has a spatial effect. Testing these spatial effects using the Breusch-Pagan test aims to be used to see the spatial heterogeneity of each location. Based on the results of Table 1, data obtained from tropical rainforests had the effect of spatial heterogeneity. This

is because each region has different environmental characteristics and diverse geographical conditions.

Furthermore, it determines the coordinates of the observation location. From the two weighting functions, gaussian kernel and bisquare kernel, the best model is a weighting Kernel Gaussian function with a smaller CV value than the CV value from the Kernel Bisquare function. The analysis is summarized in Table 1.

The best model was obtained from the Nonparametric Geospatial Model NGWR-TS with order $m = 2$ and the number of vertices 2, according to the R² value of 94,122. This value explained the area of protected forest (x_1), nature reserve and conservation (x_2), production forest (x_3), Area of each District/City (x_4), population density (x_5), economic growth rate (x_6), and Regional Development Index (x_7) is 94,122% which could explain as factors that influenced the preservation of Tropical Rain Forests in Kalimantan. Nonparametric Geospatial Model NGWR-TS model, order $m = 2$ and knot point $h = 2$ given by.

$$y_i = \beta_0(u_i, v_i) + \sum_{p=1}^7 \sum_{k=1}^2 \beta_{pk}(u_i, v_i)x_{pi}^k + \sum_{p=1}^7 \sum_{h=1}^3 \delta_{p,m+h}(u_i, v_i)(x_{pi} - K_{ph})_+^m + \varepsilon_i \quad (3)$$

with number of predictor variables $p = 7$ yaitu $x_1, x_2, x_3, x_4, x_5, x_6$ and x_7 . Next step is to complete the model (3), the optimum K_{ph} knot point selection was determined based on the smallest Generalized Cross Validation (GCV) value.

The next step for the Nonparametric Geospatial Model NGWR-TS (3) is to determine the points of knots K_{ph} . This modelling K_{ph} is the point where sub-intervals change in the regression curve. Selecting knot points K_{ph} used GCV method. The smaller the GCV value, the more optimum the knots selected. GCV values for the Nonparametric Geospatial Model NGWR-TS with order $m=2$ and the number of $K_{ph}=2$ are shown in Supplementary Data. The smallest GCV value obtained was 1.5749 with optimum knot points.

$K_{11}=4.42$; $K_{12}= 4.58$; $K_{21}=6.96$; $K_{22}=7.21$; $K_{31}=1.10$; $K_{32}=1.14$; $K_{41}=23900$; $K_{42}=24751$; $K_{51}=5339.3$; $K_{52}=5529$; $K_{61}=4.77$; $K_{62}=4.92$; $K_{71}=71.695$; $K_{72}=72.059$.

It could be concluded that general NGWR-TS Nonparametric Geospatial model model is as follows.

$$\hat{\beta} = (\beta_0(u_i, v_i) + \beta_{11}(u_i, v_i)x_{1i} + \beta_{12}(u_i, v_i)x_{2i}^2 + \dots + \beta_{71}(u_i, v_i)x_{7i} + \beta_{72}(u_i, v_i)x_{7i}^2 + \delta_{13}(u_i, v_i)(x_{1i} - K_{11})_+^m + \delta_{14}(u_i, v_i)(x_{1i} - K_{12})_+^2 + \dots + \delta_{73}(u_i, v_i)(x_{7i} - K_{72})_+^2 + \varepsilon_i) \quad (4)$$

The equation (4) for tropical rainforest data for 56 Districts/Cities could be expressed in the equation as follows:

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{56} \end{bmatrix} = \begin{bmatrix} \beta_0(u_1, v_1) + \sum_{p=1}^7 \sum_{k=1}^2 \beta_{pk}(u_1, v_1) x_{pk}^k \\ \beta_0(u_2, v_2) + \sum_{p=1}^7 \sum_{k=1}^2 \beta_{pk}(u_2, v_2) x_{pk}^k \\ \vdots \\ \beta_0(u_{56}, v_{56}) + \sum_{p=1}^7 \sum_{k=1}^2 \beta_{pk}(u_{56}, v_{56}) x_{pk}^k \end{bmatrix} + \begin{bmatrix} \sum_{p=1}^7 \sum_{h=1}^2 \delta_{p,m+h}(u_1, v_1) (x_{pn} - K_{ph})_+^m \\ \sum_{p=1}^7 \sum_{h=1}^2 \delta_{p,m+h}(u_2, v_2) (x_{pn} - K_{ph})_+^m \\ \vdots \\ \sum_{p=1}^7 \sum_{h=1}^2 \delta_{p,m+h}(u_{56}, v_{56}) (x_{pn} - K_{ph})_+^m \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_{56} \end{bmatrix} \quad (5)$$

It is given modeling for a particular area, an area that has an area of the largest tropical rainforest in Kalimantan, Malinau district. The modeling would use a parameter estimator $\hat{\beta}$ and $\hat{\delta}$ NGWR-TS and the optimal knots shown in Supplementary Data.

$$y_{malinau} = 2.93 - 0.343x_1 - 2.41x_1^2 + 2.44x_2 + 0.029x_2^2 - 0.00019x_3 - 8.90x_3^2 + 0.00024x_4 + 1.57x_4^2 + 4.14x_5 + 2.91x_5^2 - 0.003x_6 - 0.0016x_6^2 + 2.33x_7 + 0.019x_7^2$$

and Truncated function is

$$y_{malinau} = -0.005(x_1 - 4.43) - 0.0061(x_1 - 4.58) + 0.0004(x_2 - 6.96) + 0.0004(x_2 - 7.21) - 0.0003(x_3 - 1.10) - 0.0003(x_3 - 1.40) + 0.94(x_4 - 239) + 102.165(x_4 - 247) + 0.00065(x_5 - 5339.3) + 0.0006(x_5 - 5529.9) - 7.29(x_6 - 4.77) - 5.75(x_6 - 4.92) + 3.40(x_7 - 7.69) + 2.38(x_7 - 72.059) \quad (6)$$

Model (6) shows that there were estimators of NGWR Geospatial regression parameters that had a positive contribution and negative contribution to the preservation of tropical rainforests. Equation (6) shown the factors that contributed positively to tropical forests in Malinau district were nature reserves and conservation (x_2), Area of each District/City (x_4), population density (x_5), Regional Development Index (x_7). Positive contributions had an explanation that with increasing x_2, x_4, x_5 and x_7 will increase the preservation of tropical rainforests in Malinau district. The value of the contribution was in accordance with the parameter estimator coefficient $\hat{\beta}$ and $\hat{\delta}$ NGWR-TS.

The negative contribution for tropical forests in Malinau district based on equation (6) was Area of Protected Forest (x_1), Production forest (x_3), Economic

growth rate (x_6). The height of protected forests and production forests will threaten the preservation of tropical rainforests in Kalimantan. The rate of economic growth increases will reduce the preservation of tropical rainforests. Predictor variables that have a large influence on tropical rainforests in Districts Malinau were protected forest area, population density and regional development index.

NGWR-TS nonparametric geospatial regression modeling produced a different parameter estimator $\hat{\beta}$ and $\hat{\delta}$ for each District/City. This shows that each predictor variable had a different relationship between one District/City and another District/City so that it was difficult to interpret. Therefore, each predictor variable was grouped into three regional groups based on the regression coefficient or parameter estimator $\hat{\beta}$ and $\hat{\delta}$ to facilitate interpretation. NGWR-TS modeling for 56 districts/cities in Kalimantan was described in supplementary data.

Mapping NGWR-TS Results for Factors that Affect Tropical Rain Forests in each District/City in Kalimantan

Estimator Parameter of NGWR-TS Geospatial Regression for each District/City had a different value from other Districts/Cities so that each region will have different modelling and different estimator results. The following was a spatial mapping based on the diversity of NGWR-TS Geospatial regression coefficient estimators.

The largest regression coefficient estimator value gave explanation that the predictor variable x also had a large influence on Tropical Rain Forests. Fig. 2 shows that the Districts /City had the largest regression coefficient estimator value for the Extent of Protection Forest (x_1) were Districts of Citywaringin Barat, Sukamara, Lamandau, Bengkayang, Kayong Utara, Ketapang, Pontianak, Singkawang, Kubu raya, Landak, Melawi, Mempawah, Sambas Sanggau dan Sintang. The highest estimator value was held by Mempawah district, which means that the Protected Forest Area of Mempawah district has a strong relationship affecting Tropical Rain Forests in Kalimantan compared to 55 other Districts / Cities.

The largest regression coefficient estimator value on natural reserve and conservation variables (x_2) were Districts of Malinau, Bulungan, Tana Tidung, Nunukan, Tarakan city, Kutai Kartanegara, Kutai Timur, Berau, Penajam Paser Utara, Balikpapan City, Samarinda City and Bontang City. The highest estimator value was held by Nunukan district which means nature sanctuary and conservation (x_2) Nunukan district has a strong relationship affecting Tropical Rain Forests in Kalimantan compared to 55 other Districts/Cities. The largest regression coefficient estimator values in the Production Forest (x_3) were Malinau, Bulungan, Tana Tidung, Nunukan, Tarakan, Bengkayang, Pontianak,




Fig. 2. Map of Spatial Diversity Estimator of NGWR-TS Parameters for Variable Tropical Rain Forests in Kalimantan

Singkawang, Kubu Raya, Mempawah, Sambah, Kutai Kartanegara, Kutai Timur, Berau, Balikpapan, Samarinda and Bontang. The highest estimator value was held by Nunukan, which means production forest (x_3) Nunukan has a strong relationship affecting Tropical Rain Forests in Kalimantan compared to 55 other Districts / Cities.

Regression coefficient estimator value in district (x_4) variables were Tanah Laut, Kotabaru, Banjar, Barito Kuala, Tapin, Hulu sungai selatan, Hulu sungai tengah, Hulu sungai utara, Tabalong, Tanah bumbu, Belangan, Banjarmasin, Banjar baru, and Kapuas. The highest

estimator value was held by Barito Selatan, which means area of each district (x_4) Districts of Barito Selatan has a strong relationship affecting Tropical Rain Forests in Kalimantan compared to 55 other Districts/Cities.

Districts/City which has the largest regression coefficient estimator value for population density variable (x_5) were Districts Banjarmasin, Citywaringin Barat, Citywaringin Timur, Kapuas, Sukamara, Lamandau, Seruyan, Katingan, Pulau Pisau, Gunung Mas, City Palangka Raya, Bengkayang, Kapuas Hulu, Kayong Utara, Ketapang, Pontianak City, Singkawang

Table 2. Test results for partial parameter significance

Groups Area	Districts/Cities	Factors affecting Tropical Rain Forests in each Districts / City
1	Bulungan district	protected forest area (x_1), nature reserve and conservation (x_2) and total area of each District / city (x_4) Predictor variables for forest productivity factors are Production Forest (x_3) Economic Growth Rate (x_6), Regional Development Index (x_7) and Population Density (x_5).
	Tanah Laut district	
	Kotabaru district	
	Hulu Sungai Selatan district	
	Banjarmasin city	
	Banjarbaru city	
	Kapas Hulu district	
	Ketapang district	
	Pontianak city	
	Singkawang city	
2	Kubu Raya district	$x_1, x_2, x_3, x_4, x_5, x_7$
	Tabalong district	
3	Balangan district	$x_1, x_2, x_3, x_4, x_5, x_7$
	Bengkayang district	
4	Murung Raya district	$x_1, x_2, x_3, x_4, x_5, x_6, x_7$
	Penajam Paser Utara district	
5	Mahakam Ulu district	x_1, x_2, x_5, x_6, x_7
	Gunung Mas district	
6	Kutai Kartanegara district	x_1, x_2, x_4, x_6
	Samarinda city	
7	Malinau district	x_2, x_4, x_5, x_6
	Tarakan city	
8	Tana Tidung district	x_1, x_2, x_5, x_6
	Bontang city	
	Sekadau district	
9	Sukamara district	x_1, x_2, x_4, x_5
	Barito Selatan district	
10	Kutai Barat district	x_1, x_2, x_6
	Kutai Timur district	
	Balikpapan city	
	Berau district	
11	Nunukan district	x_1, x_2, x_5
	Palangka Raya city	
12	Banjar district	x_5, x_6
	Tapin district	
	Kapas district	
	kotawaringin Timur district	
13	Pulang Pisau district	x_4, x_5
	Barito Kuala district	
	Hulu Sungai Tengah district	
	Hulu Sungai Utara district	
	Katingan district	
	Landak district	
	Melawi district	
	Mempawah district	
	Barito Timur district	
	Barito Utara district	
14	Tanah Bumbu district	x_4
	Lamandau district	
15	Seruyan district	x_5
	City waringin Barat district	
16	Kayong Utara district	Not significant
	Sambas district	
	Sanggau district	
	Sintang district	
	Paser district	
	Paser district	

City, Kubu Raya, Landak, Melawi, Mempah, Sambas, Sanggau, Sekadau and Sintang. The highest estimator value was held by Citywaringin Barat district, which means population density (x_5) in Citywaringin Barat district has a strong relationship affecting Tropical Rain Forests in Kalimantan compared to 55 other Districts / Cities. The Districts/City with the largest regression

coefficient estimator value on the Economic Growth Rate variable (x_6) were Districts of Tanah laut, Kotabaru, Banjar, Barito Kuala, Tapin, Hulu sungai selatan, Hulu sungai tengah, Hulu sungai utara, Tabalong, Tanah bumbu, Balangan, Barito Selatan, Barito Timur, Paser, Kutai Barat, Penajam paser utara, Banjarmasin City, Banjarbaru City and Balikpapan City. The highest estimator value was held by Kotabaru district, which means Economic growth rate (x_6) in Kotabaru district has a strong relationship affecting Tropical Rain Forests in Kalimantan compared to 55 other Districts/Cities.

Districts/City which has the largest regression coefficient estimator value in the Regional Development Index variable (x_7) in Fig. 2 were Districts of Malinau, Bulungan, Tana Tidung, Nunukan, Kotabaru, Hulu sungai selatan, Hulu sungai tengah, Hulu sungai utara, Tabalong, Tanah bumbu, Balangan, Barito utara, Barito timur, Murung raya, Paser, Kutai barat, Kutai kartanegara, Kutai timur, Berau, Penajam Paser Utara, Mahakam Ulu, Balikpapan, Tarakan, Samarinda dan Bontang. The highest estimator value was held by Malinau district, which means Regional Development Index (x_7) in Malinau district has a strong relationship affecting Tropical Rain Forests in Kalimantan compared to 55 other Districts/Cities.

The purpose of the NGWR-TS parameter significance test is to determine the factors that have a significant effect on the preservation of tropical rainforests in each District/City in Kalimantan. Statistical Test (Fotheringham et al. 1997a, 1997b) for partial testing of NGWR-TS.

Partial test using significance level $\alpha = 0.05$ obtained results of 56 regencies / Cities which were divided into 18 regional groups based on predictor variables related to tropical rainforests of each District / city. The 18 groups were shown in Table 2 and the results obtained were then mapped on the predictor variable spatial diversity map in Fig. 2.

CONCLUSION

The Nonparametric Geospatial Regression Model NGWR-TS with a Gaussian kernel weighting produces an R^2 value of 94.12. This value explained that 94, 12% of the variables of Protected Forests, nature reserves and nature conservation, Production Forests, Area of each Districts/City, Population Density, Economic Growth Rate and Regional Development Index affected the preservation of Tropical Rain Forests in Kalimantan.

ACKNOWLEDGEMENTS

The authors acknowledge The Research Center of Medicine and cosmetic from Tropical Rainforest Resources for generously supporting this project.

REFERENCES

- BPS Indonesia (2017) Statistik Produksi Kehutanan 2017, Indonesian Central Bureau of Statistics.
- BPS Kalimantan Barat (2019) Provinsi Kalimantan Barat dalam Angka, Badan Pusat Statistika Provinsi Kalimantan Barat.
- BPS Kalimantan Selatan (2019) Provinsi Kalimantan Selatan dalam Angka, Badan Pusat Statistika Provinsi Kalimantan Selatan.
- BPS Kalimantan Tengah (2019) Provinsi Kalimantan Tengah dalam Angka, Badan Pusat Statistika Provinsi Kalimantan Tengah.
- BPS Kalimantan Timur (2019) Provinsi Kalimantan Timur dalam Angka, Badan Pusat Statistika Provinsi Kalimantan Timur.
- BPS Kalimantan Utara (2019) Provinsi Kalimantan Utara dalam Angka, Badan Pusat Statistika Provinsi Kalimantan Utara.
- Dinas Kehutanan (2013) Indeks Kelola Hutan dan Lahan Daerah, Kinerja Daerah Dalam Pengelolaan Hutan dan Lahan di Indonesia.
- Fotheringham AS, Brundson C, Charlton M (2002) Geographically Weighted Regression: The Analysis of Spatially Varying Relationships. England: John Wiley & Sons Ltd.
- Fotheringham AS, Charlton M, Brundson C (1997a) Measuring Spatial Variations in Relationships with Geographically Weighted Regression. In MM Fischer and A Getis (Eds.), Recent Developments in Spatial Analysis, Springer, Berlin/New York: 60-82. https://doi.org/10.1007/978-3-662-03499-6_4
- Fotheringham AS, Charlton M, Brundson C (1997b) Two techniques for exploring non-stationarity in geographical data. Geographical Systems, 4: 59-82.
- Fotheringham AS, Pitts TC (1995) Directional Variation in Distance Decay. Environment and Planning A, 27: 715-729. <https://doi.org/10.1068/a270715>
- Sifriyani (2018) Multivariable Nonparametric Regression Truncated Spline in The Geographically Weighted Regression Models (Unpublished PhD Dissertation), Universitas Gadjah Mada, Dept of Mathematics.
- Sifriyani, Budiantara IN, Kartiko SH, Gunardi (2018a) A New Method of Hypothesis Test for Truncated Spline Nonparametric Regression Influenced by Spatial Heterogeneity and Application. Abstract and Applied Analysis. <https://doi.org/10.1155/2018/9769150>
- Sifriyani, Budiantara IN, Kartiko SH, Gunardi (2019) Evaluation of Factors Affecting Increased Unemployment in East Java Using NGWR-TS Method. International Journal Of Sciences: Basic and Applied Research, 49(1): 123-142.
- Sifriyani, Haryatmi, Budiantara IN, Gunardi (2017) Geographically Weighted Regression with Spline Approach'. Far East Journal of Mathematical Sciences, 101(6): 1183-1196. <https://doi.org/10.17654/MS101061183>
- Sifriyani, Kartiko SH, Budiantara IN, Gunardi (2018b) Development Of Nonparametric Geographically Weighted Regression Using Truncated Spline Approach. Songklanakarin Journal of Science And Technology, 40(4): 909-920.

Evaluation of forest productivity and governance on the preservation of tropical rain forests in Kalimantan using the NGWR-TS nonparametric geospatial method

ORIGINALITY REPORT

9%	6%	6%	4%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to School of Business and Management ITB Student Paper	3%
2	jurnal.fmipa.unmul.ac.id Internet Source	2%
3	imadeputrawan.wordpress.com Internet Source	2%
4	Submitted to Yeungnam University Student Paper	1%
5	www.carijasa.co.id Internet Source	1%
6	www.productiontype.com Internet Source	1%
7	resits.its.ac.id Internet Source	1%
8	Erwin Erwin, Widar Ristiyani Pusparohmana, Indah Permata Sari, Rita Hairani, Usman	1%

Usman. "Phytochemical and antioxidant activity evaluation of the bark of Tampoi (*Baccaurea macrocarpa*)", F1000Research, 2018

Publication

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

Samarinda, 19 Desember 2019

✍ Mengetahui,
Ketua Tim Validasi Artikel Ilmiah
FMIPA Unmul



Dr. Sri Wahyuningsih, M.Si
NIP. 19690413 200012 2 001