



Indian Journal of Public Health Research & Development

An International Journal

SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development
Scopus coverage years: from 2010 to 2017 Publisher:
R.K. Sharma, Institute of Medico-Legal Publications
ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine:
Public Health, Environmental and Occupational Health
CiteScore 2015-0.02
SJR 2015-0.105
SNIP 2015-0.034



Website:

www.ijphrd.com

Indian Journal of Public Health Research & Development

EXECUTIVE EDITOR

Prof Vidya Surwade

Prof Dept of Community Medicine SIMS, Hapur

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Dr. Abdul Rashid Khan B. Md Jagar Din**, (*Associate Professor*)
Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
2. **Dr. V Kumar** (*Consulting Physician*)
Mount View Hospital, Las Vegas, USA
3. **Basheer A. Al-Sum**,
Botany and Microbiology Deptt, College of Science, King Saud University,
Riyadh, Saudi Arabia
4. **Dr. Ch Vijay Kumar** (*Associate Professor*)
Public Health and Community Medicine, University of Buraimi, Oman
5. **Dr. VMC Ramaswamy** (*Senior Lecturer*)
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
6. **Kartavya J. Vyas** (*Clinical Researcher*)
Department of Deployment Health Research,
Naval Health Research Center, San Diego, CA (USA)
7. **Prof. PK Pokharel** (*Community Medicine*)
BP Koirala Institute of Health Sciences, Nepal

NATIONAL SCIENTIFIC COMMITTEE

1. **Dr. Anju Ade** (*Associate Professor*)
Navodaya Medical College, Raichur, Karnataka
2. **Dr. E. Venkata Rao** (*Associate Professor*) Community Medicine,
Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
3. **Dr. Amit K. Singh** (*Associate Professor*) Community Medicine,
VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
4. **Dr. R G Viveki** (*Professor & Head*) Community Medicine,
Belgaum Institute of Medical Sciences, Belgaum, Karnataka
5. **Dr. Santosh Kumar Mulage** (*Assistant Professor*)
Anatomy, Raichur Institute of Medical Sciences Raichur(RIMS), Karnataka
6. **Dr. Gouri Ku. Padhy** (*Associate Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Dr. Ritu Goyal** (*Associate Professor*)
Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar
8. **Dr. Anand Kalaskar** (*Associate Professor*)
Microbiology, Prathima Institute of Medical Sciences, AP
9. **Dr. Md. Amirul Hassan** (*Associate Professor*)
Community Medicine, Government Medical College, Ambedkar Nagar, UP
10. **Dr. N. Girish** (*Associate Professor*) Microbiology, VIMS&RC, Bangalore
11. **Dr. BR Hungund** (*Associate Professor*) Pathology, JNMC, Belgaum.
12. **Dr. Sartaj Ahmad** (*Assistant Professor*),
Medical Sociology, Department of Community Medicine, Swami Vivekananda Subharti
University, Meerut, Uttar Pradesh, India
13. **Dr Sumeeta Soni** (*Associate Professor*)
Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

NATIONAL EDITORIAL ADVISORY BOARD

1. **Prof. Sushanta Kumar Mishra** (Community Medicine)
GSL Medical College – Rajahmundry, Karnataka
2. **Prof. D.K. Srivastava** (*Medical Biochemistry*)
Jamia Hamdard Medical College, New Delhi
3. **Prof. M Sriharibabu** (*General Medicine*) GSL Medical College, Rajahmundry,
Andhra Pradesh
4. **Prof. Pankaj Datta** (*Principal & Prosthodontist*)
Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

5. **Prof. Samarendra Mahapatro** (*Pediatrician*)
Hi-Tech Medical College, Bhubaneswar, Orissa
6. **Dr. Abhiruchi Galhotra** (*Additional Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Prof. Deepti Pruthvi** (*Pathologist*) SS Institute of Medical Sciences &
Research Center, Davangere, Karnataka
8. **Prof. G S Meena** (*Director Professor*)
Maulana Azad Medical College, New Delhi
9. **Prof. Pradeep Khanna** (*Community Medicine*)
Post Graduate Institute of Medical Sciences, Rohtak, Haryana
10. **Dr. Sunil Mehra** (*Paediatrician & Executive Director*)
MAMTA Health Institute of Mother & Child, New Delhi
11. **Dr Shailendra Handu**, *Associate Professor*, Phrma, DM (Pharma, PGI
Chandigarh)
12. **Dr. A.C. Dhariwal**: *Directorate* of National Vector Borne Disease
Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of
India, Delhi

Print-ISSN: 0976-0245-**Electronic-ISSN:** 0976-5506, **Frequency:** Quarterly
(Four issues per volume)

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website : www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Printed, published and owned by

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Published at

Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Spatial Distribution of Dengue Haemorrhagic Fever (DHF) Vulnerability Level based on Population Density, Rainfall, Drainage Condition, Natural Water Body, and Vector Control Program in Tanjung Redeb Sub-District, District of Berau, East Kalimantan

Risva¹, Siswanto¹, Subirman¹

¹Department of Public Health, Mulawarman University, Samarinda, Indonesia

ABSTRACT

Dengue hemorrhagic fever (DHF) has still become a major health concern in the world particularly in the tropical and sub-tropical area such as South-east Asia including Indonesia. In 2010, about 1.317 mortality due to dengue out of 150.000 cases has made Indonesia become the highest case of DHF country in South-east Asia. The province of East Kalimantan represents the endemic region of DHF and contributes the highest cases in Indonesia especially in the sub-district of Tanjung Redeb, district of Berau. This study investigated the level of vulnerability of DHF at sub-district scale based on population density, rainfall, drainage condition, natural water body and vector control program. A geographical information system (GIS) software were used for mapping the spatial distribution of vulnerability area of DHF in sub-district of Tanjung Redeb for every three months period of 2014. The level of DHF vulnerability in Tanjung Redeb was fluctuate. At the first three months was medium, and continues to become high level at the second and third three months, then become medium level at the last three months during 2014-time period. This result indicates that the spatial distribution based on climate, landscape, and human factors might be used to predict the vulnerability area of DHF at the fine scale such as sub-district level. This information will in turn may help government to conduct a better strategy for controlling the transmission of DHF in each area.

Keywords: *Level of Vulnerability, DHF, GIS, Tanjung Redeb-Berau*

INTRODUCTION

Dengue has still become a major health concern in the world particularly in the tropical and sub-tropical areas such as South-East Asia including Indonesia. Dengue Hemorrhagic Fever has impact on the environment and economic in this region¹. It is estimated over 50 million people were infected annually with dengue and up to 500.000 people develop a potentially lethal complication called dengue hemorrhagic fever/dengue shock syndrome². Mosquito vector, *Aedes aegypti* are endemic to most of the tropical and sub-tropical region over the world.

Prevention and control program are the main methods of reducing the prevalence of the diseases (DHF). Hence, good strategy for the prevention and control system is needed. Several studies have been conducted in various regions over the world to study the link of environmental factors and the prevalence of dengue fever by using spatial analysis. There were approximately fifty million dengue hemorrhagic fever (DHF) infection and about 2,5 million people live in the dengue endemic regions³. Dengue fever (DF) associated dengue hemorrhagic fever / dengue shock syndrome has emerged as an important public health problem in the countries of the South East-Asia and Western Pacific region.

Corresponding author:

Risva

E-mail: risvaskm@gmail.com




Figure. 1 Study area with principal cities and town

Geographical Information System (GIS) and SPSS are utilized to deliver more information to prevent DHF disease. The objective of this study was to investigate the DHF vulnerability level based on the population density, rainfall, drainage condition, natural water body, vector control using GIS.

Dengue fever (DF) and dengue hemorrhagic fever (DHF) are caused by the dengue virus. Virus of dengue is a mosquito-borne infection which in recent years has become a mayor international public health concern. Several studies have been applied in term of Applying geographical information system (GIS) corresponding to the dengue hemorrhagic fever such as mosquito habitat and dengue risk potential in Hawaii; a conceptual framework and GIS application⁴. There are few studies have analyzed spatial distribution vulnerability level based on population density, rainfall, drainage condition, natural water body, and vector control. In this context, application of GIS modeling for dengue fever to the prone area based on socio-cultural and environmental factors⁵. Application of GIS in modeling of dengue risk based on sociocultural data, found that any steps taken to improve any of the social and cultural practices would have favorable effects on reducing dengue case⁶. The use of GIS in Ovitrap monitoring for dengue control Singapore, found that GIS was used to plan vector surveillance and control operations⁷. Modeling dengue fever risk based on socioeconomic parameters, nationality and age group; GIS and remote sensing based case study, found that there was a strong positive association between dengue fever cases and socio-economic factors⁸. Numerous reviews have broadly addressed the used of geographical information systems (GIS), mapping and spatial and space-time modeling approaches in operational control program⁹⁻¹¹. The weather variability such as monthly maximum,

minimum temperature, rainfall, and relative humidity identified as meaningful and significant indicators for the increasing occurrence of dengue fever¹². Incorporating GIS mapping and google earth technologies in a dengue surveillance system for developing countries might be used to prioritize control strategies and to target interventions to highest risk areas in order to eliminate the likely origin of the mosquito vector¹³. Geographical information systems is also utilized to assessing the spreading patterns of dengue infection and chikungunya fever outbreaks¹⁴.

This study aims to investigate the level of vulnerability of DHF at sub-district scale based on population density, rainfall, drainage condition, natural water body and vector control program. A geographical information system (GIS) software were used for mapping the spatial distribution of vulnerability area of DHF in sub-district of Tanjung Redeb for every three months period of 2014. To know the level of vulnerability of dengue hemorrhagic fever, this study analyzes the data using an SPSS approach and then the result would be converted to the Geographical Information System (GIS). With those facts in mind, there are few studies have been published using GIS methods to model the spatial relationship between dengue fever cases, population, population density, to create a predictable mapping of level of vulnerability based on environmental, climate, and human factors. This study hypothesizes that the spatial distribution based on climate, landscape, and human factors might be used to predict the vulnerability level of dengue haemorrhagic fever (DHF) at fine scale and could help the government to conduct a better strategy for controlling the transmission of DHF in each area.

MATERIALS AND METHOD

Study Area

This research was conducted in Tanjung Redeb, District of Berau since of the major dengue epidemic that occurred in East Kalimantan, Indonesia. Tanjung Redeb is the capital city of Berau on the island of East Kalimantan. Around 83.942 inhabitants, consist of 45.127 men and 38.856 women distributed over an area of 23,76 km², with a population density of around 2.848,36 inhabitants/ km² and a number of 235,178 household. In every year, wide range spread of dengue hemorrhagic fever (DHF) occurs in East-Kalimantan and caused high mortality. Actually, prevention of DHF

has been conducted but the diseases still exist in the area. DHF is significantly increasing during 2010-2011 in Tanjung Redeb where there were 144 cases in 2012. Distribution of the population of sub-district of Tanjung Redeb shown in Table 1.

Table 1. Distribution of population based on sub-distribution in Berau Districts

No	Sub-District	Men	Women	Total
	Sungai Bedungun	4.553	3.867	8.380
1	Tanjung Redeb	12.741	10.865	23.606
2	Bugis	8.053	6.886	14.919
3	Gunung Panjang	5.311	4.550	9.861
4	Karang Ambun	7.203	6.242	13.445
5	Gayam	7.266	6.466	13.732
Total		45.127	38.856	83.942

Data Collection and Analysis

Data have been collected consist of primary and secondary data. Primary data were collected through a filed survey. The questionnaire covered about gender, occupational pattern, awareness, and knowledge about dengue, mosquito protection practice. Secondary data were obtained from three government office including health office, city office and statistic office. The collecting data was conducted on March 2015 to September 2015. The data were collected in Tanjung Redeb such as population density, rainfall, drainage condition, natural water body and vector control used in the building of GIS layer to create the dengue potential map. Geographical Information System (GIS) is an automated computer-based system with the ability to capture, retrieve, manage, display and analyze large quantities of spatial and temporal data in a geographical context. In this study, GIS is utilized to demonstrate the spatial distribution of dengue hemorrhagic fever (DHF) vulnerability level based on population density, rainfall, drainage condition, natural water body and vector control program. This research used analysis epidemiology with ArcGIS 10.

RESULTS AND DISCUSSION

The questionnaires were administered during the survey to the randomly selected respondents for data collection. SPSS software was employed for the

parameter estimation and GIS then used to create the dengue potential map. We divided three-part subsections to describe the main result.

Spread of DHF

DHF data in District of Berau is provided by Abdul Rival Hospital, it is an official hospital of Berau. The data were reported in weekly and monthly. The DHF was significantly increase in Tanjung Redeb during 2013 and it might become a serious problem if not addressed properly. The following is a mapping of DHF based on the data during 2013 in Tanjung Redeb. As can be seen in Fig. 2 that the red color is an area of DHF case, the number of DHF case in the district of Berau is 169 cases in 2013, and there were 111 cases in Tanjung Redeb. It means 65.68 % of DHF in District of Berau exist in Tanjung Redeb.




Fig. 2 Spread of DHF in Tanjung Redeb at 2013

The Level of Vulnerability

The level of Vulnerability was obtained by analyzing the determinant variable such as population density, landscape, rainfall and larva free index. The variables which have been given value (signed) and weighting according to the magnitude of the effect of vulnerability of time and area toward DHF and then over layer. Its result then divided within three class categories/level namely low risk, medium risk, and high risk of DHF. This prediction would be a reference point for future research since the population density, landscape, rainfall, and larva free index would not be significantly changing in short time, therefore this research is assumed relevant for preventing based on time and area, and risk each other. The following is a mapping of the vulnerability of DHF during 2014 and divided into four quarters. Fig. 3 is a mapping of vulnerability level of DHF for the first quarter.




Fig. 3 Vulnerability level of DHF for First quarter

Fig. 3 shows that there are some areas with vulnerability level medium and some areas exist in high vulnerability level as well as a few areas exist in low level. The vulnerability level of DHF for 2nd quarter is presented in Figure 4.




Fig. 5 Vulnerability level of DHF for third quarter

Figure 5 shows that the vulnerability level of DHF almost same between high, moderate, and low level. The vulnerability level of DHF for 4th quarter is depicted in figure 6.




Fig. 4 Vulnerability level of DHF for 2nd quarter

As can be seen in figure 4, there are several areas exist in high vulnerability level and there were some areas exist in medium vulnerability level. The vulnerability level of DHF for 3rd quarter is shown in figure 5.




Fig. 6 Vulnerability level of DHF for fourth quarter

In figure 6, it shows that the low vulnerability level of DHF is higher than moderate and high level.

The main objective of the study is to investigate the spatial distribution dengue haemorrhagic fever (DHF) vulnerability level based on population density, rainfall, drainage condition, natural water body, vector control program. Analysis of the survey data revealed that the level of DHF vulnerability in Tanjung Redeb was fluctuate as described in a map by GIS. Applying of geographical information system analysis has improved corresponding between socioeconomic factors and dengue haemorrhagic fever (DHF) prevalence. In line with previous study (AlphanaBohra et. Al, Korine N. Kolires, Krishna Prasad et.al), the application of GIS in modeling of dengue risk based on sociocultural data; case of Jalore, Rajasthan, India has found that the sociocultural factors such as the housing patterns, limited use of mosquito protection measures, irregular water supplies, poor management of waste disposal, storage of water on the premises due to inadequate water supplies in summer months, and prolonged storage of water for domestic and other purposes significantly affected the incidence of dengue⁴. This result to be relevant with the recent study. Mosquito habitat and dengue risk potential in Hawaii: A conceptual framework and GIS application argued that by modifying inputs through a variable weighting process, decision regarding mosquito and disease control can be adjusted based on specific temporal and local spatial variation². Application of GIS modeling for the dengue fever prone area based on socio-cultural and environmental factors—A case study of Delhi City Zone. It was found that GIS modeling was done to generate risk map of dengue incidence³. This result revealed that there is relevant between previous and recent study. Advances in mapping, geographical information systems technologies, and progress in spatial as well as space-time modeling could be harnessed to prevent and control the diseases⁹. Hence, this study showed that the spatial distribution of vulnerability level of Dengue Haemorrhagic Fever DHF based on population density, rainfall, drainage condition, natural water body, and vector control program could be predicted. Even though the examined models were supported and the results were generally in line with previous studies, there are limitations associated with the present study¹⁵⁻¹⁷.

CONCLUSION

Geographical Information system modeling was done to create map potential of vulnerability level of dengue haemorrhagic fever, with three levels such

as high, medium, and low-level risk. This study also revealed that dengue haemorrhagic fever generally occurred in areas with high density population. The data analysis and modeling show that the population density, rainfall (climate change), drainage condition (landscape), natural water body, and vector control have higher influence on dengue haemorrhagic fever incidences. A geographical information system software was used for mapping the spatial distribution of vulnerability area of DHF in sub-district of Tanjung Redeb for every three months period of 2014. Level of DHF vulnerability in Tanjung Redeb was fluctuate. At the first three months was medium, and continues to become high level at the second and third three months, then become medium level at the last three months during 2014-time period. This result indicates that the spatial distribution based on climate, landscape, and human factors might be used to predict the vulnerability area of DHF at the fine scale such as sub-district level. Therefore, this result will in turn may help the government to conduct a better strategy for controlling the transmission of DHF in each area. Given the importance of the information of vulnerability level of dengue haemorrhagic fever, development of geographical information system in prevention of public health area still need to be explored further.

Conflict of Interest: All authors declare that there is no any conflict of interest within this research and publication including the financial agency

Ethical Clearance: Ethical clearance obtained from university committee and respondent agreement.

Source Funding: Mulawarman University

REFERENCES

- [1] D. . Gubler, “Dengue and dengue hemorrhagic fever,” *Pediatr. Infect. Dis. J.*, vol. 11, no. 4, pp. 311–317, 1992.
- [2] J. L. Kyle and E. Harris, “Global spread and persistence of dengue,” *Annu. Rev. Microbiol.*, vol. 62, pp. 71–92, 2008.
- [3] “WHO | Dengue and severe dengue,” WHO, 2017.
- [4] K. N. Kolivras, “Mosquito habitat and dengue risk potential in Hawaii: A conceptual framework and GIS application,” *Prof. Geogr.*, vol. 58, no. 2, pp. 139–154, 2006.
- [5] K. Bhandari Prasad, P. Raju, and Sokhi.B.S, “Application of Gis Modeling for Dengue

- Fever Prone Area Based on Socio-Cultural and Environmental Factors – a Case Study of Delhi City Zone,” *Heal. San Fr.*, vol. xxxvii, no. 1, pp. 165–170, 1995.
- [6] A. Bohra and H. Andrianasolo, “Application of GIS in Modelling of Dengue Risk based on Socio-Cultural Data: Case of Jalor, Rajasthan, India,” *22nd Asian Conf. Remote Sens.*, vol. 25, no. November, pp. 5–9, 2001.
- [7] G. T. Ai-Leen and R. J. Song, “The use of GIS in ovitrap monitoring for dengue control in Singapore,” *Dengue Bull.*, vol. 24, no. 65, pp. 110–116, 2000.
- [8] H. M. Khormi and L. Kumar, “Modeling dengue fever risk based on socioeconomic parameters, nationality and age groups: GIS and remote sensing based case study,” *Sci. Total Environ.*, vol. 409, no. 22, pp. 4713–4719, 2011.
- [9] L. Eisen and S. Lozano-Fuentes, “Use of mapping and spatial and space-time modeling approaches in operational control of *Aedes aegypti* and dengue,” *PLoS Negl. Trop. Dis.*, vol. 3, no. 4, pp. 1–7, 2009.
- [10] M. Booman, D. N. Durrheim, K. La Grange, C. Martin, A. M. Mabuza, A. Zitha, F. M. Mbokazi, C. Fraser, and B. L. Sharp, “Using a geographical information system to plan a malaria control programme in South Africa,” *Bull World Heal. Organ*, vol. 78, no. 12, pp. 1438–1444, 2000.
- [11] A. Maran and N. Nurjazuli, “Studi Deskriptif Kejadian Demam Berdarah Dengue (DBD) Dengan Pendekatan Spasial Di Kota Kupang (Analisis Data sekunder Tahun 2010-2011),” *J. Kesehat.*, 2013.
- [12] S. C. Chen, C. M. Liao, C. P. Chio, H. H. Chou, S. H. You, and Y. H. Cheng, “Lagged temperature effect with mosquito transmission potential explains dengue variability in southern Taiwan: Insights from a statistical analysis,” *Sci. Total Environ.*, vol. 408, no. 19, pp. 4069–4075, 2010.
- [13] A. Y. Chang, M. E. Parrales, J. Jimenez, M. E. Sobieszczyk, S. M. Hammer, D. J. Copenhaver, and R. P. Kulkarni, “Combining Google Earth and GIS mapping technologies in a dengue surveillance system for developing countries,” *Int. J. Health Geogr.*, vol. 8, p. 49, 2009.
- [14] T. Ditsuwan, T. Liabsuetrakul, V. Chongsuvivatwong, S. Thammaphalo, and E. McNeil, “Assessing the Spreading Patterns of Dengue Infection and Chikungunya Fever Outbreaks in Lower Southern Thailand Using a Geographic Information System,” *Ann. Epidemiol.*, vol. 21, no. 4, pp. 253–261, 2011.
- [15] D. J. Gubler, “Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century,” *Trends Microbiol.*, vol. 10, no. 2, pp. 100–103, 2002.
- [16] Stang Abdul Rahman, Amran Rahim and Anwar Mallongi, 2017. Forecasting of dengue disease incident risks using non-stationary spatial of geostatistics model in Bone Regency Indonesia. *J. Entomol.*, 14: 49-57.
- [17] Amran, Stang, and Anwar Mallongi, *AIP Conference Proceedings* 1825, 020002 (2017); doi: 10.1063/1.4978971

Call for Papers / Article Submission

The editor invites scholarly articles that contribute to the development and understanding of all aspects of Public Health and all medical specialities. All manuscripts are double blind peer reviewed. If there is a requirement, medical statistician review statistical content. Invitation to submit paper: A general invitation is extended to authors to submit papers papers for publication in IJPHRD.

The following guidelines should be noted:

- The article must be submitted by e-mail only. Hard copy not needed. Send article as attachment in e-mail.
- The article should be accompanied by a declaration from all authors that it is an original work and has not been sent to any other journal for publication.
- As a policy matter, journal encourages articles regarding new concepts and new information.
- Article should have a Title
- Names of authors
- Your Affiliation (designations with college address)
- Abstract
- Key words
- Introduction or back ground
- Material and Methods
- Findings
- Conclusion
- Acknowledgements
- Interest of conflict
- References in Vancouver style.
- Please quote references in text by superscripting
- Word limit 2500-3000 words, MSWORD Format, single file

All articles should be sent to: **editor.ijphrd@gmail.com**

Our Contact Info:

Institute of Medico-Legal Publications

501, Manisha Building, 75-76, Nehru Place, New Delhi-110019,

Mob: 09971888542, E-mail: editor.ijphrd@gmail.com

Website: www.ijphrd.com



Indian Journal of Public Health Research & Development

CALL FOR SUBSCRIPTIONS

About the Journal

Print-ISSN: 0976-0245 **Electronic - ISSN:** 0976-5506, **Frequency:** Monthly

Indian Journal of Public Health Research & Development is a double blind peer reviewed international Journal. The frequency is half yearly. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, Public Health Laws and covers all medical specialities concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and south east Asia.

The journal has been assigned international standards (ISSN) serial number and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases.

Subscription Information

Journal Title	Pricing of Journals		
	Print Only	Print+Online	Online Only
IJPHRD			
Indian	INR 7000	INR 9000	INR 5500
Foreign	USD 450	USD 550	USD 350

Note for Subscribers

Advance payment required by cheque/demand draft in the name of " **Institute of Medico-Legal Publications** payable at New Delhi.

Cancellation not allowed except for duplicate payment.

Claim must be made within six months from issue date.

A free copy can be forwarded on request.

Send all payment to :

Institute of Medico-Legal Publications

501, Manisha Building, 75-76, Nehru Place, New Delhi-110019,

Mob: 09971888542, E-mail: editor.ijphrd@gmail.com,

Website: www.ijphrd.com

