

# Application of organic acids in *Clarias gariepinus* Burchell, 1822 aquafeed: impacts on fish relative growth rate, condition factor and cannibalism ratio

*by* Rudy Agung Nugroho

---

**Submission date:** 13-Feb-2020 01:19PM (UTC+0700)

**Submission ID:** 1256652285

**File name:** P3.pdf (486.78K)

**Word count:** 2799

**Character count:** 14698

**PAPER • OPEN ACCESS**

2

**Application of organic acids in *Clarias gariepinus* Burchell, 1822 aquafeed: impacts on fish relative growth rate, condition factor and cannibalism ratio**

1

To cite this article: L Asriqah *et al* 2019 *J. Phys.: Conf. Ser.* **1277** 012032View the [article online](#) for updates and enhancements.

**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

2

## Application of organic acids in *Clarias gariepinus* Burchell, 1822 aquafeed: impacts on fish relative growth rate, condition factor and cannibalism ratio

L Asriqah, R A Nugroho, R Aryani\*

Animal Physiology, Development, and Molecular Laboratory, Department of Biology,  
Faculty of Mathematics and Natural Sciences Mulawarman University. Samarinda,  
East Kalimantan, Indonesia

\*Corresponding Author: [retno\\_ar@yahoo.co.id](mailto:retno_ar@yahoo.co.id)

**Abstract.** The present research was performed to determine the relative growth rate (RGR), condition factor (K) and cannibalism ratio (CR) of Catfish (*Clarias gariepinus* Burchell, 1822) fed different level and types of organic acid addition in the fish. Eighty fish were randomly assigned into five group triplicates, reared, and fed different type organic acid supplemented viz: control diet with no organic acid addition (B1); mixture of formic, acetic and propionic acid at level 0.05% (B2); mixture of formic, acetic and propionic acid at level 0.1% (B3); 0.05% butyric acid (B4); 0.1% butyric acid (B5), for 8 weeks. At the end of the 8<sup>th</sup> week, RGR, K, and CR were determined. Fish fed 0.1% mix of formic, acetic and propionic acid or 0.1% butyric acid in the diet had the significantly higher ( $p < 0.05$ ) RGR than control or other groups. Meanwhile, the K and CR of fish during the trial were not affected by any various and levels of the organic acid. In conclusion, the supplementation of 0.1% mix organic acid or 0.1% butyric acid in the diet of *C. gariepinus* for 8 weeks enhanced the RGR of the catfish but no effect on the condition factor and cannibalism ratio.

### 1. Introduction

In fish culture, optimum and balanced nutrition are the very important aspect which contributes up to 40–60% of total cost production in farmed fish [1, 2]. Achieving the balance of nutrition in a commercial diet that supports optimum fish growth and health has a main goal of the farmer and much researcher. Further, many researchers have conducted some studies to develop a specific diet formulation [1]. It is already a major concern that the application of antibiotics or chemical substances can be used as a growth booster in the fish fed which may help to enhance growth, survival, and feed utilization. Nevertheless, many concerns regarding the negative effects of antibiotics and/or chemical substances to the environment have led to an avoid or even a ban of the application of chemical substances in aquaculture practice [3].

A past study revealed that the use of non-chemical compound, namely acidifiers, to enhance growth performance has been applied in many fish. The application of citric acid/formic acid enhances the metabolism minerals, such as phosphorus, magnesium, calcium and iron in *Oncorhynchus mykiss*, sea *Pagrus major* and *Labeo rohita* [4, 5]. Some researchers also stated that supplementation acidifiers in the diet of fish decrease the pH in the digestive system such as stomach and foregut, which



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

increasing pepsin activity, boosting protein catabolism and mineral absorption by intestines [4, 6]. Moreover, a short-chain organic acid is commonly absorbed by the intestinal epithelia through passive diffusion, resulting in energy for maintaining the intestinal epithelia and gut health [7].

One of the strains of African catfishes, namely *Clarias gariepinus* Burchell, 1822, is currently an important species for fish culture industry in Asian countries. In Indonesia, the production of catfish placed in rank two after tilapia, which was from 144,755 MT in 2009 to 644,221 MT in 2013 [8]. The catfish is a pseudo-lungs, has relatively long bodies and a high capacity to produce mucous as a form of protective adaptation to living in stagnant environments or drought conditions. Moreover, the catfish is grouped in omnivorous fish which consumes plant material, plankton, arthropods, molluscs, fish, reptiles, and even amphibians [9]. The catfish is also more resistant to diseases and has a high growth performance [10]. In addition, cannibalistic behaviour eventually occurs at the larval and juvenile stages of the catfish which impacts on the economic benefit [11].

To boost growth status, such as relative growth rate of this catfish, aquaculturists and researchers have supplemented various compound to the diet of catfish [12-14]. Moreover, the condition factor of the catfish which is a method to understand their physical condition the well-being of an individual fish should be known. However, the information regarding dietary addition of organic acid (formic, acetic, propionic and butyric acid) on the relative growth rate (RGR), condition factor (K), and cannibalism ratio (CR) in catfish is very limit. Thus, the purpose of the present research was to determine the RGR, K, and CR of catfish fed different types of diet, containing an organic acid.

## 2. Materials and Methods

### 2.1. Site and time

The research was done at PT Suri Tani Pemuka Unit Research and Development, Ciranjang, West Java, Indonesia from March to May 2018. All *C. gariepinus* were obtained from PT Suri Tani Pemuka, Cisarua, Tegal Waru, HIAT Purwakarta Regency, West Java 41165, Indonesia. The fishes were maintained in the oxygenated polythene bag and transported by truck to PT Suri Tani Pemuka, Research and Development Farm, Ciranjang West Java, Indonesia. The fish had been acclimated and grown under the PT Suri Tani protocols and conditions.

### 2.2. Experimental design

Five groups in five triplicates separate tanks, viz: B1, control diet without organic acid addition; B2, control diet added with 0.05% formic, acetic, and propionic acid; B3, control diet added with 0.1% formic, acetic, and propionic acid; B4, control diet added with 0.05% butyric acid; B5, control diet added with 0.1% butyric acid. The control diet was formulated from a commercial diet which was obtained from PT Suri Tani Pemuka, Purwakarta, West Java, Indonesia. The commercial diet contained 35% crude protein, 8.58% crude fat, and 2.75% fibre. All fishes were reared in a plastic tank (vol. 520 L) at a stocking density of 80 fish per tank for 8 weeks.

### 2.3. Fish culture and feeding trial

In total 1200 fishes (initial average weight 8.78 g) were placed randomly into 15 plastic tanks (80 fish/tank) with a volume of 520 L. Each tank was filled with fresh water up to 500 L. The fishes were fed with B1-B5 three times per day (01:00, 05:00 and 09:00 GMT) using satiation methods for 8 weeks.

### 2.4. Measured parameters

Biomass (g) fishes per tank were determined on the first day and the final day of the experiment. The RGR was calculated using equation as previously used by Dharmaraj and Phevendaran [15]:  $[(\text{final mean mass} - \text{initial mean mass}) / (\text{initial mean mass})] \times 100$ . Meanwhile, the K was calculated according to the equation proposed by Hun-Han [16]:  $K = (W \times 100) / L^3$ , Where, W=weight of fish (g), L=Length of fish (cm). Meanwhile, the survival rate (SR) was measured by calculating live fish using equations

that were previously used by Muchlisin et al and Nugroho et al [17, 18] as  $SR = (Nt/N0) \times 100\%$ , where  $Nt$  is total fish at the final day of trial and  $N0$  is total fish at the first day of the trial.

### 2.5. Data analysis

The data of RGR, K, and CR are expressed as means  $\pm$  standard error (SE) and subjected to analysis of variance (ANOVA), followed by Duncan post hoc test to evaluate significant differences among the groups of treatments. All significant tests were at  $P < 0.05$ . All analysis was performed using SPSS version 22 (SPSS, Inc., USA).

### 3. Results

Present findings revealed that the application of 0.1% mix formic, acetic and propionic acid or 0.1% butyric acid in the diet of fish resulted higher RGR than control. However, any various of organic acid supplementation in the diet of fish had no effects on the K and CR (Table 1).

**Table 1.** Mean and standard error of relative growth rate, Condition Factor and cannibalism of Mutiara catfish (*Clarias gariepinus* Burchell, 1822) fed various organic acid supplementation for 56 days

Parameters	Diets				
	B1	B2	B3	B4	B5
RGR	15.91 $\pm$ 0.42 <sup>a</sup>	16.49 $\pm$ 0.08 <sup>ab</sup>	17.20 $\pm$ 0.52 <sup>b</sup>	15.75 $\pm$ 0.03 <sup>a</sup>	16.98 $\pm$ 0.21 <sup>b</sup>
K	0.77 $\pm$ 0.01 <sup>a</sup>	0.77 $\pm$ 0.02 <sup>a</sup>	0.79 $\pm$ 0.01 <sup>a</sup>	0.79 $\pm$ 0.01 <sup>a</sup>	0.75 $\pm$ 0.06 <sup>a</sup>
CR	6.25 $\pm$ 1.91 <sup>a</sup>	4.58 $\pm$ 2.73 <sup>a</sup>	3.75 $\pm$ 0.72 <sup>a</sup>	2.08 $\pm$ 1.50 <sup>a</sup>	5.00 $\pm$ 0.00 <sup>a</sup>

Different alphabets (a, b) indicate significantly different means for different group of diets at  $p < 0.05$ . B1= Control diet without organic acid supplementation; B2= supplemented control diets with 0.05% mix formic, acetic and propionic acid; B3= supplemented control diets with 0.1% mix formic, acetic and propionic acid; B4= supplemented control diets with 0.05% butyric acid; B5= supplemented control diets with 0.1% butyric acid; RGR= Relative growth rate; K= Condition factor; CR= Cannibalism rate.

### 4. Discussion

Organic acid has been used as supplementation in practical fish culture [19, 20]. Past researcher Da Silva et al.[21] found that the use of butyrate acid in shrimp culture has been applied as feed attractants which enhance feed utilization. The butyric acid has also improved the feed utility, digestive tract activity and reduced pH milieu in the gastrointestinal tract of a red hybrid tilapia, *Oreochromis* sp. [3, 22]. In addition, butyric acid has a property as attractant in the diet of shrimp, enhancing growth rate [23]. In contrast to current finding, the use of fumaric acid (1.5-2 g kg<sup>-1</sup>) for *C. gariepinus* significantly culture decreased not only growth statues but also feed utilities [24]. These results can be correlated with pH balance in the digestive tract of the fish fed with high levels of fumaric acid. In addition, the supplementation of propionic acid at 10<sup>-3</sup> M may decrease feeding activity of the fish. Meanwhile, previous research also revealed that the addition of acetic acid at 10<sup>-5</sup> M had a negative effects on the fish feeding activity. As the results of the effects of organic acid on the fish is different from fish to fish, Lim et al.[25] stated that the organic acid supplementation in the diet of fish may differ from fish to fish, inconsistent, depending on the feed ingredient, culture model, and water quality parameters.



## 5. Conclusion

This research highlighted the addition of organic acid in the diet has positive results on the relative growth rate but no effects on the condition factor and cannibalism ratio. The supplementation of mixture of 0.1% formic, acetic and propionic acid; or 0.1% butyric acid in the diet of *C. gariepinus* is suggested to increase relative growth rate.

## Acknowledgement

8

The author grateful to Department of Biology, Faculty of Mathematics and natural Sciences, Mulawarman University for any kinds of support to present this paper. All authors also grateful to PT Suri Tani Pemuka Unit Research and Development, Cianjur, West Java for supporting and facilitating this research.

## References

- [1] Craig S, Helfrich LA, Kuhn D, Schwarz MH 2017 Understanding fish nutrition, feeds, and feeding.
- [2] Fadri S, Muchlisin Z, Sugito S 2016 Growth performance, survival rate and feed utilization of Nile tilapia, *Oreochromis niloticus* fed experimental diet contains jaloh leaf, *Salix tetrasperma* Roxb at different levels of EM-4 probiotic. *Jurnal Ilmiah Mahasiswa Kelautan dan Perikanan Unsyiah*. **1**(2) pp 210-21.
- [3] Luckstadt C 2008 The use of acidifiers in fish nutrition. Perspectives in Agriculture, *Veterinary Science, Nutrition and Natural Resources* **3**(044) pp 1-8.
- [4] Jun-sheng L, Jian-lin L, Ting-ting W 2006 Ontogeny of protease, amylase and lipase in the alimentary tract of hybrid juvenile tilapia (*Oreochromis niloticus* × *Oreochromis aureus*) *Fish Physiology and Biochemistry* **32**(4) pp 295-303.
- [5] Vielma J, Lall S 1997 Dietary formic acid enhances apparent digestibility of minerals in rainbow trout, *Oncorhynchus mykiss* (Walbaum) *Aquaculture Nutrition*. **3**(4) pp 265-8.
- [6] Lückstädt C 2008 Effect of organic acid containing additives in worldwide aquaculture-sustainable production the non-antibiotic way *Acidifiers Anim Nutr*. **71**.
- [7] Abu Elala NM, Ragaa NM 2015 Eubiotic effect of a dietary acidifier (potassium diformate) on the health status of cultured *Oreochromis niloticus* *Journal of advanced research* **6**(4) pp 621-9.
- [8] Fauji H, Budiardi T, Ekasari J 2018 Growth performance and robustness of African Catfish *Clarias gariepinus* (Burchell) in biofloc-based nursery production with different stocking densities. *Aquaculture Research* **49**(3) pp 1339-46.
- [9] Vitule JR, Umbria S, Aranha J 2006 Introduction of the African catfish *Clarias gariepinus* (BURCHELL, 1822) into Southern Brazil *Biological Invasions* **8**(4) p 677.
- [10] Putra I, Rusliadi R, Fauzi M, Tang U, Muchlisin Z 2017 Growth performance and feed utilization of African catfish *Clarias gariepinus* fed a commercial diet and reared in the biofloc system enhanced with probiotic. *F1000Research*. **6**(1545).
- [11] Kawamura G, Bagarinao T, Yong ASK, Sao PW, Lim LS, Senoo S 2017 Optimum low salinity to reduce cannibalism and improve survival of the larvae of freshwater African catfish *Clarias gariepinus*. *Fisheries Science* **83**(4) pp 597-605.
- [12] Chris UO, Singh N, Agarwal A 2018 Nanoparticles as feed supplement on Growth behaviour of Cultured Catfish (*Clarias gariepinus*) fingerlings. *Materials Today: Proceedings* **5**(3): pp 9076-81.
- [13] El-Husseiny OM, Hassan MI, El-Haroun ER, Suloma A 2018 Utilization of poultry by-product meal supplemented with L-lysine as fish meal replacer in the diet of African catfish *Clarias gariepinus* (Burchell, 1822) *Journal of Applied Aquaculture* **30**(1) pp 63-75.
- [14] Omosowone O, Dada A, Adeparusi E 2018 Comparison of dietary butyric acid supplementation effect on growth performance and body composition of *Clarias gariepinus* and *Oreochromis niloticus* fingerlings *Iranian Journal of Fisheries Sciences* **17**(2) pp 403-12.
- [15] Dharmaraj S, Dhevendaran K 2010 Evaluation of Streptomyces as a probiotic feed for the growth of ornamental fish *Xiphophorus helleri* *Food Technology and Biotechnology* **48**(4) pp 497-504.
- [16] Hun-Han M 1978 The reproductive biology of the dab *Limanda limanda* (L.) in the North Sea: gonadosomatic index, hepatosomatic index and condition factor. *J Fish Biol* **13** pp 369-78.
- [17] Muchlisin ZA, Arisa AA, Muhammadar AA, Fadli N, Arisa II, Siti-Azizah MN 2016 Growth performance and feed utilization of keureling (*Tor tambra*) fingerlings fed a formulated diet with different doses of vitamin E (alpha-tocopherol). *Archives of Polish Fisheries* **24**(1) pp 47-52.
- [18] Nugroho RA, Manurung H, Nur FM, Prahistika W 2017 *Terminalia catappa* L. extract improves survival,

- hematological profile and resistance to *Aeromonas hydrophila* in *Betta* sp. Archives of Polish Fisheries **25**(2) pp 103-15.
- [19] Lin Y-H, Cheng M-Y 2017 Effects of dietary organic acid supplementation on the growth, nutrient digestibility and intestinal histology of the giant grouper *Epinephelus lanceolatus* fed a diet with soybean meal. Aquaculture **469** pp 106-11.
- [20] Zhu Y, Qiu X, Ding Q, Duan M, Wang C 2014 Combined effects of dietary phytase and organic acid on growth and phosphorus utilization of juvenile yellow catfish *Pelteobagrus fulvidraco*. Aquaculture **430** pp 1-8.
- [21] da Silva BC, do Nascimento Vieira F, Mourinho JLP, Ferreira GS, Seiffert WQ 2013 Salts of organic acids selection by multiple characteristics for marine shrimp nutrition Aquaculture **384** pp 104-10.
- [22] Ng WK, Koh CB, Sudesh K, Siti-Zahrah A 2009 Effects of dietary organic acids on growth, nutrient digestibility and gut microflora of red hybrid tilapia, *Oreochromis* sp., and subsequent survival during a challenge test with *Streptococcus agalactiae* Aquaculture Research **40**(13) pp 1490-500.
- [23] Da Silva BC, Vieira FdN, Mourinho JLP, Bolivar N, Seiffert WQ 2016 Butyrate and propionate improve the growth performance of *Litopenaeus vannamei*. Aquaculture research **47**(2) pp 612-23.
- [24] Omosowone O, Dada A, Adeparusi E 2015 Effects of dietary supplementation of fumaric acid on growth performance of African catfish *Clarias gariepinus* and *Aeromonas sobria* challenge Croatian Journal of Fisheries **73**(1) pp 13-9.
- [25] Lim C, Luckstadt C, Klesius P 2010 Use of organic acids, salts in fish diets Global Aquaculture Advocate **13**(5):45-6.

# Application of organic acids in *Clarias gariepinus* Burchell, 1822 aquafeed: impacts on fish relative growth rate, condition factor and cannibalism ratio

## ORIGINALITY REPORT

12%

SIMILARITY INDEX

8%

INTERNET SOURCES

8%

PUBLICATIONS

7%

STUDENT PAPERS

## PRIMARY SOURCES

1

[publications.aston.ac.uk](http://publications.aston.ac.uk)

Internet Source

4%

2

[sintadev.ristekdikti.go.id](http://sintadev.ristekdikti.go.id)

Internet Source

1%

3

Surjya Narayan Datta, Vaneet Inder Kaur, Asha Dhawan, Geeta Jassal. "Estimation of length-weight relationship and condition factor of spotted snakehead *Channa punctata* (Bloch) under different feeding regimes", SpringerPlus, 2013

Publication

1%

4

J. R. S. Vitule, S. C. Umbria, J. M. R. Aranha. "Introduction of the African Catfish *Clarias gariepinus* (BURCHELL, 1822) into Southern Brazil", Biological Invasions, 2006

Publication

1%

5

Submitted to University of Duhok

Student Paper

1%



6	Hassan, S.M.. "Haemolytic and antimicrobial activities of saponin-rich extracts from guar meal", Food Chemistry, 20100315 Publication	1 %
7	sadpas.co.za Internet Source	1 %
8	biodiversitas.mipa.uns.ac.id Internet Source	1 %
9	Submitted to National University of Singapore Student Paper	1 %
10	S. Xie. "Effects of several organic acids on the feeding behavior of Tilapia nilotica", Journal of Applied Ichthyology, 8/2003 Publication	<1 %
11	M.S. Hassaan, M.A. Soltan, S. Jarmołowicz, H.S. Abdo. " Combined effects of dietary malic acid and on growth, gut microbiota and blood parameters of Nile tilapia ( ) ", Aquaculture Nutrition, 2018 Publication	<1 %
12	www.ftb.com.hr Internet Source	<1 %
13	Submitted to Federal University of Technology Student Paper	<1 %
14	Submitted to University of Adelaide Student Paper	<1 %

---

Exclude quotes	On
Exclude bibliography	On

Exclude matches	Off
-----------------	-----