

In silico identification of natural products from *Centella asiatica* as severe acute respiratory syndrome coronavirus 2 main protease inhibitor

by Herman Herman

Submission date: 13-Jul-2022 10:51PM (UTC-0400)

Submission ID: 1870298413

File name: JAdvPharmTechRes123261-2587578_071115.pdf (1.74M)

Word count: 3792

Character count: 20938

ORIGINAL ARTICLE

In silico identification of natural products from *Centella asiatica* as severe acute respiratory syndrome-coronavirus 2 main protease inhibitor

Putu Gita Maya,
Widyaswari Mahayasih,
Harizal Herman¹, Islamudin Ahmad²

⁵ Department of Pharmacy, Faculty of Health Sciences, Universitas Esa Unggul, West Jakarta, DKI Jakarta, ¹Laboratory of Pharmaceutical Research and Development of FARMAKA TROPIS, Faculty of Pharmacy, Universitas Mulawarman, Samarinda, ²Department of Pharmaceutical Sciences, Faculty of Pharmacy, Universitas Mulawarman, Samarinda, East Kalimantan, Indonesia

J. Adv. Pharm. Technol. Res.

ABSTRACT

Severe acute respiratory syndrome-coronavirus 2 (SARS-CoV-2) main protease (S-CoV-2 Mpro) is one of the main targets in designing antiviral against SARS-CoV-2. *Centella asiatica* contains several triterpenoids, polyacetylenes, and benzoic ester derivative with various biological activities including anti-inflammation and antiviral. Triterpenoids from *C. asiatica* could act as inhibitors of S-CoV-2 Mpro. The main objective of this study was to identify potential natural products from *C. asiatica* as S-CoV-2 Mpro inhibitor with better pharmacokinetic through in silico molecular docking method. : As much as 11 compounds from *C. asiatica* were docked with S-CoV-2 Mpro (PDB ID: 6LU7) using AutoDock v4.2.6. Pharmacokinetic parameters of these compounds were assessed using SwissADME (free access webserver). Molecular docking results of 11 natural products indicated that asiatic acid 6 and asiatic acid 10 have strong interaction with quite similar binding free energy compared to native ligand (-9.00 and -9.58 kcal/mol compared to -9.18 kcal/mol, respectively) with proper interaction to the catalytic dyad (His41 and Cys145). Pharmacokinetic analysis revealed that asiatic acid 4, asiatic acid 10, and asiatic acid 11 have poor pharmacokinetic properties. These results indicated that asiatic acid 6 could be recommended for further study as S-CoV-2 M^{pro} inhibitor.

Key words: Asiaticoside, coronavirus disease-2019, isothankunic acid, molecular docking, polyacetylenes, triterpenoids

Address for correspondence:

Dr. Islamudin Ahmad,
Department of Pharmaceutical Sciences,
Faculty of Pharmacy, Universitas Mulawarman,
Samarinda, East Kalimantan, Indonesia.
E-mail: islamudinahmad@farmasi.unmul.ac.id
⁵ ss. Putu Gita Maya Widyaswari Mahayasih,
Department of Pharmacy,
Faculty of Health Sciences,
Universitas Esa Unggul, West Jakarta,
DKI Jakarta.
Email: putu.gitamaya@esaunggul.ac.id

Submitted: 08-Dec-2020

Revised: 08-Feb-2021

Accepted: 28-Mar-2021

Published: 16-Jul-2021

Access this article online

Quick Response Code:	Website: www.japtr.org
	DOI: 10.4103/japtr.JAPTR_297_20

INTRODUCTION

Coronavirus disease-2019 outbreak is caused by severe acute respiratory syndrome coronavirus-2 (S-CoV-2), a betacoronavirus strain. Development of antiviral drugs has been performed by targeting certain proteins either in the virus including structural or nonstructural proteins in S-CoV-2 or host targets.^[1]

Main protease (M^{pro}) or 3-chymotrypsin-like protease is a nonstructural protein that has a significant role in viral replication and governing the response of the cell

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Maya PG, Mahayasih W, Harizal, Herman, Ahmad I. In silico identification of natural products from *Centella asiatica* as severe acute respiratory syndrome-coronavirus 2 main protease inhibitor. J Adv Pharm Technol Res 2021;12:261-6.

host. Together with papain-like protease (M^{pro}), M^{pro} cleaves and transforms two large polyproteins (pp1 and pp1ab) translated from the viral RNA selectively at the Leu-Gln (Ser, Ala, Gly) cleavage site.^[2] Due to its crucial role and specific cleavage site, inhibiting the activity of M^{pro} enzyme would stop viral replication with no undesired toxic effect on human.

Several potential synthetic drugs have been evaluated clinically against S-CoV-2 such as ritonavir/lopinavir^[3] and cobicistat/darunavir.^[4] However, these drugs still have not been approved by FDA for S-CoV-2 patients due to several limitations such as poor pharmacokinetic,^[5] low effectivity,^[4] and certain adverse effects.^[5] On the other side, repurposing of various classes of natural products in several herbs for S-CoV-2 M^{pro} inhibitor also showed promising results. These repurposings were mainly conducted through *in silico* approaches such as terpenoids from *Cacospongia mycofijiensis*,^[6] diterpenoids and bioflavonoids from *Torreya nucifera*,^[7] glycosylated flavonoids from various herbs, and saponins and tannins from various herbs.^[8]

Centella asiatica is a traditional herb from Southeast Asia that is widely used as medicines, cosmetics, foods, and beverages. This herb is known with different local names in different countries such as tapak kuda (Indonesia), pegaga (Indonesia and Malaysia), and gotu kola (Sri Lanka). *C. asiatica* contains various classes of active compounds (triterpenes, polyacetylenes, carotenoids, flavonoids, etc.) with different pharmacological activities including antioxidant, anti-inflammation, antidiabetic, and antimicrobial.^[9] *C. asiatica* also has antiviral activity against several viral strain such as human immunodeficiency virus (HIV), hepatitis B virus (HBV), and herpes simplex virus (HSV).^[10,11]

Among these classes of active compounds, aglycone and glycoside forms of triterpenoids were considered as the most active ingredients in this herb, especially for therapeutic purposes. Several triterpenoids also have been identified *in silico* as active S-CoV-2 inhibitors in several reports such as asiatic acid derivatives and asiaticoside derivatives.^[12] In this report, we tried to identify potential triterpenoids, polyacetylenes, and benzoic ester derivatives from *C. asiatica* as S-CoV-2 M^{pro} inhibitor through molecular docking as part of repurposing of active ingredients in *C. asiatica*.

MATERIALS AND METHODS

Materials

Materials used in this investigation were 11 known compounds from *C. asiatica* obtained from various literatures^[13,14] and receptor protein S-CoV-2 M^{pro} (PDB ID:6LU7). The *in silico* molecular docking study was conducted using Autodock v4.2.6 and AutodockTools (<http://autodock.scripps.edu/>), ChemOffice Pro v15.00 PerkinElmer, Python Molecular Viewer (PMV 1.5.6), Open Babel graphical user interface (GUI), and Discovery Studio Visualizer.

Native ligand and protein receptor preparation

Protein structure of S-CoV-2 M^{pro} complexed with N3 as native ligand at PDB 6LU7 (2.16 Å resolution) was obtained from the Protein Data Bank via website <https://rcsb.org>. Native ligand from each Protein Data Bank was separated using PMV 1.5.6. Receptor and native ligand were separated in PDBQT (.pdbqt) format using AutodockTools^[15] and Open Babel GUI programs.^[16]

Test ligands preparation

The structure of test ligands from *C. asiatica* was obtained from some literatures [as shown in Figure 1]. Preparation of ligands structure was performed using ChemDraw® Pro v15 and Chem3D® Pro v15 then process, as described by Wang *et al.*^[17]

In silico molecular docking analysis

Autodock v4.2.6 (The Scripps Research Institute) was used for *in silico* molecular docking of 11 compounds from *C. asiatica* as S-CoV-2 M^{pro} inhibitor. Lamarckian parameters were used to perform the docking simulation. The grid box size and position were also validated^[3] analyze the 11 known compounds from *C. asiatica* and visualized using Accelrys Discovery Studio Visualizer 4.0.

Pharmacokinetic properties analysis

Pharmacokinetic parameters of test and native ligands were predicted using SwissADME (<http://swissadme.ch/>).^[18]

RESULTS

In silico molecular docking study

Inhibition activity of 11 natural products isolated *C. asiatica* against S-CoV-2 M^{pro} has been conducted through *in silico* molecular docking study. This investigation aimed to identify activity of S-CoV-2 M^{pro} inhibition based on the bonding form and interaction between the ligands and active site of receptor. The structure of S-CoV-2 M^{pro} consisted of 3 domains including domain I (residues 8–101), domain II (residues 102–184), and domain III (residues 185–200). Residue histidine 41 (His41) located in domain I and cysteine 145 (Cys145) located in domain II were the active sites of S-CoV-2 M^{pro}.^[19]

Docking results of ligands-receptor were evaluated by verifying re-docking result of native ligand (N3) and S-CoV-2 M^{pro}. Re-docking verification was performed based on the RMSD value of lowest interaction energy between backbone atoms and active site of S-CoV-2 M^{pro} of 1.112 Å (with Gibbs free energy of -6.91 kcal/mol at 86 clusters) and 0.978 Å (with Gibbs free energy of -9.18 kcal/mol at 36 clusters) for 100 runs with 2 Å RMSD tolerance, as described in Figure 2.

According to the interactions shown in Table 1, the interaction pattern of native ligand (N3)-amino acids located inside the active site after the re-docking was still similar to the interaction before the re-docking in which the native




Figure 1: Structure of test and native ligands (a) centellin, (b) asiaticin, (c) cenetellicin, (d) betulinic acid, (e) madasiatic acid, (f) isothankunic acid, (g) terminolic acid, (h) asiatic acid, (i) madecassic acid, (j) asiaticoside, (k) madecassoside, (l) N3, native ligand


ligand (N3) had closely interacted with His41 in domain I dan Cys145 in domain II. Native ligand (N3) was formed pi-alkyl interaction with His41, covalent bonding with Cys145, and also interaction with other residues. Molecular docking results of 11 compounds obtained from *C. asiatica* indicated that asiaticate 6 and asiaticate 10 [Figure 3] have strong interaction with quite similar binding free energy to native ligand (-9.00 and -9.58 kcal/mol compared to -9.18 kcal/mol, respectively). Asiaticate 11 even had stronger interaction with S-CoV-2 M^{pro} active site (binding free energy of -10.98 kcal/mol); however, it did not interact with His41 residue in domain I as one of the residues of catalytic dyad.

Pharmacokinetics properties analysis

Pharmacokinetics properties of native and test ligands were analyzed using free web server SwissADME.^[18] Molecular weight of 11 compounds showed a wide range of values from 250 to 1000 g/mol. In this range, 9 compounds (asiaticate 1-9) have MW below 505 g/mol which are easy to be transported, diffused, and absorbed in the human body, while 2 compounds (asiaticate 10-11) have about 950 g/mol which are hard to be distributed inside the body. These also could be figured out from high topological polar surface area ($>140 \text{ \AA}^2$),^[20] low bioavailability score (0.17 for small molecules that fail to pass rule of five),^[21] low gastrointestinal absorption,

Table 1: Characteristics of ligands-receptor interactions

Compound	Binding free energy (kcal/mol)	Inhibition constant	Interaction
Asiatate 11	-10.98	8.95 nM	Gln189, His172, Gly170, Pro168, Leu167, Glu166, Met165, His164, His163, Cys145, Ser144, Gly143, Asn142, Leu141, Phe140, Met49, Thr45, Thr26, Thr25, Thr24
Asiatate 10	-9.58	95.01 nM	Gln189, His172, Pro168, Glu166, Met165, His164, His163, Cys145, Gly143, Asn142, Leu141, Phe140, Met49, His41, Leu27, Thr26, Thr25, Thr24
N3	-9.18	186.57 nM	Ala191, Thr190, Gln189, Arg188, Asp187, His172, Pro168, Leu167, Glu166, Met165, His164, His163, Cys145, Ser144, Gly143, Asn142, Leu141, Phe140, Tyr54, Met49, His41, Thr26
Asiatate 6	-9.00	254.38 nM	Gln192, Thr190, Gln189, Arg188, Pro168, Leu167, Glu166, Met165, His164, His163, Cys145, Ser144, Gly143, Asn142, Leu141, Met49, His41, Leu27, Thr25
Asiatate 4	-8.89	303.25 nM	Gln189, Glu166, Met165, His164, His163, Cys145, Asn142, Met49, Thr45, His41, Leu27, Thr26, Thr25, Thr24
Asiatate 5	-8.07	1.21 μ M	Gln192, Thr190, Gln189, Arg188, Pro168, Leu167, Glu166, Met165, His164, Cys145, Ser144, Gly143, Asn142, Leu141, Met49, His41, Leu27
Asiatate 7	-8.02	1.33 μ M	Gln192, Ala191, Thr190, Gln189, Arg188, Pro168, Leu167, Glu166, Met165, His164, Cys145, Ser144, Gly143, Asn142, Met49, His41, Leu27
Asiatate 3	-7.88	1.67 μ M	Gln192, Thr190, Gln189, Arg188, Pro168, Leu167, Glu166, Met165, His164, His163, Cys145, Ser144, Gly143, Asn142, Leu141, Phe140
Asiatate 1	-7.84	1.79 μ M	Gln189, His172, Glu166, Met165, His164, His163, Cys145, Ser144, Gly143, Asn142, Leu141, Phe140, Met49, His41, Leu27, Thr26, Thr25, Thr24
Asiatate 2	-7.71	2.24 μ M	Gln192, Thr190, Gln189, Arg188, Pro168, Leu167, Glu166, Met165, Cys145, Ser144, Gly143, Asn142, Leu141, Met49, His41, Leu27
Asiatate 8	-7.63	2.56 μ M	Gln189, Arg188, Asp187, Glu166, Met165, His164, Cys145, Gly143, Asn142, Met49, His41, Leu27, Thr26, Thr25, Thr24
Asiatate 9	-7.52	3.07 μ M	Arg188, Asp187, Glu166, Met165, His164, Cys145, Gly143, Asn142, Met49, Ser46, His41, Leu27, Thr26, Thr25, Thr24

**Figure 2:** Native ligand (blue) and native ligand re-docked (red) comparisons

and no blood–brain barrier (BBB) permeation values that indicated that asiataste 10 and asiataste 11 have poor pharmacokinetic properties. The remaining molecules (asiataste 1–asiataste 9) showed better pharmacokinetic parameters except for asiataste 4 with high lipophilicity ($\log P > 5$) which would cause rapid metabolic turnover, low solubility, and poor absorption.^[22]

DISCUSSION

C. asiatica is a traditional herb containing various active compounds such as terpenoids and polyacetylenes. with antiviral activity against certain viruses such as HIV, HSV, and HBV.^[10,11] In this report, it was revealed that several active compounds might have anti-S-CoV-2 activity by inhibiting S-CoV-2 M^{pro}.

In general, all of the test ligands showed high affinity toward S-CoV-2 M^{pro} with binding free energy range from -7.52 to -10.98 kcal/mol; however, only asiataste 6 ($E_{\text{bind}} = -9.00$ kcal/mol) and asiataste 10 ($E_{\text{bind}} = -9.58$ kcal/mol) showed the highest affinity with proper interaction with His41 and Cys145 as catalytic dyad residues. Although both asiataste 6 and asiataste 10 were triterpenoid derivatives, they interacted with amino acid residues in the active site of S-CoV-2 M^{pro} with different parts of molecules. Asiataste 6 (a nonglycosylated triterpenoid) entered the active site and interacted with the residue through its pentacyclic triterpenoid backbone, while asiataste 10 (a glycosylated triterpenoid) entered the active site through glycosides moiety. This phenomenon was probably due to the bigger size of the asiataste 6 and asiataste 10 compared to the size of active site. Higher affinity of asiataste 10 and asiataste 11 toward S-CoV-2 M^{pro} was also probably due to the formation

of many hydrogen bonds between amino acids residues and glycoside moiety.^[23]

From pharmacokinetic viewpoint, asiatic acid 6 showed better pharmacokinetic properties than asiatic acid 10 in which the later molecule violated 3 of 5 Lipinski rules, while the former one only violated 1 of 5 Lipinski rules. Other poor pharmacokinetic parameters also showed for asiatic acid 10 such as lower gastrointestinal absorption and smaller bioavailability score [Table 2]. This high affinity-poor pharmacokinetic properties of asiatic acid 10 also can be observed from glycyrrhizin which have the same structural features as asiatic acid 10.^[24] From *in silico* molecular docking and pharmacokinetics analysis, it can be stated that asiatic acid 6 was the most recommended compound for further *in vitro* study.

Both asiatic acid 6 and asiatic acid 10 were pentacyclic triterpenoid derivatives. This class of compound showed a high affinity

toward S-CoV-2 M^{pro} in some reports including several test ligands used in these investigations such as asiatic acid (asiatic acid 8) and asiatic acid derivatives (asiatic acid 10).^[25,26] Other triterpenoid derivatives either from semisynthetic or natural also showed strong interaction including withaferin, acetylated asiatic acid,^[25] soyasaponin I, glycyrrhizin,^[24] and quinone-methide triterpenes.^[26] From these reports, it was revealed that triterpenoid derivatives need special attention for S-CoV-2 M^{pro} inhibitor.

CONCLUSIONS

Of 11 natural products isolated from *C. asiatica*, asiatic acid 6 indicated high potency as S-CoV-2 M^{pro} inhibitor. Asiatic acid 6 has quite similar binding free energy with S-CoV-2 M^{pro} as compared to native ligand (N3) with proper interaction to the catalytic dyad residues. Pharmacokinetic parameters' analysis also revealed that asiatic acid 6 has good pharmacokinetic performance. From these results, asiatic acid




Figure 3: Three dimension of S-CoV-2 M^{pro} with (a) native ligand and (b) asiatic acid 10

Table 2: Pharmacokinetics properties of native ligand (N3) and test ligands

Ligands	E _{Bind}	MW	B _{rot}	H _{Don}	H _{Acc}	TPSA	log P	GI _{Abs}	BBB	NLV	BS
N3	-9.18	684.82	22	5	9	197.83	2.50	Low	No	2	0.17
Asiatic acid 1	-7.84	250.33	8	1	3	46.53	3.17	High	Yes	0	0.55
Asiatic acid 2	-7.71	312.36	9	0	4	52.60	4.13	High	Yes	0	0.55
Asiatic acid 3	-7.88	284.35	10	3	5	86.99	1.72	High	No	0	0.55
Asiatic acid 4	-8.89	456.71	2	2	3	57.53	6.14	Low	No	1	0.85
Asiatic acid 5	-8.07	488.70	1	4	5	97.99	4.27	High	No	1	0.56
Asiatic acid 6	-9.00	504.70	2	5	6	118.22	3.65	High	No	1	0.65
Asiatic acid 7	-8.02	504.70	2	5	6	118.22	3.87	High	No	1	0.56
Asiatic acid 8	-7.63	488.70	2	4	5	97.99	4.28	High	No	0	0.56
Asiatic acid 9	-7.52	504.70	2	5	6	118.22	3.62	High	No	1	0.56
Asiatic acid 10	-9.58	959.13	10	12	19	315.21	-0.30	Low	No	3	0.17
Asiatic acid 11	-10.98	975.13	10	13	20	335.44	-1.26	Low	No	3	0.17

E_{Bind}: Binding free energy (kcal/mol), MW: Molecular weight (g/mol), B_{rot}: Number of rotatable bonds, H_{Don}: Number of hydrogen bond donors, H_{Acc}: Number of hydrogen bond acceptor, TPSA: Topological polar surface area (Å²), log P: Predicted octanol/water partition coefficient, GI_{Abs}: Gastrointestinal absorption, BBB: Blood-brain barrier permeation, NLV: Number of Lipinski rule violation, BS: Bioavailability score

6 could be recommended for further evaluation as S-CoV-2 M^{pro} inhibitor.

Financial support and sponsorship

This research was supported by Ministry of Research and Technology/National Research and Innovation Agency in funding aspect through Inter-Higher Education Collaborative Research grant with contract number 183/SP2H/AMD/LT/DRPM/2020.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Gil C, Ginex T, Maestro I, Nozal V, Barrado-Gil L, Cuesta-Geijo MÁ, et al. COVID-19: Drug targets and potential treatments. *J Med Chem* 2020;63:12359-86.
2. Hilgenfeld R. From SARS to MERS: Crystallographic studies on coronavirus proteases enable antiviral drug design. *FEBS J* 2014;281:4085-96.
3. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A trial of lopinavir-ritonavir in adults hospitalized with severe Covid-19. *N Engl J Med* 2020;382:1787-99.
4. Chen J, Xia L, Liu L, Xu Q, Ling Y, Huang D, et al. Antiviral activity and safety of darunavir/cobicistat for the treatment of COVID-19. *Open Forum Infect Dis* 2020;7:ofaa241.
5. Osborne V, Davies M, Lane S, Evans A, Denyer J, Dhanda S, et al. Lopinavir-ritonavir in the treatment of COVID-19: A dynamic systematic benefit-risk assessment. *Drug Saf* 2020;43:809-21.
6. Sepay N, Sekar A, Halder UC, Alarifi A, Afzal M. Anti-COVID-19 terpenoid from marine sources: A docking, admet and molecular dynamics study. *J Mol Struct* 2020;1228:129433.
7. Ghosh R, Chakraborty A, Biswas A, Chowdhuri S. Computer aided identification of potential SARS CoV-2 main protease inhibitors from diterpenoids and biflavonoids of *Torreya nucifera* leaves. *J Biomol Struct Dyn* 2020:1-6.
8. Falade VA, Adelusi TI, Adedotun IO, Abdul-Hammed M, Lawal TA, Agboluaje SA. *In silico* investigation of saponins and tannins as potential inhibitors of SARS-CoV-2 main protease (M^{pro}). *In Silico Pharmacol*. 2021;9:9. doi: 10.1007/s40203-020-00071-w.
9. Sardrood SG, Saadatmand S, Assareh MH, Satan TN. Chemical composition and biological activity of essential oils of *Centella asiatica* (L.). *Toxicol Environ Health Sci* 2019;11:125-31.
10. Thanigaivel S, Durgadevi H, Balasubramaniam J, Mythily V, Elanchezhian M. Comparative evaluation of the anti-Hepatitis B virus activity of *Centella asiatica* and *Camellia sinensis* (green tea). *BMC Infect Dis* 2014;14:P21.
11. Lamorde M, Tabuti JR, Obua C, Kukunda-Byobona C, Lanyero H, Byakika-Kibwika P, et al. Medicinal plants used by traditional medicine practitioners for the treatment of HIV/AIDS and related conditions in Uganda. *J Ethnopharmacol* 2010;130:43-53.
12. Sun B, Wu L, Wu Y, Zhang C, Qin L, Hayashi M, et al. Therapeutic potential of *Centella asiatica* and its triterpenes: A review. *Front Pharmacol* 2020;11:1-24.
13. Azerad R. Chemical structures, production and enzymatic transformations of saponinins and saponins from *Centella asiatica* (L.) Urban. *Fitoterapia* 2016;114:168-87.
14. Siddiqui BS, Aslam H, Ali ST, Khan S, Begum S. Chemical constituents of *Centella asiatica*. *J Asian Nat Prod Res* 2007;9:407-14.
15. Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, et al. AutoDock4 and AutodockTools4: Automated docking with selectivity receptor flexibility. *J Comput Chem* 2009;30:2785-91.
16. Boyle NM, Banck M, James CA, Morley C, Vandermeersch T, Hutchison GR. Openbabel : An open chemical toolbox. *J Cheminform* 2011;3:1-4.
17. Wang L, Wu Y, Deng Y, Kim B, Pierce L, Krilov G, et al. Accurate and reliable prediction of relative ligand binding potency in prospective drug discovery by way of a modern free-energy calculation protocol and force field. *J Am Chem Soc* 2015;137:2695-703.
18. Daina A, Michielin O, Zoete V. SwissADME: A free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci Rep* 2017;7:1-3.
19. Sk MF, Roy R, Jonniya NA, Poddar S, Kar P. Elucidating biophysical basis of binding of inhibitors to SARS-CoV-2 main protease by using molecular dynamics simulations and free energy calculations. *J Biomol Struct Dyn*. 2020:1-13.
20. Matsson P, Kihlberg J. How big is too big for cell permeability? *J Med Chem* 2017;60:1662-4.
21. Martin YC. A bioavailability score. *J Med Chem* 2005;48:3164-70.
22. Arnott JA, Planey SL. The influence of lipophilicity in drug discovery and design. *Expert Opin Drug Discov* 2012;7:863-75.
23. Chen D, Oezguen N, Urvil P, Ferguson C, Dann SM, Savidge TC. Regulation of protein-ligand binding affinity by hydrogen bond pairing. *Sci Adv* 2016;2:e1501240.
24. Narkhede RR, Pise AV, Cheke RS, Shinde SD. Recognition of natural products as potential inhibitors of COVID-19 main protease (M^{pro}): *In silico* evidences. *Nat Products Bioprospect* 2020;10:297-306.
25. Musfiroh I, Azura AR, Rahayu D. Prediction of Asiatic acid derivatives affinity against SARS-CoV-2 main protease using molecular docking. *Pharm Sci Res* 2020;7:57-64.
26. Ryu YB, Park SJ, Kim YM, Lee JY, Seo WD, Chang JS, et al. SARS-CoV 3CL^{pro} inhibitory effects of quinone-methide triterpenes from *Tripterygium regelii*. *Bioorganic Med Chem Lett* 2010;20:1873-6.

In silico identification of natural products from Centella asiatica as severe acute respiratory syndromecoronavirus 2 main protease inhibitor

ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

8%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Zeynab Fakhar, Shama Khan, Aijaz Ahmad. "ABBV-744 and Onalespid as Potential Inhibitors of SARS-CoV-2 main Protease Enzyme: A Promising Therapeutics in COVID-19?", Research Square, 2020
Publication 2%
- 2** www.researchsquare.com
Internet Source 1%
- 3** phcogj.com
Internet Source 1%
- 4** Rajesh Ghosh, Ayon Chakraborty, Ashis Biswas, Snehasis Chowdhuri. " Computer aided identification of potential SARS CoV-2 main protease inhibitors from diterpenoids and biflavonoids of leaves ", Journal of Biomolecular Structure and Dynamics, 2020
Publication 1%
- 5** www.jpionline.org
Internet Source 1%

6	Submitted to Purdue University Student Paper	1 %
7	Abhik Kumar Ray, Parth Sarthi Sen Gupta, Saroj Kumar Panda, Satyaranjan Biswal, Uddipan Bhattacharya, Malay Kumar Rana. "Repurposing of FDA-approved drugs as potential inhibitors of the SARS-CoV-2 main protease: Molecular insights into improved therapeutic discovery", Computers in Biology and Medicine, 2021 Publication	1 %
8	jonuns.com Internet Source	1 %
9	Submitted to Napier University Student Paper	1 %
10	Dibyendu Mondal, Arie Warshel. "Exploring the Mechanism of Covalent Inhibition: Simulating the Binding Free Energy of α -Ketoamide Inhibitors of the Main Protease of SARS-CoV-2", Biochemistry, 2020 Publication	1 %
11	www.pubfacts.com Internet Source	<1 %

Exclude bibliography On