

COPY RIGHT TRANSFER, AGREEMENT AND DECLARATION FORM

(Take printout of this form. Fill the yellow highlighted space . Scan and send along with your manuscript)

Without this completed form, your manuscript will not be considered for publication

To.

The MANAGING EDITOR,
INTERNATIONAL JOURNAL OF PHARMA AND BIO SCIENCES.

TITLE	BIOACTIVITY TEST OF COMPOUND (Hexa-Tetra Contana) FROM SPONGE (Callyspongia pseudoreticulata) AS ANTIBACTERIAL OF WITHERED DISEASE ON POTATO PLANT
TYPE	RESEARCH
SUBJECT	Pharmaceutical Science
BRANCH	Know the Hexa-tetra contana compound concentration's influence toward the inhibitor effect of Ralstonia solanacearum grows for the potato plant
www.ijpbs.net	

In reference to the above , I as a corresponding author along with other authors mentioned below table, submit the manuscript for publication in International Journal of Pharma and Bio sciences (IJPBS). Contributing authors refer to all the authors mentioned in the submitted manuscript .We — contributing authors, satisfy and declare the following criteria

- The animal/human study (if any) was taken after the prior approval of country/institutional relevant ethical committee and mentioned in the manuscript, along with co authors ensure the authenticity and relevance of this work and the content of the manuscript will no way harm the reputation of the journal (UPBS). A written consent was taken (for human study-if any) from the human subjects or patients involved in the study ensuring their acceptance in the study and publication of datas/photos/images in the journal and should be mentioned in the manuscript. We declare that UPBS Journal will not be responsible if any of the above not followed and the editors decision would be final if any litigation arises during processing for publication or after publication.
- The undersigned author declares that along with the co authors warrants the submitted work is
 original and has no unlawful or defamatory to the reputation of the journal or invites litigation in
 individual policies or any copy right or proprietary right. The authors and coauthors contributed
 significantly to the content and preparation of this manuscript and they have seen and approved
 the content, authorship and order of author representation. The Revision of manuscript during
 publication process rests on the contributors.
- The content if taken from Tolong an open access repository or in the public domain includes a citation and URL.
- The contributors agree that the content of the manuscript has not been published or communicated to any journal or in any citation database or in any book or chapter etc.
- All authors confirms commercial affiliations, stock or equity interests or patent licensing related
 to this manuscript that could have a financial conflict, needs to be declared to the editorial office
 separately bythe authors.
- We the contributing authors takes the responsibility for getting permission for the use of any
 material in the work that would be under copy right and for unpublished or uncopy righted work
 that are provided by others and if included in the manuscript.
- We agree that status of indexing and impact factor(I.F) or value of I.F are subject to change at any
 time and the journal does not held responsible for the changes. The changes may or may not be
 informed in time to the contributing authors by the UPBS office or its web page. Continuance of
 indexing depends upon the decision of Managing editor.
- We the contributing authors has no fictitious name or content was given in the manuscript
- If any problem or litigations arise in relevant to the manuscript or its content, the contributing authors (inclusive of corresponding author) are fully responsible to solve the issues in favour of International journal of pharma and bio sciences (www.iipbs.net) and should ensure that no harm to be made to the journal reputation. If the contribution was prepared jointly, the contributor agrees to inform the co-contributors of the terms of this Agreement and to obtain their signature to this Copyright Transfer Agreement or their written permission to sign on their behalf. The contribution is submitted only to this Journal and has not been published before, except for "preprints" as permitted above. The contributor also warrants that the contribution contains no

libellous or unlawful statements, doesn't infringe upon the rights (including without limitation the copyright, patent or trademark rights) or the privacy of others, or contain material or instructions that might cause harm or injury to any person or society or to the law of land.

COPYRIGHT TRANSFER

- 1. The contributor assigns to "International Journal of Pharma and Bio Sciences" during the full term of copyright and any extensions or renewals of that term, all copyright in and to the contribution, including but not limited to the right to publish, republish, transmit, sell, distribute and otherwise use the contribution and the material contained therein in electronic and print editions(if any) of the Journal and in derivative works, in all languages and in all media of expression now known or later developed, and to license or permit others to do so.
- Reproduction, posting, transmission or other distribution or use of the contribution or any material contained therein, in any medium as permitted hereunder, requires a citation to the Journal and an approximate credit to International Journal of Pharma and Bio Sciences as Publisher, suitable in form and content.

RETAINED RIGHTS:

Notwithstanding the above ,the contributors retains all proprietary rights other than copyright , such as patent right ,in any process ,procedure or article of manufacture described in the contribution, and the right to make oral presentation of material from the contribution.

OTHER RIGHTS OF CONTRIBUTOR

- 1. The right to share with colleagues print or electronic "preprints" of the unpublished contribution, in form and content as accepted by International Journal of Pharma and Bio Sciences, for publication in the journal. Such preprints may be posted as electronic files on the contributors own website for personal or professional use, or on the contributor's internal university or corporate networks/internet or secure external website at contributor's institution, but not for commercial sale or for any systematic external distribution by a third party. Priorto publication the contributor must include the following notice on the preprint. This is preprint of an article accepted for publication in International Journal of Pharma and Bio Sciences copyright year (copy right owner as specified in the journal).
- 2. The right, without change, to photocopy or to transmit online or to download, print out and distribute to a colleague a copy of the published contribution in whole or in part, for the colleague's personal or professional use, for the advancement of scholarly or scientific research or study.
- The right to republish, without charge, in print format, all or part of the material from the published contribution in a book written or edited by the Contributor.
- 4. The right to use selected figures and tables, and selected text (up to 200 words, exclusive of the abstract) from the contribution, for the Contributor's own teaching purposes, or for incorporation within another work by the Contributor i.e. made part of an edited work published (in print or electronic format) by a third party, or for presentation in electronic format or an external website of the Contributor.

I, the corresponding author along with the other contributing author, confirm that the authors listed on the manuscript are aware of their authorship status and qualify to be authors on the manuscript and acting with consent of all authors listed on the manuscript to international journal of pharma and bio sciences for publication.

IJPBS thank you for submitting your work for publication. In order to expedite the editing and publishing process and enable ijpbs to disseminate your work to the fullest extent, we need to have this Copyright Transfer Agreement signed and returned to us as soon as possible.

NAME	SIGNATURE											E	MAII	_ID											
OF THE AUTHORS	0																								
1 M. NURDIN	(VF	N	U	R	D	1	N	Τ.	K	1	M	@	G	М	Α	1	L		С	0	М	П	T	Т	
2 USMAN	\	S	Α	П	N	U	S	М	A	N	@	Y	M	Α	Tī	T	Π.	С	0	М	T		1	+	\dagger
3								T	T	T	T	十	┪						\dagger				一一		\vdash

4				
		 1 1 1 1 1		

- (if more authors, kindly add additional row to the above table)

 If you could not get signature from any of the author, you/corresponding author can provide your signature prefixing the word "for" after getting permission from that author and by this you/corresponding author takes the responsibility for any litigation if raised by that author in future.

制



Internationally indexed journal

Indexed in Chemical Abstract Services (USA), Index coppernicus, Ulrichs Directory of Peroidicals, Google scholar, PSOAR, EBSCO, Open J gate, Proquest, EMBASE, etc.



Rapid and Easy Publishing

The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quartely. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences.

Indexed in Elsevier Bibliographic Database (EMBASE)
SCImago Journal Rank 0.329
Impact factor 6.268* (SJIF)



Pharmaceutical Sciences

- Pharmaceutics
- Novel drug delivery system Biomaterial sciences
- Nanaotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics

- Polymer sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics



Biological Sciences

- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

ELSEVIER INDEXED JOURNAL

SNIP VALUE - 0.538 SJR - 0.274

CITESCORE - 0.35

SNIP - Source Normalised Impact per Paper SJR - SCImago Journal rank Source - www.journalmetrics.com (powered by scopus (ELSEVIER)



ISSN 0975-6299

CODEN 11PB12












International Journal of Pharma and Bio Sciences













The Plagiarism Checker







CAS Source Index (CASSI) Search Tool

AGF Arbeitsgemeinschaft der deutschen Familienorganisationen e.V.









International Journal of Pharma and Bio Sciences

ISSN 0975-6299

BIOACTIVITY TEST OF COMPOUND (Hexa-Tetra Contana) FROM SPONGE (Callyspongia pseudoreticulata) AS ANTIBACTERIAL OF WITHERED DISEASE ON POTATO PLANT (Ralstonia solanacearum)

M. NURDIN*1 AND USMAN2

¹Department of Chemistry, Islamic University, Makssar, Indonesia.
²Department of Education Mathematic and Natural Science, Mulawarman University, Samarinda, Indonesia

ABSTRACT

Sponge is one of the marine organisms of coral reefs that is able to produce secondary metabolite compounds with varieties of bioactivity and structural variations; one of its biological activities is as an antimicrobial. This study is aiming to test the bioactivity of *hexa-tetracontane* compounds that have been isolated from sponge of *Callyspongia pseudoreticulata*. An antibacterial activity test was performed on the *Ralstonia solanacearum* bacteria found in potato by using agar diffusion method with the use of filter paper discs. The result of bioactivity of *hexa-tetracontane* compounds with varied concentrations; 400 mg/mL, 500 mg/mL, 600 mg/mL with incubation periode of 1-7 days, showing that the diameter of the inhibition zone is at the peak of the bacterial growth under the concentration of 400 mg/ml and incubation period of 7 days and the diameter of inhibition zone is 21,33 mm with great difference compared to the other concentrations.

KEYWORDS: Bioactivity, Hexa-tetra contane, Abtrivacterial, Callyspongia Pseudoreticulata, Ralstonia solanacearum.



M. NURDIN*

Department of Chemistry, Islamic University, Makssar, Indonesia.

Received on: 23-05-2017

Revised and Accepted on: 26-09-2017

DOI: http://dx.doi.org/10.22376/ijpbs.2017.8.4.p184-187



Creative commons version 4.0

INTRODUCTION

Callyspongia pseudoreticulata is one of sponge that is easy to find in Indonesian ocean. Sponge is one of sea biota that contains many secondary metabolite compounds. The isolate of the sponge was reported has antimicrobial activity and antiparasite and some of its secondary metabolite compounds group successfully identified like terpenoid, steroid, and hydrocarbon.¹ There are not many researches about the exploration of secondary metabolite compounds of Callyspongia pseudoreticulata, especially the determination of the structure as the important step in developing the biogenetic pathway analysis as the guidance of compounds synthesis in industry. 3-6 Therefore, we need a further research in order to identify the secondary structure metabolite molecule and bioactivity characteristic that can be used in any sides, especially the prevention of the plant cultivation disorder that appears in sponge Callyspongia pseudoreticulata in order to obtain the optimum data related to the previous research. ⁷⁻⁸ That research is about the advantages and identifying the secondary metabolite compound group that exists in that sponge and also can be used for the food plant. One of the common disease that suffering the plant is bacteria Ralstonia solanacearum, mainly for the potato. Potato is an essential horticultural commodity in Indonesia that becomes an alternative food currently due to its potential as carbohydrate source with high protein. It encourages the food diversification program and also be as the industry raw materials. This plant is becoming one of the commodities that obtain the development priority because the need of potato tends to increase in line with

the population escalation, income improvement, and the developing of food cause. That condition causes the expansion of potato farming, also the increasing of highquality potato seeds demand.9 Two sorts of products that show the tendency to be more popular in the society consumption pattern are french fries and potato chips. The main obstacle to potato production in the tropical country including Indonesia is the existence of any dangerous diseases. This disease inflicts the significant damage to the decreasing of the crops and the limitation of good quality potato seeds by the farmer. Bacterial withered disease by Ralstonia solanacearum is able to give huge disadvantages potentially because it tend to reduce the quality and the quantity or even eliminate the plant. Generally, the plant that suffered the withered disease is those more than 6 weeks old. The plant become withered firstly that started to form the bud into the bottom part, so the mucous vessels will be seen The diseased potato tubers' color will change into brown and will also be smelly. One of the management methods is the good seeds choice, plant rotation, favorable water and air regulation around the plant. But the potent product that can restrain the disease has not found yet until today. Thus further research is very necessary to take, especially in order to know the Callyspongia pseudoreticulata sponge bioactivity toward the Ralsitonia solanacearum bacteria. One of the compounds that has been successfully isolated from Callyspongia pseudoreticulata sponge is *hexa*tetrcontane as shown in Figure 1. The compound is solid white crystalline-shaped with melting point of 79-80°C with high enough toxicity towards Artemia salina with LC_{50} 61,5 µg/ml.




Figure 1
Hexa-tetra contana compound structure.

The purpose of the research is to know the *Hexa-tetracontane* concentration's influence towards the inhibitory effect of *Ralstonia solanacearum* grows for the potato plant.

MATERIALS AND METHODS

Media Making

20 grams of Nutrient Agar (NA) medium powder and 1 liter of aquades was heated inside an Erlenmeyer flask until it's completely dissolved. After that, the cylindrical neck was covered with a gauze and was sterilized inside an autoclave with the temperature of 121 °C for 15 minutes.

Culture of Microorganism Test

Culture of Ralsitonia solanacearum that is used as

bacterial test is collected from the Laboratory of Microbiology of Agriculture Faculty Hasanuddin University. The bacteria culture is rejuvenated by transferring bacteria with inoculation loop from the stock culture to the NA medium and is incubated at the temperature of 37 $^{\circ}$ C for 18 – 24 hours.

Preparation of Sample Solution

The concentrations of hexa-tetracontane sample solution that is used to test the activity of the antibacterial are 400 μ g/ml, 500 μ g/ml dan 600 μ g/ml, which is made by weighing as much as 5 milligrams of hexa-tetracontane and dissolving it into 0.5 mL methanol.

Bacterial Activivity Test

The culture of Ralsitonia solanacearum was suspended

into sterile physiological saline, then each culture was dropped onto solid NA media surface in a sterile petri dish as they were diffused into the whole media, and was kept for 15 minutes. Having it done, 3 filter paper discs were dyed into sample solution of hexatetracontane for each concentrations; which are 400, 500, and 600 μ g/ml, and another two discs were dyed into methanol (negative control) and tetracycline (positive control) simultaneously and was dried at room temperature for 10 minutes. The dried filter paper discs were put on the surface of the test media and were incubated at 37°C for 24 hours. The antibacterial activity

can be seen by measuring the diameter of the growth inhibition zone (clear zone) around the filter paper disc.

RESULT AND DISCUSSION

The result of antibacterial activity and the diameter measurement of inhibition zone of *hexa-tetracontane* with varied concentrations; 400 mg/ml, 500 mg/mL dan 600 mg/ml, with incubation period of 1 – 7 days, can be seen at Table 1.

Table 1
Average of inhibition zone diameter of hexa-tetracontane towards the growth of Ralstonia solanacearum

Concentration	Incubatio	n period (p	er day) and	l average o	f inhibition	zone diam	eter (mm)
(mg/mL)	1	2	3	4	5	6	7
400	9.66	11.33	13.33	15.33	17.33	19.33	21.33
500	13.33	11.33	13.66	15.66	18.88	20.33	21.00
600	7.66	9.00	10.33	11.66	13.33	14.33	15.33

a- Diameter of inhibition zone (mm); b- 1 - 7 incubation period shown in days

According to the result of bioactivity test of hexatetracontane as seen in Table 1, showing that at the concentration of 400 mg/ml and incubation period of 7 days, the hexa-tetracontane reached the highest inhibition zone diameter towards the bacterial growth of Ralstonia solanacearum for 21,33 mm. Whereas the lowest inhibitory capacity is at 600 mg/ml and incubation period of 1 day for 7,66 mm. If the data above is shifted into diagrams, these are the following profiles of the inhibition zones towards the growth of the used bacteria as the effect of addition of several concentrations of hexa-tetracontane (Figure 2).




Figure 2.

Diagram of the diameter of inhibition zone profile towards the growth of Ralstonia solanacearum bacteria

The result of regression analysis in Figure 2 shows that the incubation time on day 1 until day 7 has a linier positive correlation with diameter of inhibition zone of the test bacteria. The longer the incubation period, the bigger the diameter of inhibition zone become and follow linier equations, y = 1,9643x + 7,5238; $R^2 = 0.9994$ for concentration 400 mg / ml, y = 1.8452x + 7.8571; $R^2 = 0.9604$ for concentration 500 mg / ml, y = 1.3095x + 6.4286; $R^2 = 0.9957$ for concentration 600 mg / ml.

Based on the criteria of the capacity of an antibacterial activity, the hexa-tetracontane constituents are included in the strong category to inhibit the growth of *Ralstonia solanacearum* bacteria with inhibition zone diameter of > 20 mm. ¹². The positive result that is shown towards the bacteria is indicating that the compound has a wide spectrum for the bacteria itself. Generally the groups of gram-positive bacteria are more sensitive to compounds that have antimicrobial activity compared to the gram-

negative bacteria. This sensitivity difference of grampositive and gram-negative bacteria is due to the fact that there is a difference between structures of each of the bacteria's cell walls. Positive-Gram bacteria have one layer cell wall structure containing peptidoglycan, a thin laments of teapixic acid and theuric acid. While Gram-negative bacteria have layer outside the cell wall containing 5 -10% of peptidoglycan, It also consists of proteins, lipopolysaccharides and lipoproteins. Gramnegative bacteria have two layers of lipid (lipid bilayer) called the lipopolysaccharide layer (LPS). In addition, the gram-negative bacteria's peptidoglycan walls are quite solid and compact and have the capability of "efflux-pump mechanism" by removing unnecessary compounds in the bacterial cellular biotransformation process, thus inhibiting the process of internalizing the compounds to affect the cellular mechanisms of bacteria. Therefore, antimicrobial substances are easier to penetrate into Gram-positive bacterial cells than Gram negative bacteria. 13

REFERENCES

- Achmad SA. Fourty Years in Organic Chemistry of Natural Materials, Tropical Plant of Indonesia, Recollection and Prospects. Bulletin of the Indonesian Society of Nat Prod Chem. 2004;4(2):50-4.
- 2. Alam GP, Astuti D, Sari S, Wahyuono, and M. T. Hamman. Structure Elucidation of Bioactive Compounds Isolated from Sponge Petrosia sp. Collected Bunaken bat Menado. *Indones, J, Chem.* 2005 May;5(2):177-81.
- 3. Amirl, and Budianto A. Know Sea Sponge (Demospongia) Generally. *Oseana*. Pusat Penelitian Osenografi-LIPI Jakarta. 1996;21(2):15-30. ISSN 0216-1877.
- 4. Barnes R, Calon P, and Olive P. The Invertebrate. Blackkywell Scientific Pub. Oxford, London, Endiburg, Boston, Melborne, 1989. Five Pub. p 53 49.
- 5. Rahmaniar. An Introduction to Oseana. *J. Toxin Marine*. 1992;62(1):1-11.
- 6. Razak AR, dan Ridhay A. (Collaboration between National Education Department and Teacher Training and Education Science Faculty of Tadulako University) Screening of Anti-Microbial Compounds of Some Kinds of Coral (Porifera) by Thin Layer Chromatography.Final report ;2004. The report of research Number 89/P2IPT/DPPM/P10/III/2004.
- 7. Schemitz FJ, Bowden BF, and Toth SI. Antitumour and Cytotoxic Compounds from Marine Organism, *Pharma and Bioac Nat Prod.* New York: Plenum Press; 1993.

CONCLUSION

According to the research result of *Callyspongia* pseudoreticulata, we can conclude that *Hexa-tetra* contana give diameter of inhibition zone to the bacteria *Ralstonia solanacearum* in potato plant. The compound with concentration 400 mg/ml show the best diameter of inhibition zone, 21,33 mm, and significantly different with other concentration.

ACKNOWLEDGMENTS


The authors wish to thank Imran for valuable discussions. The research was performed using the facilities of Chemical Laboratory of FMIPA UNHAS Makassar and LIPI Serpong.

CONFLICT OF INTEREST

Conflict of interest declared none.

- 8. Soest RWM van. The Indonesian Sponges Fauna, A Status Report Netherlands. *J. Sea Res.* 1989:23:223-30.
- Nurdin M, Usman H, Ala A, and Baharuddin. Structural Elucidation of Secondary Metabolites in Sponge (Callyspongia pseudoreticulata) with N-Hexane Extract. *Int. J. Agr. Syst.- IJAS*. 2014 Jun; 2(1):69-75.p-ISSN:2337-9782, e-ISSN: 2580-6815
- Mora Enda, Emrizal, and MulyantikaEriya. Isolation of Compound from EtilAsetate of Kepayang Leaves (*Panglumedule Reinw*) and Its Antibacterial Activity Test. *J. Farma Sains*. 2014 April;2(3).
- Suryaningrum T.D, Pramadhany W.W, danWikanta T. Screening of Antibacterial Compounds and Sponge Toxicity Origin of Bonerate Island of South Sulawesi.J. Pascapanen Bioteknologi Kelautan dan Perikanan. 2007 Jun;2(1):45-53.ISSN:1907-9133, e-ISSN:2406-9264.
- Wawengkang D.S, Sumilat D.A, Rotinsulu H. Characterization and Antibacterial Activity Test of Sponge Compound Haliclona sp. From Manado Bay. *Jurnal LPPM Bidang Sains dan Tekanologi*. 2014 Okt ;1(1):71-85. pISSN:2407-6074, ejournal.unsrat.ac.id
- Usman, Soekamto N.H., Usman H., Ahmad A. Phytochemical and Antibacterial Test of Leaf Methanol Extract Melochiaumbellata (Houtt) StapfvarDegrabrata (Paliasa) Against Pathogenic Bacteria.Jurnal Kimia Sains.2012 Feb ;6(1):30-33.ISSN:2085-7594.
- 14. Pooole K. Efflux Pumps as Antimicrobial. *Ann Med.* 2007;39(3):162-176

Reviewers of this article



Dr. Erwin, M.Si

Department of Chemistry, Mulawarman University, Jalan, Barong Tongkok 4, Samarinda , Indonesia



Prof.Dr.K.Suriaprabha

Asst. Editor, International Journal of Pharma and Bio sciences.



Prof.Dr.Prapurna Chandra Rao

Assistant Proffessor, KLE University, Belgaum, Karnataka, India



Prof.P.Muthuprasanna

Managing Editor, International
Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript

International Journal of Pharma and Bio Sciences is also cataloged in the following universities libraries

United States Universities



UC Berkeley Libraries, California, United States



University of Washington Libraries, Seattle, Washington, United States



Branford Price Millar Library, Oregon, United States



Colombus, Ohio, United States



Tucson, Arizona, United States



Houston Cole Library Alabama, United States



Miami, Florida, United States



Ralph Brown Draughon Library, Auburn, United States



Oklahoma, United States

United Kingdom Universities





New Britain, Connecticut



Bodleian Libraries,
Oxford, United Kingdom

Birmingham University Library, Birmingham, United Kingdom

Library



Lancaster, England



Bristol, England

Other Universities



James Cook University Library, Singapore



The University of Manchester

The university of Manchester Library, Singapore



NUS Libraries





THE UNIVERSITY OF BRITISH COLUMBIA

UBC Libraries, Vancouver, Canada



Glasgow, Scotland



Heidelberg University, Germany



Lund University Libraries, Sweden







Tübingen, Germany

And indexed /catalogued in many more indexing bodies and universities

For complete list of indexing kindly visit our journal webpage (www.ijpbs.net)
For any querry / submit mannuscript /feedback kindly visit www.ijpbs.net
Authors/readers/users may visit our journal web page (home page and instruction to authors)
for any further details



Submit your manuscript/feedback/querries to "contact" of www.ijpbs.net