

Dynamics of Water Quality in an Effluent-dominated Reach of the Santa Cruz River, AZ

Hamdhani*^{1,2}

¹ Department of Aquatic Resources Management, University of Mulawarman, Samarinda, Indonesia

² School of Natural Resources and the Environment, University of Arizona, Tucson, AZ, USA

*Corresponding author at: Email address: hamdhani@fpik.unmul.ac.id

Abstract

In Arizona, the Santa Cruz River system is one of several river systems that have experienced groundwater drops and loss of perennial flow due to groundwater pumping and drought. Along with Gila River and Salt River, the Santa Cruz River currently has artificial, perennial reaches that are supported by effluent discharge from wastewater treatment plants. The return of perennial reaches has brought back habitat for numerous of aquatic and riparian taxa. However, since these reaches are artificially supported by effluent they no longer have a natural flow regime or natural water quality gradients. Little research has been conducted on the water quality gradients along these perennial effluent-dominated reaches. The objective of this study is to quantify physical and chemical water quality gradients downstream from the effluent outfall. Beginning in August 2017, we have collected water quality data (e.g. DO, pH, conductivity, nutrient levels) monthly from six reaches spaced ~2 miles apart along the effluent-dominated lower Santa Cruz River, and will continue to do so through 2019. Additionally, we will expand our study to include other effluent-dominated reaches of Santa Cruz, Salt, and Gila rivers, as well as reference streams unaffected by effluent input. The findings of this study will help us understand how effluent can be used to restore perennial river habitats in arid climates.

Keywords: effluent; wastewater treatment plant; aquatic organism; Santa Cruz River.

1. Introduction

The discharge of effluent into dewatered streams incidentally helps re-create aquatic and riparian habitat that was previously lost (Boyle and Fraleigh, 2003; Luthy, et al., 2015), and can even support endangered aquatic and riparian species. For example, endangered California red-legged frogs (*Rana draytonii*) and San Francisco garter snakes (*Thamnophis sirtalis tetrataenia*) are found in effluent-dependent streams in California (Halaburka et al., 2013; Luthy et al., 2015) and endangered Gila topminnow (*Poeciliopsis occidentalis*) are found in similar streams in Arizona (Minckley, 1999; Sonoran Institute, 2017). However, the quality of wastewater treatment processes may limit the potential uses of effluent for ecosystem restoration in many cases (Canobbio et al., 2009; Matamoros and Rodríguez, 2017; Hamdhani et al., 2020).

Water quality in effluent-dependent streams can be quite different from that in natural streams (Brooks et al., 2006; Hamdhani et al., 2020). Differences noted in effluent-dependent or effluent-dominated streams include elevated temperatures (Boyle and Fraleigh, 2003; Canobbio et al., 2009) and nutrient levels, such as nitrate (Hur et al., 2007; Chen et al., 2009), ammonium/ammonia (Gafny et al., 2000; Boyle and Fraleigh, 2003), and phosphate (Birge et al., 1989; Chen et al., 2009). Additionally, low dissolved oxygen levels are frequently observed in stream reaches closest to effluent outfalls (Birge et al., 1989; Matamoros and Rodríguez, 2017).

Conversely, as effluent flows through a stream channel, natural remediation processes (e.g. aeration, nutrient uptake) can occur and lead to improved water quality conditions (Boyle and Fraleigh, 2003). Previous studies demonstrated that longitudinal distance from the effluent source is an important factor in these processes (Growth et al., 2009; Hughes et al., 2014), which suggests that the degree of remediation might differ depending on the length of effluent-dependent reaches. In natural stream systems, other factors such as seasonality and climate also play important roles in influencing water quality (e.g. Pionke et al., 1999; Morrill et al., 2005), so such factors likely are important in effluent-dependent streams as well. However, few studies have examined how water quality changes with varying longitudinal distances from effluent inputs and how those changes may differ by season or climate zone.

The purpose of this study was to investigate the dynamics of basic water quality parameters (e.g. temperature, dissolved oxygen, pH, nutrients) in six effluent-dependent reaches of differing lengths in three river systems across a climate gradient in Arizona (USA). Our goals were to investigate: (1) how water quality changes with distance from effluent outfalls, (2) how water quality and longitudinal patterns change with season and climate, and (3) whether water quality conditions in these systems are sufficient to support native aquatic species. We hypothesized that: (1) water quality would improve with increasing distance downstream from effluent outfalls due to natural remediation processes, (2) water quality conditions would deteriorate in summer, especially in reaches with hotter climates, and (3) water quality conditions would provide suitable habitat for most of aquatic biotas.

2. Methodology

2.1. Site description

Three reaches in the Santa Cruz River basin were sampled. The Santa Cruz River basin (22,000 km²) is located in southern Arizona and northern Mexico (Webb et al., 2014). The climate is characterized by hot summers (average high July temp: 38°C) and moderately cool winters (average low January temp: 5°C). Annual average precipitation is ~300 mm and rainfall is bimodal, with the months of January and August being the wettest and May and November being the driest

(data provided by NOAA ESRL Global Monitoring Division, Boulder, Colorado, USA [<http://esrl.noaa.gov/>]). The uppermost study reach in the Santa Cruz Basin (~25 km long) is supported by the Nogales International Wastewater Treatment Plant. This plant was constructed in 1943, upgraded in 2009, and discharges approximately 57 million liters of tertiary-treated effluent into the river each day (Santa Cruz County, 2019). The second Santa Cruz reach is shorter (~9 km) and is supported by the Agua Nueva Water Reclamation Facility (WRF), which was constructed in 1951, upgraded and modernized in 2014, and discharges approximately 30 million liters of tertiary-treated effluent into the river each day (Sonoran Institute, 2017). The third and longest (~31 km) reach is supported by the Tres Rios WRF, which was built in 1978, upgraded and modernized in 2014, and discharges approximately 115 million liters of tertiary-treated effluent each day (Sonoran Institute, 2017). In addition to baseflow from the treatment plants, all three study reaches in the Santa Cruz River experience seasonal flood events from precipitation runoff, which can surpass 280 m³/s. The three perennial study reaches are separated by dry reaches and are only briefly connected during large flood events.

Table 1. Sampling locations by river and reach, with information about the length of the reach, the number of sites sampled within each reach, and the number of monthly and seasonal sampling events for each reach.

| River | Reach/WRF | Length (km) | # Sites sampled | # Sample months | Sampling events by season |
|------------------|--------------|-------------|-----------------|-----------------|--|
| Santa Cruz River | Tres Rios | 31 | 8 | 12 | <ul style="list-style-type: none"> • Winter (3x) • Spring (3x) • Summer (3x) • Fall (3x) |
| | Nogales Int. | 24 | 5 | 6 | <ul style="list-style-type: none"> • Winter (2x) • Spring (1x) • Summer (2x) • Fall (1x) |
| | Agua Nueva | 9 | 3 | 4 | <ul style="list-style-type: none"> • Winter (1x) • Spring (1x) • Summer (1x) • Fall (1x) |

2.2. Data collection

The location and number of sampling sites for each reach were selected based on the availability of public access points and the length of reach, resulting in a range of 3 to 8 sampling sites per reach (Table 1). A minimum of 0.8 km separated each progressive downstream sampling site within a reach, with an average distance of 3.5 km between sites. To examine seasonal variation in water quality, we took measurements (described below) at each site at least once per season (winter = January to March, spring = April to June, summer = July to September, fall = October to December) in the year of 2018. To provide additional temporal resolution, one reach was sampled bi-monthly (i.e. 6x per year) and one reach was sampled monthly (i.e. 12x per year) (Table 1).

We measured basic water quality parameters (dissolved oxygen, temperature, pH, specific conductance, and alkalinity) and nutrients (total phosphorus, ammonia, and nitrate). Specific conductance is hereafter referred to as conductivity. Basic water quality parameters were measured using in situ water quality probes: dissolved oxygen (Apera Instruments AI480 DO850 probe), pH, temperature, and conductivity (Apera Instruments SX823-B multiprobe). For nutrient concentrations (total phosphorus, total ammonia nitrogen, and nitrate), we collected 500 mL water samples from each site, immediately transported them to the laboratory at University of Arizona, and stored at 4°C until analyzed using a YSI EcoSense Model 9500 Photometer (within 48 hours). When nutrient analysis results exceeded the test's calibration range, samples were diluted and rerun, and the appropriate dilution factor correction applied.

2.3. Data Analysis

Histogram analysis indicated that there were no outliers in the dataset and all variables were approximately normally distributed, except for ammonia, which was log-transformed prior to analysis (data not shown). Finally, to visualize how overall water quality varied across the year at all sites, we included all 8 water quality parameters in a principal component analysis (PCA). The resulting plot allowed us to identify data clustering across river basins, reaches, and seasons. PCA was run with PC-ORD version 7.08 (McCune and Mefford, 2018). Each of the eight water quality parameters was modeled with LMEs (Pinheiro & Bates 2000) using distance from effluent outfall (Distance), Season, and the interaction between distance and season (Distance*Season) as predictors with WWTP (Site) as a random factor: [water quality] ~ Distance * Season, random = Site. Models were run in the statistical program R (version 3.5.1: R Core Team 2019) with the package 'glmmTMB' (version 1.0.2.1; Brooks et al. 2017).

3. Results

Our Linear Mixed-Effect model indicated that temperature generally decreased with increasing distance from effluent outfalls. There was a significant interaction between distance and season in, where temperatures actually increased with distance from outfall in spring and summer. Across all study reaches, water temperature was usually elevated in spring and summer, and reduced in fall and winter. However, this pattern of increasing temperatures with distance from outfall in warmer months was not observed in the higher elevation Nogales reach, located 70 km upstream of the Tres Rios reach.

Our model indicated that dissolved oxygen also responded significantly to both season and distance from outfall (Table 2). Concentrations of dissolved oxygen were low during summer and spring and high during winter and fall. Furthermore, dissolved oxygen concentrations were consistently lowest at sampling sites closest to effluent outfalls, and then increased with the distance downstream in each season. The lowest dissolved oxygen concentration at the Tres Rios reach in the summer dropped as low as ~3 mg/L.

Table 2. Summary of Linear Mixed-Effect model results explaining variation in **temperature, dissolved oxygen, and ammonia** with predictors distance from outfall in km (Distance), season (Spring, Summer, Winter), and the interaction between distance and season with river (Site) as a random factor in the model. Results include Beta estimates (Estimates) with corresponding 95% confidence intervals (CI) and p values (p) as well as within site residual variance (σ^2), between site variance (τ_{00}), and intra-class correlation coefficient (ICC) for the random factor. Both marginal and conditional pseudo R^2 values are listed. Bold indicates significant p values of predictors ($\alpha \leq 0.05$).

| Predictors | Temperature | | | Dissolved Oxygen | | | Ammonia | | |
|-------------------|-------------|---------------|------------------|------------------|---------------|------------------|-----------|---------------|------------------|
| | Estimates | CI | p | Estimates | CI | p | Estimates | CI | p |
| (Intercept) | 20.33 | 17.57 – 23.09 | <0.001 | 8.66 | 7.39 – 9.92 | <0.001 | -0.65 | -0.99 – -0.32 | <0.001 |
| Distance | -0.14 | -0.23 – -0.06 | 0.001 | 0.09 | 0.04 – 0.15 | 0.001 | -0.02 | -0.04 – -0.01 | 0.003 |
| Spring | 4.97 | 3.15 – 6.80 | <0.001 | -1.89 | -3.08 – -0.64 | 0.003 | -0.16 | -0.52 – 0.19 | 0.364 |
| Summer | 7.70 | 5.89 – 9.50 | <0.001 | -3.63 | -4.83 – -2.42 | <0.001 | 0.15 | -0.20 – 0.49 | 0.413 |
| Winter | -1.34 | -3.14 – 0.45 | 0.143 | -0.50 | -1.70 – 0.70 | 0.418 | -0.19 | -0.54 – 0.15 | 0.274 |
| Distance*Spring | 0.18 | 0.06 – 0.30 | 0.003 | 0.00 | -0.08 – 0.08 | 0.945 | 0.00 | -0.02 – 0.02 | 0.937 |
| Distance*Summer | 0.15 | 0.03 – 0.27 | 0.016 | 0.03 | -0.05 – 0.11 | 0.437 | 0.01 | -0.01 – 0.03 | 0.348 |
| Distance*Winter | 0.06 | -0.05 – 0.18 | 0.262 | -0.03 | -0.11 – 0.04 | 0.377 | -0.01 | -0.03 – 0.02 | 0.593 |
| Random Effects: | | | | | | | | | |
| σ^2 | 7.37 | | | 3.29 | | | 0.28 | | |
| τ_{00} Site | 9.31 | | | 1.36 | | | 0.08 | | |
| ICC | 0.56 | | | 0.29 | | | 0.22 | | |
| N_{Site} | 6 | | | 6 | | | 6 | | |
| N | 170 | | | 170 | | | 170 | | |
| Marginal R^2 | 0.550 | | | 0.349 | | | 0.227 | | |
| Conditional R^2 | 0.801 | | | 0.540 | | | 0.397 | | |

Overall, conductivity values were stable across seasons and with increasing distance from effluent outfalls across most reaches. However, the Salt River at 91st Avenue reach had much higher initial conductivity values, which then increased by about 70% within 6 km from the effluent outfall.




Fig. 1. Ammonia (A), nitrate (B), and phosphorus (C) levels with increasing distances from the effluent outfall in the Tres Rios reach of the Santa Cruz River. Each point represents a sampling unit, with points coded by season. Solid lines represent the best fit regression corresponding to each season.

The highest ammonia concentrations observed across all sites were at the long Tres Rios reach of the Santa Cruz River, with values as high as 5.36 mg/L N during the fall season. However, these high ammonia concentrations decreased with the distance downstream and reached undetectable levels at the end of the reach, ~30km downstream from the outfall. In other seasons,

ammonia concentrations remained very low or undetectable across all sampling sites. Nitrate and phosphorus exhibited only minor changes with distance, with nitrate increasing slightly and phosphorus decreasing slightly.

4. Discussion

4.1. Distance and season effect

Effluent is generally warmer than natural baseflow temperatures, so heat exchange between the atmosphere and stream likely led to cooling trends (Boyle and Fraleigh, 2003). As temperatures cooled, dissolved oxygen levels rose. Increased oxygen levels were likely due to a combination of cooler temperatures, photosynthesis by macrophytes and algae, and atmospheric reaeration (O'Connor, 1967; Correa-González et al., 2014). However, the effect of longitudinal distance from effluent outfalls on water quality varied widely among measured factors.

Conversely, longitudinal variation in pH was not significant, the magnitude of change was small and values remained circumneutral to slightly basic. These results are in line with other studies of streams receiving effluent, which reported relatively stable pH values downstream of effluent outfalls (Chen et al., 2009; Prat et al., 2013; Matamoros and Rodríguez, 2017). Relatedly, we also found insignificant longitudinal variation in alkalinity, which indicates the buffering capacity to neutralize acids and bases of a water body. Some previous studies have reported a decrease in alkalinity with distance downstream (Birge et al. 1989; Boyle and Fraleigh, 2003), a result likely due to dilution by other water sources. However, our effluent-dependent study reaches are not fed by any tributaries that can dilute the effluent during baseflow.

In natural streams, conductivity tends to be low in headwater reaches and then increase in the downstream reaches, especially when streams pass through urban areas (Chusov et al., 2014). However, previous studies of effluent-fed streams have reported elevated conductivities near outfalls and then relatively stable values downstream (Chen et al., 2009; Prat et al., 2013; Matamoros and Rodríguez, 2017). In general, our results show stable longitudinal conductivity values, with two exceptions.

Our data indicated that across different reach lengths on the nutrient levels shows that phosphorus and nitrate variations were not significantly affected by longitudinal distance from the outfalls. Previous phosphorus studies in the other effluent-fed streams (Birge et al. 1989; Chen et al., 2009) indicated concentration decreased with distance downstream. When we look at our long reaches separately we observed similar decreasing trends. This reduction is likely due to biotic and abiotic nutrient uptake along the reaches (Davis and Minshall, 1999; Stutter et al., 2010).

The ammonia levels decreased longitudinally as nitrate levels were relatively stable. High levels of ammonia have commonly been reported near effluent outfalls, with decreasing levels downstream (Sebenik et al., 1972; Hamdhani et al., 2020). In the long reach of Tres Rios, the average rate of ammonia loss in the fall season was about ten times greater than nitrate gain (-0.11 mg/L/km of N ammonia; 0.01 mg/L/km of N nitrate). In summer, the average rate of ammonia loss was double that of nitrate gain (-0.08 mg/L/km of N ammonia; 0.04 mg/L/km of N nitrate). The increased nitrate levels are likely due to nitrification occurring in the stream channel (Sebenik et al., 1972) but do not fully account for ammonia removal, suggesting mechanisms besides oxidation of ammonia to nitrate contributed to ammonia loss during river transport. These other mechanisms include volatilization, adsorption by cation exchange complex, fixation by clays, and fixation by organic matter (Van Schreven, 1968; Weiler, 1979; Freney et al., 1981; Simon and Kennedy, 1987; Kowalchuk et al., 1998; Chen et al., 2007; Lin et al., 2019). Since pH at all sites

was always <9 (see Supplementary Material), total ammonia nitrogen was present mainly as ammonium ion (NH_4^+), thus volatilization is not expected to have contributed significantly to ammonia loss in any of the study reaches.

This effect is usually stronger in unregulated stream systems (Bowles et al., 1977) and relatively weak in effluent-dependent stream systems, especially near effluent outfalls (Hamdhani et al., 2020). However, we did observe a significant effect of season on water temperatures across all six study reaches, with warmer water observed in spring and summer than in fall and winter, regardless of reach length. This pattern demonstrates the strong influence of atmospheric and climatic conditions on water temperature despite the stable, warm temperature of effluent at its point of release. Trends in dissolved oxygen levels across seasons were generally opposite to those of temperature, with the lowest values observed in summer; however this pattern was only significant in three of the six reaches. In other reaches, dissolved oxygen levels could be more strongly driven by non-temperature factors, such as variation in biochemical oxygen demand, water volume, and photosynthetic activity (Guasch and Subater, 1995; Mulholland et al., 2005; Mandal et al., 2010).

We found no clear pattern on conductivity and pH, but we could not detect any consistent patterns. For example, conductivity was often highest in the summer, but in some cases it was lowest in the summer, and similar inconsistencies were observed with pH (data shown in supplementary material). Both factors can be influenced by precipitation and runoff, especially in urban areas where runoff may carry pollutants (Vega et al., 1998; Laudon et al., 2000). However, we avoided sampling during periods of precipitation runoff, thus all water quality measurements reflect conditions when flow was 100% effluent. Further study is needed to understand why these factors vary inconsistently by season in the six study reaches.

Similarly, we found higher variation and inconsistency in nutrient levels than in other measured parameters (Table 2) that we attribute mainly to seasonal variation in wastewater treatment plant influent and performance. The most substantial seasonal variations in ammonia, nitrate, and phosphorus occurred at the Tres Rios WRF, which discharges to our longest study reach of the Santa Cruz River. In summer and fall, ammonia levels were much higher and nitrate levels were at their lowest, relative to concentrations observed in winter and spring, especially nearer to the effluent outfall. This finding was unexpected because conversion of ammonia to nitrate during wastewater (Siripong and Rittmann, 2007) and drinking water treatment (Liu et al., 2017) is usually highest in warmer seasons. We suspect that manipulation in treatment operations at Tres Rios WRF contributed to lessened nitrification capacity in summer (Hegg et al., 1979; Liu et al., 2012). Finally, phosphorus levels discharged in effluent from the Tres Rios WRF were higher during summer and spring, and lower during winter and fall. We speculate that this may be related to the type of *Accumulibacter* involved in phosphorus removal processes during wastewater treatment. Some types of *Accumulibacter* alter their activity patterns with seasonal-induced temperature changes (Flowers et al., 2013). Clearly, more research is needed to disentangle the complex seasonal changes in nutrient levels in effluent-dependent streams.

4.2. Geography, climate and potential for effluent to support native aquatic species

Geographic and climatic settings strongly influence the structural and functional features of natural rivers and streams (Gasith and Resh, 1999; Shi et al., 2019), but it is unclear how strongly effluent-dependent systems might be influenced. For example, we expected that water temperature would be fairly consistent across all reaches, regardless of climate zone, given the warm temperature of effluent resulting from the treatment process (Brooks et al., 2006; Hamdhani et al.,

2020). However, when considering all measured factors across all reaches, sampling sites, and sampling dates together, distinct clusters were observed for high elevation sites (>1,000 m), the medium elevation sites (~700 m), and the low elevation site (<300 m). As previously mentioned, the lowest elevation reach (91st Ave, Salt River) occurs in a stream basin with natural salt deposits (Harper, 1931). The high mineral content in the water likely contributed to the distinctness of that reach. Higher nitrate levels were observed in the higher elevation reaches. Because rates of biochemical reactions, including denitrification, are temperature-dependent (Dawson and Murphy, 1972; Xu et al., 2019), lower air temperatures may reduce nitrate removal capacity of the treatment plants in the high elevation sites.

Our six study reaches demonstrated that the effluent has restored flow to rivers that had been dry for decades due to extraction of groundwater and surface water (e.g. Webb et al. 2014). However, prior to wastewater treatment plant upgrades, water quality was generally poor and aquatic biodiversity was low (Cordy et al., 2000; Walker et al., 2005; Sonoran Institute, 2017). The ability of effluent to enhance or re-create habitat for native aquatic species is still actively debated, but some conservation successes have been noted in recent years (Halaburka et al., 2013; Luthy et al., 2015). In particular, high water temperatures, low dissolved oxygen levels, and elevated concentrations of ammonia are frequently cited as the cause of low biodiversity in effluent-dependent streams (e.g. Monda et al., 1995; Hamdhani et al., 2020; Eppehimer et al., 2020).

Among 11 native Arizona fish species tested by Carveth et al. (2006), the most sensitive to high water temperatures was the Speckled dace (*Rhinichthy osculus*), which becomes disoriented at 34°C and perishes at 36°C. One of the most tolerant fishes identified was the Gila topminnow (*Poeciliopsis occidentalis*), which can withstand temperatures as high as 38-39°C for short periods of time (Carveth et al., 2006). The highest temperature we recorded across all of our sites and seasons was 34.2°C, but in our study not all instantaneous measurements were taken at the hottest times of the day for each reach, so it is possible that temperatures might increase a further 1 to 3°C (Lowney, 2000) above our measured values. Taking that into account, the maximum temperatures that organisms might experience in some of these reaches could reach 37.2 during summer, which would be higher than Speckled dace could tolerate (Carveth et al. 2006). However, since the Lethal Thermal Method study conducted by Carveth et al. (2006) was generated with faster temperature rate of change in the laboratory (0.3°C/min) than typically occur in the stream system, so fish in stream system should better escape potentially lethal conditions, unless temperature increases at a similar rate. This result suggests that studied effluent-dependent streams in the state are hypothetically thermally suitable for native fish, but sensitive species could be negatively impacted in the warmest reaches. In these cases, a gradient in fish composition could occur, where thermally sensitive species are only found further downstream from the outfalls. A similar pattern has been reported for the thermally-sensitive Greenside Darter (*Etheostoma blennioides*) in a Canadian effluent-fed stream (Brown et al., 2011). Many aquatic animals are adapted to specific temperature ranges (Carveth et al., 2006; Eliason et al., 2011) and warmer water can affect their growth, behavior and survival (Crawshaw, 1977; Schneider and Connors, 1982; Marine and Cech, 2004).

We occasionally observed low dissolved oxygen levels near effluent outfalls in summer, with concentrations as low as 2.9 mg/L, as has been observed in similar systems (Birge et al. 1989; Boyle and Fraleigh, 2003; Matamoros and Rodríguez, 2017). As well as temperature, in our study not all instantaneous measurements were taken at the lowest times of the day for each reach, so it is possible that dissolved oxygen might decrease 2 to 3°C (Wang et al., 2003). Mortality or loss of equilibrium for fishes and other aquatic organisms can occur at concentrations between 1 and 3

mg/L (US EPA, 1986), and chronic exposure can cause behavioral changes that make individuals vulnerable to predation or other risk factors (Dean and Richardson, 1999). For native Arizona fishes, dissolved oxygen levels ranging from 0.22 to 1.47 mg/L have been reported as being lethal (Lowe et al., 1967). These findings suggest that oxygen levels were probably stressful for native fishes across all of our study sites near outfall and warm seasons. A gradient in fish composition could occur, where low oxygen sensitive species are only found further downstream from the outfalls.

Nitrogen species such as ammonia has direct toxic effects on aquatic species (e.g., Richardson, 1997; Hickey and Vickers, 1994), with concentrations greater than 2 mg/L as N (pH 7.0 and temperature 20°C) causing impairment of aquatic life (Constable et al. 2003; Yeom et al., 2007; US EPA, 2013). The vast majority of our measurements in Arizona were below this threshold, but at one site on the Santa Cruz River we did find concentrations as high as 3-5 mg/L in summer and fall. In a prior study of the Santa Cruz River, before treatment plants were upgraded, the absence of sensitive mayfly taxa was likely caused by elevated ammonia concentrations (Boyle and Fraleigh, 2003). However, recent work has shown that both mayflies and fish can even be found at locations with occasional high ammonia levels (e.g. lower Santa Cruz River: Sonoran Institute, 2017; Eppheimer et al., 2020). Together, these findings suggest that ammonia is no longer a primary concern for aquatic species in effluent-dependent streams of Arizona.

5. Conclusion

The following factors: effluent outfalls, season, and climatic and geographic factors all play important roles in the water quality dynamics of effluent-dependent streams. Water quality conditions declined somewhat in some reaches during the hottest months of the summer and, for several factors, we observed natural remediation of water quality in longer reaches. Our study also shows that effluent-dependent streams supported by high quality tertiary effluent meet or exceed water quality conditions needed to support robust assemblages of native species. However, conditions may be stressful for the most sensitive taxa at sites closest to effluent outfalls, especially in summer. Natural streams across arid and semi-arid regions are continuing to dry up due to climate change and water abstraction (Seager et al., 2007; de Graaf et al., 2019), but effluent-dependent streams are becoming more common (Luthy et al., 2015; Hamdhani et al., 2020). In conclusion we suggest that these systems have the capacity to serve as refuges for native biota, and they may become the only aquatic habitat available in many urbanizing arid regions.

Acknowledgements

We thank K. Hollien for helping in the field and laboratory works, as well as E. McGee, S. Wasko, and M. Grageda for providing useful feedback.

References

- Arizona Department of Environmental Quality (n.d.) West Central Phoenix - North Plume | Site Hydrogeology. Retrieved September 5, 2020, from <http://www.azdeq.gov/west-central-phoenix-north-plume-site-hydrogeology>
- Bills, D. & Enyedy, K. (2015). The rio de flag. In Plateau Magazine, *Picture Canyon- People, Petroglyphs, and Place*, Vol 8. Flagstaff: Museum of Northern Arizona.

- Birge, W. J., Black, J. A., Short, T. M., & Westerman, A. G. (1989). A comparative ecological and toxicological investigation of a secondary wastewater treatment plant effluent and its receiving stream. *Environmental Toxicology and Chemistry*, 8(5), 437-450.
- Bowles, D. S., Fread, D. L., & Greeney, W. J. (1977). Coupled dynamic streamflow-temperature models. *Journal of the Hydraulics Division*, 103(ASCE 12929).
- Boyle, T. P., & Fraleigh Jr, H. D. (2003). Natural and anthropogenic factors affecting the structure of the benthic macroinvertebrate community in an effluent-dominated reach of the Santa Cruz River, AZ. *Ecological Indicators*, 3(2), 93-117.
- Brooks, B. W., Riley, T. M., & Taylor, R. D. (2006). Water quality of effluent-dominated ecosystems: ecotoxicological, hydrological, and management considerations. *Hydrobiologia*, 556(1), 365-379.
- Brown, C. J., Knight, B. W., McMaster, M. E., Munkittrick, K. R., Oakes, K. D., Tetreault, G. R., & Servos, M. R. (2011). The effects of tertiary treated municipal wastewater on fish communities of a small river tributary in Southern Ontario, Canada. *Environmental Pollution*, 159(7), 1923-1931.
- Canobbio, S., Mezzanotte, V., Sanfilippo, U., & Benvenuto, F. (2009). Effect of multiple stressors on water quality and macroinvertebrate assemblages in an effluent-dominated stream. *Water, Air, and Soil Pollution*, 198(1-4), 359-371.
- Carlson, M. A., Lohse, K. A., McIntosh, J. C., & McLain, J. E. (2011). Impacts of urbanization on groundwater quality and recharge in a semi-arid alluvial basin. *Journal of Hydrology*, 409(1-2), 196-211.
- Carveth, C. J., Widmer, A. M., & Bonar, S. A. (2006). Comparison of upper thermal tolerances of native and nonnative fish species in Arizona. *Transactions of the American Fisheries Society*, 135(6), 1433-1440.
- Chen, B., Nam, S. N., Westerhoff, P. K., Krasner, S. W., & Amy, G. (2009). Fate of effluent organic matter and DBP precursors in an effluent-dominated river: A case study of wastewater impact on downstream water quality. *Water Research*, 43(6), 1755-1765.
- Chen, Z. H., Chen, L. J., Wu, Z. J., Zhang, Y. L., & Juan, Y. H. (2007). Ammonia volatilization from rice field under different water conditions in lower Liaohe River Plain. *Ying yong sheng tai xue bao= The Journal of Applied Ecology*, 18(12), 2771-2776.
- Chusov, A. N., Bondarenko, E. A., & Andrianova, M. J. (2014). Study of electric conductivity of urban stream water polluted with municipal effluents. In *Applied Mechanics and Materials* (Vol. 641, pp. 1172-1175). Trans Tech Publications Ltd.
- City of Flagstaff (2020): <https://www.flagstaff.az.gov/120/Water-Reclamation-Plants>
- Constable, M., Charlton, M., Jensen, F., McDonald, K., Craig, G., & Taylor, K. W. (2003). An ecological risk assessment of ammonia in the aquatic environment. *Human and Ecological Risk Assessment*, 9(2), 527-548.

- Cordy, G. E., Gellenbeck, D. J., Gebler, J. B., Anning, D. W., Coes, A. L., Edmonds, R. J., ... & Sanger, H. W. (2000). *Water Quality in the Central Arizona Basins, 1995-98* (Vol. 1213). US Geological Survey.
- Correa-González, J. C., del Carmen Chávez-Parga, M., Cortés, J. A., & Pérez-Munguía, R. M. (2014). Photosynthesis, respiration and reaeration in a stream with complex dissolved oxygen pattern and temperature dependence. *Ecological Modelling*, 273, 220-227.
- Crawshaw, L. I. (1977). Physiological and behavioral reactions of fishes to temperature change. *Journal of the Fisheries Board of Canada*, 34(5), 730-734.
- Davis, J. C., & Minshall, G. W. (1999). Nitrogen and phosphorus uptake in two Idaho (USA) headwater wilderness streams. *Oecologia*, 119(2), 247-255.
- Dawson, R. N., & Murphy, K. L. (1972). The temperature dependency of biological denitrification. *Water Research*, 6(1), 71-83.
- de Graaf, I. E., Gleeson, T., van Beek, L. R., Sutanudjaja, E. H., & Bierkens, M. F. (2019). Environmental flow limits to global groundwater pumping. *Nature*, 574(7776), 90-94.
- Dean, T. L., & Richardson, J. (1999). Responses of seven species of native freshwater fish and a shrimp to low levels of dissolved oxygen. *New Zealand Journal of Marine and Freshwater Research*, 33(1), 99-106.
- Du, B., Haddad, S.P., Scott, W.C., Chambliss, C.K. & Brooks, B.W. (2015). Pharmaceutical bioaccumulation by periphyton and snails in an effluent-dependent stream during an extreme drought. *Chemosphere*, 119, 927-934.
- Eliason, E. J., Clark, T. D., Hague, M. J., Hanson, L. M., Gallagher, Z. S., Jeffries, K. M., ... & Farrell, A. P. (2011). Differences in thermal tolerance among sockeye salmon populations. *Science*, 332(6025), 109-112.
- Eppehimer, D. E., Hamdhani, H., Hollien, K. D., & Bogan, M. T. (2020). Evaluating the potential of treated effluent as novel habitats for aquatic invertebrates in arid regions. *Hydrobiologia*, 847(16), 3381-3396.
- Flowers, J. J., Cadkin, T. A., & McMahon, K. D. (2013). Seasonal bacterial community dynamics in a full-scale enhanced biological phosphorus removal plant. *Water Research*, 47(19), 7019-7031.
- Fovet, O., Humbert, G., Dupas, R., Gascuel-Oudou, C., Gruau, G., Jaffrézic, A., ... & Grimaldi, C. (2018). Seasonal variability of stream water quality response to storm events captured using high-frequency and multi-parameter data. *Journal of Hydrology*, 559, 282-293.
- Freney, J. R., Simpson, J. R., & Denmead, O. T. (1981). Ammonia volatilization. *Ecological Bulletins*, 291-302.

- Gafny, S., Goren, M., & Gasith, A. (2000). Habitat condition and fish assemblage structure in a coastal mediterranean stream (Yarqon, Israel) receiving domestic effluent. In *Assessing the Ecological Integrity of Running Waters* (pp. 319-330). Springer, Dordrecht.
- Gardner, K. K., & McGlynn, B. L. (2009). Seasonality in spatial variability and influence of land use/land cover and watershed characteristics on stream water nitrate concentrations in a developing watershed in the Rocky Mountain West. *Water Resources Research*, 45(8).
- Gasith, A., & Resh, V. H. (1999). Streams in Mediterranean climate regions: abiotic influences and biotic responses to predictable seasonal events. *Annual Review of Ecology and Systematics*, 30(1), 51-81.
- Gleeson, T., Wada, Y., Bierkens, M. F., & Van Beek, L. P. (2012). Water balance of global aquifers revealed by groundwater footprint. *Nature*, 488(7410), 197-200.
- Goodrich, D. C., Kepner, W. G., Levick, L. R., & Wigington Jr, P. J. (2018). Southwestern intermittent and ephemeral stream connectivity. *JAWRA Journal of the American Water Resources Association*, 54(2), 400-422.
- Growns, I., Reinfelds, I., Williams, S., & Coade, G. (2009). Longitudinal effects of a water supply reservoir (Tallowa Dam) on downstream water quality, substrate and riffle macroinvertebrate assemblages in the Shoalhaven River, Australia. *Marine and Freshwater Research*, 60(6), 594-606.
- Guasch, H., & Subater, S. (1995). Seasonal variations in photosynthesis-irradiance responses by biofilms in Mediterranean streams. *Journal of Phycology*, 31(5), 727-735.
- Halaburka, B. J., Lawrence, J. E., Bischel, H. N., Hsiao, J., Plumlee, M. H., Resh, V. H., & Luthy, R. G. (2013). Economic and ecological costs and benefits of streamflow augmentation using recycled water in a California coastal stream. *Environmental Science & Technology*, 47(19), 10735-10743.
- Hamdhani, H., Eppehimer, D. E., & Bogan, M. T. (2020). Release of treated effluent into streams: A global review of ecological impacts with a consideration of its potential use for environmental flows. *Freshwater Biology*.
- Harper, W. G. (1931). *Soil survey of the Salt River Valley Area, Arizona*. United States Dept. of Agriculture, Bureau of Chemistry and Soils.
- Hegg, B. A., Rakness, K. L., & Schultz, J. R. (1979). *Evaluation of operation and maintenance factors limiting municipal wastewater treatment plant performance* (Vol. 1). Environmental Protection Agency, Office of Research and Development, [Office of Energy, Minerals, and Industry], Municipal Environmental Research Laboratory.
- Hickey, C. W., & Vickers, M. L. (1994). Toxicity of ammonia to nine native New Zealand freshwater invertebrate species. *Archives of environmental contamination and toxicology*, 26(3), 292-298.

- Hughes, R. M., Dunham, S., Maas-Hebner, K. G., Yeakley, J. A., Schreck, C., Harte, M., ... & Schaeffer, J. (2014). A review of urban water body challenges and approaches:(1) rehabilitation and remediation. *Fisheries*, 39(1), 18-29.
- Hur, J., Schlautman, M. A., Karanfil, T., Smink, J., Song, H., Klaine, S. J., & Hayes, J. C. (2007). Influence of drought and municipal sewage effluents on the baseflow water chemistry of an upper Piedmont river. *Environmental Monitoring and Assessment*, 132(1-3), 171-187.
- Kowalchuk, G. A., Bodelier, P. L., Heilig, G. H. J., Stephen, J. R., & Laanbroek, H. J. (1998). Community analysis of ammonia-oxidising bacteria, in relation to oxygen availability in soils and root-oxygenated sediments, using PCR, DGGE and oligonucleotide probe hybridisation. *FEMS Microbiology Ecology*, 27(4), 339-350.
- Laudon, H., Westling, O., & Bishop, K. (2000). Cause of pH decline in stream water during spring melt runoff in northern Sweden. *Canadian Journal of Fisheries and Aquatic Sciences*, 57(9), 1888-1900.
- Lencioni, V., Bernabo, P., Vanin, S., Di Muro, P., & Beltramini, M. (2008). Respiration rate and oxy-regulatory capacity in cold stenothermal chironomids. *Journal of Insect Physiology*, 54(9), 1337-1342.
- Lin, J., Tang, Y., Liu, D., Zhang, S., Lan, B., He, L., ... & Qu, Y. (2019). Characteristics of organic nitrogen fractions in sediments of the water level fluctuation zone in the tributary of the Yangtze River. *Science of The Total Environment*, 653, 327-333.
- Liu, H., Zhao, F., Mao, B., & Wen, X. (2012). Enhanced nitrogen removal in a wastewater treatment process characterized by carbon source manipulation with biological adsorption and sludge hydrolysis. *Bioresource Technology*, 114, 62-68.
- Liu, H., Zhu, L., Tian, X., & Yin, Y. (2017). Seasonal variation of bacterial community in biological aerated filter for ammonia removal in drinking water treatment. *Water Research*, 123, 668-677.
- Logan, M. F. (2006). *The lessening stream: an environmental history of the Santa Cruz River*. University of Arizona Press.
- Lowe, C. H., Hinds, D. H., & Halpern, E. A. (1967). Experimental catastrophic selection and tolerances to low oxygen concentration in native Arizona freshwater fishes. *Ecology*, 48(6), 1013-1017.
- Lowney, C. L. (2000). Stream temperature variation in regulated rivers: Evidence for a spatial pattern in daily minimum and maximum magnitudes. *Water Resources Research*, 36(10), 2947-2955.
- Luthy, R. G., Sedlak, D. L., Plumlee, M. H., Austin, D., & Resh, V. H. (2015). Wastewater-effluent-dominated streams as ecosystem-management tools in a drier climate. *Frontiers in Ecology and the Environment*, 13(9), 477-485.

- Mandal, P., Upadhyay, R., & Hasan, A. (2010). Seasonal and spatial variation of Yamuna River water quality in Delhi, India. *Environmental Monitoring and Assessment*, 170(1-4), 661-670.
- Marine, K. R., & Cech Jr, J. J. (2004). Effects of High Water Temperature on Growth, Smoltification, and Predator Avoidance in Juvenile Sacramento River Chinook Salmon. *North American Journal of Fisheries Management*, 24(1), 198-210.
- Martins, R. T., Stephan, N. N. C., & Alves, R. G. (2008). Tubificidae (Annelida: Oligochaeta) as an indicator of water quality in an urban stream in southeast Brazil. *Acta Limnologica Brasiliensia*, 20(3), 221-226.
- Matamoros, V. & Rodríguez, Y. (2017). Influence of seasonality and vegetation on the attenuation of emerging contaminants in wastewater effluent-dominated streams. A preliminary study. *Chemosphere*, 186, 269-277.
- McCune, B., & Mefford, M. J. (2018). PC-ORD: Multivariate Analysis of Ecological Data: Version 7.08 for Windows. *Wild Blueberry Media, Corvallis, Oregon, U.S.A.*
- Minckley, W. L. (1999). Ecological review and management recommendations for recovery of the endangered Gila topminnow. *The Great Basin Naturalist*, 230-244.
- Monda, D. P., Galat, D. L., & Finger, S. E. (1995). Evaluating ammonia toxicity in sewage effluent to stream macroinvertebrates: I. A multi-level approach. *Archives of Environmental Contamination and Toxicology*, 28(3), 378-384.
- Morrill, J. C., Bales, R. C., & Conklin, M. H. (2005). Estimating stream temperature from air temperature: implications for future water quality. *Journal of Environmental Engineering*, 131(1), 139-146.
- Mulholland, P. J., Houser, J. N., & Maloney, K. O. (2005). Stream diurnal dissolved oxygen profiles as indicators of in-stream metabolism and disturbance effects: Fort Benning as a case study. *Ecological Indicators*, 5(3), 243-252.
- O'Connor, D. J. (1967). The temporal and spatial distribution of dissolved oxygen in streams. *Water Resources Research*, 3(1), 65-79.
- Pionke, H. B., Gburek, W. J., Schnabel, R. R., Sharpley, A. N., & Elwinger, G. F. (1999). Seasonal flow, nutrient concentrations and loading patterns in stream flow draining an agricultural hill-land watershed. *Journal of Hydrology*, 220(1-2), 62-73.
- Prat, N., Rieradevall, M., Barata, C. & Munné, A. (2013). The combined use of metrics of biological quality and biomarkers to detect the effects of reclaimed water on macroinvertebrate assemblages in the lower part of a polluted Mediterranean river (Llobregat River, NE Spain). *Ecological Indicators*, 24, 167-176.
- Pratt, B., & Chang, H. (2012). Effects of land cover, topography, and built structure on seasonal water quality at multiple spatial scales. *Journal of Hazardous Materials*, 209, 48-58.

- Richardson, J. (1997) Acute ammonia toxicity for eight New Zealand indigenous freshwater species, *New Zealand Journal of Marine and Freshwater Research*, 31:2, 185-190, DOI: [10.1080/00288330.1997.9516757](https://doi.org/10.1080/00288330.1997.9516757)
- Santa Cruz County (2019). *Nogales Wash and International Outfall Interceptor Fact Sheet*. <https://www.santacruzcountyaz.gov/DocumentCenter/View/8418/Nogales-Wash-and-International-Outfall-Interceptor-Fact-Sheet>
- Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., ... & Gosling, S. N. (2014). Multimodel assessment of water scarcity under climate change. *Proceedings of the National Academy of Sciences*, 111(9), 3245-3250.
- Schneider, M. J., & Connors, T. J. (1982). Effects of elevated water temperature on the critical swim speeds of yearling rainbow trout *Salmo gairdneri*. *Journal of Thermal Biology*, 7(4), 227-229.
- Seager, R., Ting, M., Held, I., Kushnir, Y., Lu, J., Vecchi, G., ... & Li, C. (2007). Model projections of an imminent transition to a more arid climate in southwestern North America. *Science*, 316(5828), 1181-1184.
- Sebenik, P. G., Cluff, C. B., & DeCook, K. J. (1972). Nitrogen species transformations of sewage effluent releases in a desert stream channel. *Hydrology and Water Resources in Arizona and the Southwest*, Vol. 02 (1972)
- Shi, B., Bach, P. M., Lintern, A., Zhang, K., Coleman, R. A., Metzeling, L., ... & Deletic, A. (2019). Understanding spatiotemporal variability of in-stream water quality in urban environments—a case study of Melbourne, Australia. *Journal of Environmental Management*, 246, 203-213.
- Simon, N. S., & Kennedy, M. M. (1987). The distribution of nitrogen species and adsorption of ammonium in sediments from the tidal Potomac River and estuary. *Estuarine, Coastal and Shelf Science*, 25(1), 11-26.
- Sinokrot, B. A., & Stefan, H. G. (1994). Stream water-temperature sensitivity to weather and bed parameters. *Journal of Hydraulic Engineering*, 120(6), 722-736.
- Siripong, S., & Rittmann, B. E. (2007). Diversity study of nitrifying bacteria in full-scale municipal wastewater treatment plants. *Water Research*, 41(5), 1110-1120.
- Sonoran Institute. (2017). *A living river: Charting wetland conditions of the lower Santa Cruz River 2016 water year*.
- Stutter, M. I., Demars, B. O. L., & Langan, S. J. (2010). River phosphorus cycling: Separating biotic and abiotic uptake during short-term changes in sewage effluent loading. *Water Research*, 44(15), 4425-4436.
- US EPA. (1986). Ambient water quality criteria for dissolved oxygen. EPA 440/5-86-003, Washington, DC.

- US EPA. (2013). Aquatic Life Ambient Water Quality Criteria for Ammonia – Freshwater 2013. EPA 822-R-18-002, Washington, DC.
- US EPA. (2016). National Pollutant Discharge Elimination System Fact Sheet. NPDES Permit No. AZ0020524. <https://www.epa.gov/sites/production/files/2017-07/documents/az0020524-91st-ave-npdes-permit-factsheet-2016-10-04.pdf>
- Van Schreven, D. A. (1968). Ammonium fixation and availability of fixed ammonium In some Dutch loam and clay soils. *NJAS Wageningen Journal of Life Sciences*, 16(2), 91-102.
- Vega, M., Pardo, R., Barrado, E., & Debán, L. (1998). Assessment of seasonal and polluting effects on the quality of river water by exploratory data analysis. *Water Research*, 32(12), 3581-3592.
- Voss, K. A., Famiglietti, J. S., Lo, M., De Linage, C., Rodell, M., & Swenson, S. C. (2013). Groundwater depletion in the Middle East from GRACE with implications for transboundary water management in the Tigris-Euphrates-Western Iran region. *Water Resources Research*, 49(2), 904-914.
- Wada, Y., & Bierkens, M. F. (2014). Sustainability of global water use: past reconstruction and future projections. *Environmental Research Letters*, 9(10), 104003.
- Walker, D. B., Goforth, C., & Rector, S. (2005). *An exploration of nutrient and community variables in effluent dependent streams in Arizona* (pp. 05-09). Arizona Department of Environmental Quality.
- Wang, H., Hondzo, M., Xu, C., Poole, V., & Spacie, A. (2003). Dissolved oxygen dynamics of streams draining an urbanized and an agricultural catchment. *Ecological Modelling*, 160(1-2), 145-161.
- Webb, R. H., Betancourt, J. L., Johnson, R. R., Turner, R. M., & Fontana, B. L. (2014). *Requiem for the Santa Cruz: an environmental history of an Arizona River*. University of Arizona Press.
- Weiler, R. R. (1979). Rate of loss of ammonia from water to the atmosphere. *Journal of the Fisheries Board of Canada*, 36(6), 685-689.
- Xu, Z., Dai, X., & Chai, X. (2019). Effect of temperature on tertiary nitrogen removal from municipal wastewater in a PHBV/PLA-supported denitrification system. *Environmental Science and Pollution Research*, 26(26), 26893-26899.
- Yeom, D. H., Lee, S. A., Kang, G. S., Seo, J., & Lee, S. K. (2007). Stressor identification and health assessment of fish exposed to wastewater effluents in Miho Stream, South Korea. *Chemosphere*, 67(11), 2282-2292.