

Microorganisms In Ex-Coal Mining Land After The Application Of Organic Fertilizers In Saliki Village Samarinda, East Kalimantan

by Marjenah Marjenah

Submission date: 23-Sep-2021 11:47AM (UTC+0700)

Submission ID: 1655349269

File name: 10._Microorganisms_In_Ex-Coal_Mining.pdf (204.22K)

Word count: 2729

Character count: 15925

Microorganisms In Ex-Coal Mining Land After The Application Of Organic Fertilizers In Saliki Village Samarinda, East Kalimantan

Samsurianto, Wawan Kustiawan, Marjenah, Djumali Mardji

Abstract: This study was intended to determine whether organic fertilizers affect the presence of soil microorganisms after coal mining. The study location was the ex-coal mining area of PT Graha Benua Etam, in Saliki Village, Samarinda, East Kalimantan. This study used 4 kinds of organic fertilizer on the soil planted with Jabon seedlings, namely market waste vegetable fertilizer (WV), chicken manure (CM), cow dung (CD) and goat manure (GM). The results of this study indicated that before fertilizing 2 fungal genera were found in soil samples, namely *Aspergillus* and *Trichoderma*. At 3 months after fertilizing the soil with WV, 3 fungal genera were found, namely *Aspergillus*, *Penicillium*, and *Trichoderma*. At 3 months after fertilizing the soil with CM, CD, and GM found 3 genera of fungi the same with WV. At 6 months after fertilizing the soil with WV 3 genera of fungi were found that were same with 3 months, plus two other genera, namely *Mucor* and *Rhizopus*. At 6 months after fertilizing the soil with CM, CD, and GM the same 3 fungal genera as WV were found. Before fertilizing there was found 3 families of bacteria, namely *Azotobacteraceae*, *Bacillaceae*, and *Micrococcaceae*. At 3 and 6 months after fertilizing the soil with the four organic fertilizers, the same 3 families as before fertilizing were found. Nematodes were not found in soil samples before fertilizing, but 3 months after fertilizing with the four organic fertilizers there was found 3 genera, namely *Dorylaimus*, *Rhabditis* and *Rotylenchulus*. At 6 months after fertilizing, 5 genera of nematodes were found in the soil fertilized with WV and CM, namely *Aphelenchus*, *Dorylaimus*, *Rhabditis*, *Rotylenchulus*, and *Hoplolaimus*. In the soil fertilized with CD, only 3 genera of nematodes were found, namely *Dorylaimus*, *Rhabditis*, and *Rotylenchulus*. In the soil fertilized with GM, 4 genera of nematodes were found, namely *Dorylaimus*, *Rhabditis*, *Rotylenchulus*, and *Hoplolaimus*. It can be concluded that the number of genera of microorganisms increases with improved ex-mining soil conditions resulting from the application of organic fertilizers.

Index Terms: bacteria, ex-coal mining, fertilizer, fungi, microorganism, nematode

1 INTRODUCTION

COAL mining techniques commonly used in East Kalimantan are open pit mining with back-filling methods. Mining like this causes an increase in critical land, due to the loss of top soil, sub soil, land cover vegetation, heavy pressure from raindrops, greater soil erosion, direct touch of sunlight, soil compaction occurs due to heavy vehicle and soil parent material will appear on the surface of the soil [1]. The open coal mining system results in changes in landscape [2] and leaves environmental damage if it is not reclaimed [3], although landfill has been carried out with top soil, because fertile topsoil has mixed with the infertile soil at the bottom [4]. Topsoil is a micro- and macro-fauna habitat which plays an important role in supporting plant growth. These roles include helping to provide nutrients for plants, regulating nutrient cycles in the soil, synthesizing and breaking down soil organic matter, improving ground water availability and affecting plant health [5]. The loss of top soil is the main cause of declining population and soil micro-fauna activities which play an important role in the supply of nutrients [6]. The characteristics of ex-coal mining land include open soil, high temperature, low soil pH, non-vegetation and micro-fauna degradation [7], [8]. To reduce the decrease in soil quality due to mining, the addition of compost or organic fertilizer is a need on the soil. Compost can be made by utilizing organic waste from a very large number of markets in the city of Samarinda, where 815.04 tons/day of waste production is left out and has not been maximally processed. In 1 year, organic waste collected in the landfill (waste disposal site) is 382,957 m³ per year.

Organic waste that can be transported is 62% (237,210 m³), un-transported organic waste 32% (122,546 m³) and processed by the community for 6% (22,977 m³) [9]. These activities can cause environmental pollution and as a source of various infectious diseases in the community, even though organic market waste if processed into organic fertilizer or compost can improve environmental health and biological properties of land damaged by coal mining. Market organic waste still contains a lot of completely unraveled ingredients such as carbohydrates, fats and proteins [10] and water that can be used for to support plant growth [11]. Organic fertilizers have an essential function to bulge the surface of soil (top soil), increase the population of microorganisms, enhance absorption and save water which can increase soil fertility [12]. In this study, the plants used were *Anthocephalus cadamba* Miq. (jabon) seedlings, but what was reported in this article was microorganisms present in the soil before and after fertilization.

2 MATERIAL AND METHODS

This study used 4 organic fertilizers, namely market waste vegetable fertilizer (WV), chicken manure (CM), cow dung (CD) and goat manure (GM) which were given to ex-coal mining land with *A. cadamba* as experimental seedlings. The ingredients for the fertilizer are first processed into compost for 3 months. The land consisted of 3 blocks as replications, each block was divided into 5 plots, the distance between planting holes was 3x3 m. Each plot consisted of 24 seeds and in 1 block there were 120 planting holes, so the number of seedlings in the 3 research blocks were 360 plants. Planting holes were made with a size of width, length and depth of 30 x 30 x 30cm. A total of 1.5kg of organic fertilizer was included in each planting hole, then after 3 months it was fertilized again with the same weight. At 3 and 6 months after fertilizing, soil samples were taken from the planting holes to investigate the presence of microorganisms.

- Samsurianto, Wawan Kustiawan, Marjenah, Djumali Mardji
- Faculty of Forestry, Mulawarman University
- Email: sam_fmipa1@gmail.com, wawan.kusti51@gmail.com, djumalmardji@gmail.com

3. RESULTS AND DISCUSSION

TABLE 1. MICROORGANISMS ON EX-COAL MINING SOIL BEFORE FERTILIZING

Soil sample	Fungi	Bacteria
1	Aspergillus Trichoderma	Azotobacteraceae
2	Aspergillus Trichoderma	Azotobacteraceae
3	Aspergillus	Bacillaceae Micrococcaceae

TABLE 2. MICROORGANISMS ON EX-COAL MINING SOIL AT 3 AND 6 MONTHS AFTER FERTILIZING WITH ORGANIC FERTILIZER

Fertilizer	Fungi		Bacteria		Nematodes	
	3 months after fertilizing	6 months after fertilizing	3 months after fertilizing	6 months after fertilizing	3 months after fertilizing	6 months after fertilizing
WV1	Aspergillus Penicillium Trichoderma	Aspergillus Trichoderma	Azotobacteraceae	Azotobacteraceae	Dorylaimus Rotylenchullus	Aphelenchus Dorylaimus Rhabditis Rotylenchullus
WV2	Aspergillus Penicillium Trichoderma	Aspergillus Rhizopus Trichoderma	Azotobacteraceae	Azotobacteraceae	Rhabditis Rotylenchullus	Aphelenchus Hoplolaimus Rhabditis
WV3	Aspergillus Trichoderma	Aspergillus Trichoderma Penicillium Mucor	Bacillaceae Micrococcaceae	Azotobacteraceae	Rhabditis Rotylenchullus	Rhabditis Rotylenchullus
CM1	Aspergillus Penicillium Trichoderma	Aspergillus Trichoderma Penicillium	Azotobacteraceae	Azotobacteraceae	Dorylaimus Rotylenchullus	Rhabditis Rotylenchullus
CM2	Aspergillus Penicillium Trichoderma	Aspergillus Trichoderma	Azotobacteraceae	Azotobacteraceae	Rhabditis Rotylenchullus	Aphelenchus Dorylaimus Rhabditis Rotylenchullus
CM3	Aspergillus Trichoderma	Aspergillus Rhizopus Trichoderma	Bacillaceae Micrococcaceae	Micrococcaceae Bacillaceae	Rhabditis Rotylenchullus	Hoplolaimus Rhabditis Rotylenchullus
CD1	Aspergillus Penicillium Trichoderma	Aspergillus Penicillium Trichoderma	Azotobacteraceae	Azotobacteraceae	Dorylaimus Rotylenchullus	Dorylaimus Rotylenchullus
CD2	Aspergillus Penicillium Trichoderma	Aspergillus Penicillium Trichoderma	Azotobacteraceae	Azotobacteraceae	Rhabditis Rotylenchullus	Rhabditis Rotylenchullus
CD3	Aspergillus Trichoderma	Aspergillus Trichoderma	Bacillaceae Micrococcaceae	Micrococcaceae Bacillaceae	Rhabditis Rotylenchullus	Rhabditis Rotylenchullus
GM1	Aspergillus Penicillium Trichoderma	Aspergillus Penicillium	Azotobacteraceae	Azotobacteraceae	Dorylaimus Rotylenchullus	Dorylaimus Rhabditis Rotylenchullus
GM2	Aspergillus Penicillium Trichoderma	Aspergillus Trichoderma Penicillium	Azotobacteraceae	Azotobacteraceae	Rhabditis Rotylenchullus	Dorylaimus Rhabditis
GM3	Aspergillus Trichoderma	Aspergillus Penicillium Trichoderma	Bacillaceae Micrococcaceae	Bacillaceae	Rhabditis Rotylenchullus	Hoplolaimus Rhabditis Rotylenchullus

WV = waste vegetable. CM = chicken manure. CD = cow dung. GM = goat manure

3.1 Fungi

Before fertilizing, 2 fungal genera were found in soil samples, namely Aspergillus and Trichoderma (Table 1). At 3 months after fertilizing the soil with WV (WV1, WV2 and WV3) 3 fungal genera were found, namely Aspergillus, Penicillium, and Trichoderma. At 3 months after fertilizing the soil with CM (CM1, CM2 and CM3), CD (CD1, CD2 and CD3), and GM (GM1, GM2 and GM3) were found 3 genera of fungi, the same with WV1, WV2 and WV3 (Table 2). At 6 months after

fertilizing the soil with WV, 3 fungal genera were found which were same with 3 months, plus two other fungal genera, namely Mucor and Rhizopus. At 6 months after fertilizing the soil with CM (CM1, CM2 and CM3), CM (CM1, CM2 and CM3), and GM (GM1, GM2 and GM3) were found in 3 genera of fungi same with WV1, WV2 and WV3. So the number of fungi found before fertilizing were 2 genera, at 3 months after were 3 genera and 6 months after became 5 genera. This shows that organic fertilizers can enrich the genera of soil

fungi. It was a possibility that the fungi were carried away from the fertilizer, so that they developed in the soil. This indicated that the fungi were resistant to extreme habitat conditions. It is well known, that the condition of ex-coal mining land is very extreme, the land is open, so that the air and soil temperatures are high, the air and soil moisture is low, the soil pH is low. But in the presence of organic fertilizers, some species of microorganisms can still survive as observed in this study. According to Sajadah Co. [13], vegetable waste contains a lot of mineral nitrogen (N), phosphorus (P), potassium (K), and vitamin B12; cow manure has a high level of N, P, and K, so that it can supply the nutrients needed by the soil maximally. Fertile soil is favored by microorganisms and vegetation, so they can grow and develop well.

3.2 Bacteria

Before fertilizing, 3 families of bacteria were found, namely Azotobacteraceae, Bacillaceae, and Micrococcaceae (Table 1). At 3 and 6 months after fertilizing the soil with WV, CM, CD, and GM was found 3 families of bacteria that were same with their families before fertilizing (Table 2). This indicated that the three families of bacteria were resistant to the extreme habitat conditions, while other bacteria were dead. The number of families of bacteria in the ex-mining soil is possible to increase with the development of vegetation on it, especially vegetation that is resistant to extreme environmental conditions, especially species of herbs. The results of study conducted by Suryani et al. [14], bacteria found in chicken manure were *Lactobacillus achidophilus*, *Lactobacillus reuteri*, *Leuconostoc mensenteroide* and *Streptococcus thermophilus*, a small portion of which were *Actinomycetes* and *molds*. The other researcher noted that bacteria in cow dung were *Bacillus sp.*, *Vigna sinensis*, *Corynebacterium sp.*, and *Lactobacillus sp.*, fungi *Aspergillus* and *Trichoderma*, protozoa, yeast *Saccharomyces* and *Candida* [15]. These species of microorganisms are likely to be found directly in the fertilizer, so they are not affected by extreme environmental conditions such in at ex-coal mining area.

3.3 Nematode

Before fertilizing there was no nematodes were found in soil samples. At 3 months after fertilizing with WV, 3 genera were found, namely *Dorylaimus*, *Rhabditis* and *Rotylenchullus*. Likewise, the soil fertilized with CM, CD, and GM was found in the same genera as the soil fertilized with WV. At 6 months after fertilizing, 5 genera of nematodes were found on soil fertilized with WV and CM, namely *Aphelenchus*, *Dorylaimus*, *Rhabditis*, *Rotylenchullus*, and *Hoplolaimus*. In soil fertilized with CD, only 3 genera were found, namely *Dorylaimus*, *Rhabditis*, and *Rotylenchullus*. In the soil fertilized with GM found 4 genera, namely *Dorylaimus*, *Rhabditis*, *Rotylenchullus*, and *Hoplolaimus*. There was no nematode found before soil fertilizing, but after the soil is fertilized with the organic fertilizers, there were nematodes whose the number of genera differed according to the organic material used, where the soil fertilized with WV had the same number of nematode genera with CM, whereas in the soil fertilized with CD and GM there were fewer genera of nematodes. It means that soil fertilized with WV and CM was preferred by nematodes rather than soil fertilized with two other organic fertilizers. The application of organic fertilizer can increase the abundance of soil nematodes and make the soil environment tend to be healthy [16]. The results of the study by Liu et al.

[17] showed that organic fertilizers differed in effect on soil nematodes, fertilizers that have carbon C-rich plant residues support a population of free-living nematodes, whereas manure containing lots of nitrogen is more effective in controlling plant-eating nematodes. Application of fertilizers from plant waste is most effective way to increase soil biodiversity in intensively managed agroecosystems.

4. CONCLUSION

The use of organic fertilizers on ex-coal mining soil can increase in the number of species of fungi, bacteria and nematodes. The number of species of microorganisms is dependent on the type of organic fertilizer used. Soil fertilized with organic fertilizers derived from market waste (vegetable scraps) and chicken manure is more microorganisms found than in soil fertilized with cow and goat manure, although there are only small differences in the number of species.

REFERENCES

- [1] S. Harjowigeno, Peatland Development Opportunities Challenges For Agriculture. A Scientific Professor Oration of Soil Science, Faculty of Agriculture, Bogor Agricultural Institute, 22 June 1996.
- [2] N. Cavender, S. Byrd, C.L. Bechtoldt and J.M. Buman, "Vegetation Communities of A Coal Reclamation Site in Southeastern Ohio", *Northeastern Nat.*, vol. 21, no. 1, pp. 31-46, 2014.
- [3] J. Ejaz, G.A. Camer, K. Anwar and M. Asraf, "Impacts of Air Pollution Monitoring: Analysis and Histopathological Pixe Modalities in Evaluating the Relative Risks of Elemental Contamination", *Ecotoxicology*, no. 23, pp. 357-369, 2014.
- [4] A. Subandrio, Sukarman and R.T. Tambunan, "Implementation of Reclamation in PT Adaro Indonesia". Proc. of Workshop on Science and Technology Through the Forest Rescue Rehabilitation Former Coal Mine. Samarinda: Dipterocarp Research Institute, 2019.
- [5] M.U. Swift, K.A. Dvorak, K. Mulongoy, M. Musoko, N. Sangiya and G. Tian, "The Role of Soil Organisms in the Sustainable of Tropical Cropping Systems", *Soil Science and Sustainable Land Management in the Tropics*, Rimmer, ed., CAB International in Association with the British Society of Soil Science, pp. 155-172, 1994.
- [6] C.A. Siregar and T. Butarbutar, *Exposure Research Techniques Critical Land Rehabilitation and Reforestation*, Siantar: Wanariset II, Kuole, Forestry Research Institute, 1998.
- [7] Rahmawati, "Former Mine Site Restoration Ecology Based Rule", Medan: Faculty of Agriculture, University of North Sumatera, 2002.
- [8] Pratiwi, E. Santoso and M. Turjaman. "Determination of Ameliorant for Soil Improvement of Quartz Tailings as Growing Media of Forest Plants". *Journal of Forest Research and Nature Conservation*, vol. 5, no. 9, pp. 163-174, 2012.

- [9] Anonymous, "Accountability of Government Institutions (LAKIP)". Samarinda: Department of Hygiene and Yard, 2015.
- [10] R. Latifah, N. Winarsih and Y.S. Rahayu, "To Useful Organic Waste Such Fertilizer of Liquid Matter to Grow Reet Parrot (*Alternanthera ficoides*)", *Lantern Bio*, vol. 1 no. 3, pp. 139-144. September 2012.
- [11] M. Hakim, J. Wijaya and R. Sudirja, "Finding Solutions Troubleshooting Municipal Solid Waste", Bandung: Faculty of Agriculture, Padjadjaran University, cooperation with the Directorate General of Horticulture R.I. Sufficient at the Workshop "Municipal Solid Waste Management in Horticultural Development Revitalization in Indonesia", 2006.
- [12] M.M. Sutedjo, *Fertilizer and Fertilization*. Jakarta: Rinesa Cipta, 2008.
- [13] SAJADAH CO. 2018. "BAHAN-BAHAN YANG DAPAT DIJADIKAN PUPUK KOMPOS YANG HARUS ANDA KETAHUI". [HTTP://WWW.SAJADAH.CO/BAHAN-BAHAN-YANG-DAPAT-DIJADIKAN-PUPUK-KOMPOS-YANG-HARUS-ANDA-KETAHUI/](http://www.sajadah.co/bahan-bahan-yang-dapat-dijadikan-pupuk-kompos-yang-harus-anda-ketahui/). 2018.
- [14] Y. Suryani, Astuti, B. Oktavia and S. Umniyati S, "Isolasi dan Karakterisasi Bakteri Asam Laktat dari Limbah Kotoran Ayam sebagai Agensi Probiotik dan Enzim Kolesterol Reduktase", *Prosiding Seminar Nasional Biologi*, pp. 138-147, 2010.
- [15] S. Bai, M. Ravi Kumar, D.J. Mukesh Kumar, P. Balashanmugam, M.D. Balakumaran and P.T. Kalaichelvan, "Cellulose Production by *Bacillus subtilis* Isolated from Cow Dung", *Archives of Applied Science Research*, vol. 4, no. 1, pp. 269-279, 2012.
- [16] T. Liu, C.L. Ye, X.Y. Chen, W. Ran, Q.R. Shen, F. Hu and H.X. Li, "Effects of Different Organic Manure Sources and Their Combinations with Chemical Fertilization on Soil Nematode Community Structure in a Paddy Field of East China", *Ying Yong Sheng Tai Xue Bao*, vol. 24, no. 12, pp. 3508-3516, 2013.
- [17] T. Liu, X.Y. Chen, F. Hu, W. Ran, Q.R. Shen, H.X. Li and J.K. Whalen, "Carbon-rich Organic Fertilizers to Increase Soil Biodiversity: Evidence from a Meta-analysis of Nematode Communities", *Agriculture, Ecosystems & Environment*, no. 232, pp. 199-207, 2016.

Microorganisms In Ex-Coal Mining Land After The Application Of Organic Fertilizers In Saliki Village Samarinda, East Kalimantan

ORIGINALITY REPORT

7%

SIMILARITY INDEX

6%

INTERNET SOURCES

2%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	www.researchgate.net Internet Source	2%
2	www.1trajet1sourire.fr Internet Source	2%
3	Submitted to Universitas Riau Student Paper	1%
4	Ting Liu, Xiaoyun Chen, Feng Hu, Wei Ran, Qirong Shen, Huixin Li, Joann K. Whalen. "Carbon-rich organic fertilizers to increase soil biodiversity: Evidence from a meta-analysis of nematode communities", Agriculture, Ecosystems & Environment, 2016 Publication	1%
5	www.science.gov Internet Source	1%

Exclude bibliography On