

ISBN 978-602-71169-7-9

**Proceedings of
the 5th International Seminar on
New Paradigm and Innovation
on Natural Sciences and Its
Application
(5th ISNPINSA)**

**7-8 October 2015
ICT Centre,
Diponegoro University,
Semarang, Indonesia**



Table of Content

Proceedings of the 5th International Seminar on New Paradigm and Innovation on Natural Sciences and Its Application (5th ISNPINSA)	i
Preface to The Conference Proceedings	iii
Board of Reviewers	iv
Table of Content	v
Bacillus as Siderophore and Iron-bioremoval Bacteria Enny Zulaika, Septa Tri Farisna, and Nur Laili	1
Phytochemical Screening and Antibacterial Activity of Leaves Extract Balangla (<i>Litsea cubeba</i> (Lour) Pers.) from Malinau, East Borneo Hetty Manurung, Rudy Agung Nugroho, Elvi Marina	5
The Effects of Temulawak extract and Yoghurt on HDL-LDL mice blood exposed waste cooking oil Kartiawati Alipin, Walida Tanzania, Yasmi Purnamasari Kuntana	9
Drought Resistance Analysis of the North Sulawesi Local Rice Based on the Root Characters Nio Song Ai, Ludong Daniel Peter Mantilen	12
Bioavailibility of Cd, Pb, Cu, and Zn in Sediment in Garapan, Cibungur, and Ciliman Rivermouth Noverita Dian Takarina	16
Carbon Sinks of Morphologic Tree Stands in Bandung City Green Space: Case Study Taman Balai Kota, KebunBinatang, and Taman LaluLintas Ade Irma Suryani Nurvita Cundaningsih, Teguh Husodo, Herri Y. Hadikusumah	22
Effect of Growing Season on Growth and Relation of Height and Above Ground Biomass of <i>Avicennia Marina</i> Rini Budihastuti	27
Growth Improvement of Mung Bean (<i>Vigna Radiata</i> (L.) Wilczek R.) by Application of Mycofer and Phosphate Fertilizer Tia Setiawati, Mohamad Nurzaman, Asep Zainal Mutaqin, Guntur E. Adiwinata	32
Utilization of Channels Digestion Golden Snail (<i>Pomacea Canaliculata</i>) as Lytic Enzyme and Application on Yeast <i>Pichia Manshurica</i> DUC-CY15 Wijanarka, Jafron W.Hidayat, Sarjana Parman	38
Glucose Content of Sago Waste After Chloride Acid Pre-Treatment Hydrolysis For Bioethanol Production Erma Prihastanti, Widowati, Endang Kusdyantini, Agustina LINA, M.Anwar Djaelani, Priyo Sidik Sasongko, Agus Setyawan	42
Ultrastructure and Nutrient Content of Waste Sago and The Potential as Compost Block for Plant Growth Media Erma Prihastanti	45

Growth Optimization of Thermophilic Bacteria <i>Bacillus thermoamylovorans</i> and <i>Brevibacillus</i> sp. in Producing Keratinolytic Enzyme Henri Yohandini, Muharni, Eggy Lifrety Nainggolan	47
Riparian Vegetation of Suhuyon River, North Sulawesi Ratna Siahaan and Parluhutan Siahaan	51
Histological Structure of Mice (<i>Mus Musculus</i> L.) Liver after Administration of Ethanol Extract and Spinasterol from Senggugu (<i>Clerodendron Serratum</i> L) Leaves Desak Made Malini, Madihah, Euis Julaeha	55
Bacterial Colloids Silver from Slurry Of Silver Craft Industry and Its Activity as an Antibacteria Endang S. Soetarto, Fitri Nur Hidayati, Harsojo	59
<i>Bacillus</i> Resistance and Potensial as Chromium (Cr) Bioremoval Enny Zulaika, Adisya Prima, Nita Citrasari, Langkah Sembiring	65
Phytochemical Screening and Antibacterial Activity of Leaves Extract Balangla (<i>Litsea cubeba</i> (Lour) Pers.) from Malinau, East Borneo Hetty Manurung, Rudy Agung Nugroho and Elvi Marina	69
Agroforestry enhance soil moisture and fertility in rain-fed farmlands I Gede Ketut Adiputra	73
Abundance and Diversity of Coral Fish in Border Water of Unarang Reef, Nunukan, Kalimantan Utara Province Jafron W. Hidayat, Benny Diah M	77
The Improvement of Protein Content by the Use of Dried Fish Meal of <i>Oreochromis niloticus</i> in Tempeh as Aquaculture Product Diversification for Sustainable Aquaculture Lusiawati Dewi, Sapto P. Putro	83
The Use of Seaweeds <i>Sargassum</i> Sp and <i>Gracilaria Verrucosa</i> as Soil Conditioner to Enhance The Growth of <i>Vigna Radiata</i> in Sandy and Clay Soil Munifatul Izzati	86
Evaluation on The Change of Water Quality and Survival Rate of Mangrove Seedling within Silvicultural Pond at Semarang City during Early Dry Season Endah Dwi Hastuti, Rini Budihastuti	90
Magnetic Modeling of the Diwak-Derekan Geothermal Area with Extension to Bawen, Central Java Udi Harmoko, Hiska Anggit M., Tony Yulianto, Gatot Yulianto, Sugeng Widada, Achmad Widodo, Yusuf Dewantoro Herlambang, Sahid	96
A Simple Polarization for Powerful Preliminary Test of Oil Quality Level K. Sofjan Firdausi, Suryono, Priyono, Zaenul Muhlisin	100
Applications of Laser Induced Chlorophyll Fluorescence Imaging to detect Environmental Effect on Spinach Plant Minarni Shiddiq, Zulkarnain, Tengku Emrinaldi, Fitria Asriani, Iswanti Sihaloho, Heru Susanto	104

Identification Geothermal Reservoir of Telomoyo Mount from Anomaly Magnetic Data using 3D Magnetic Inversion Hiska Anggit M., Udi Harmoko, Tony Yulianto, Gatot Yulianto	109
Fabrication of NanoChiSil for Application of Fertilizer Agus Subagio, Erma Prihastanti, Ngadiwiyanana, Khasan Rowi, Ahmad Gufron	113
Synthesis Optimization of L-Aspartic acid β -hydroxamate by a novel Enzyme, β -Aspartyl- γ -glutamyl transferase Asep Awaludin Prihanto, Yuki Nonomura, Kazuyoshi Takagi, Ryosuke Naohara, Mamoru Wakayama	117
Fabrication Material Zeolite Modified by Fe with Treatment and Without High Energy Milling on Zeolite Materials Nur Farida Grafiana, Pardoyo, Agus Subagio	123
In Vitro Antioxidant Activity of Methanolic Extract of Piper retrofractum Vahl. Nurul Jadid, Sylviana R Hartanti, Nurlita Abdulgani, Wiwi Wikanta, Fitrih R Sulthoni	127
Optimization of Reaction Conditions in the Production of Gadolinium Diethylenetriamine Pentaacetate-Folate A. Mutalib, R. P. Fauzia, A. H. Gunawan, A. Anggraeni, H. Pujiastuti, R. Ukun. M.S. Soedjanaatmadja, H. H. Bahti	132
Emic and Ethic Knowledge of Bamboo's Characteristic in Process of Making Angklung Syaima Rima Saputri, Nurvita Cundaningsih, Annisa Amalia, Budi Irawan, Teguh Husodo	137
Isolation of Local Lipolytic Isolate from Domestic Compost Syifa F. Syihab, Fida Madayanti, Akhmaloka	143
Synthesis of Rice Husk-Based Zeolit using Hydrothermal Method and Its Detergent Builder Properties Alfiansyah, Arnelli, Yayuk Astuti	148
Formalin Exposure on the Rats Feeding Diet on Antioxidant Enzymatic activity and Oxidative Damage of Rats Liver Tissue Chanif Mahdi, Aulanium	154
Modification of Rice Husk-Based Activated Carbon using Sodium Lauryl Sulfat (SLS) for Lead (Pb) Ions Removal Dewi Reskiandini, Arnelli, Yayuk Astuti	159
Comparative study of encapsulated rhizome extract of <i>Alpinia purpurata</i> (Zingiberaceae) in alginate and alginate-chitosan Meiny Suzery, Dian Majid, Bambang Cahyono	165
Novel Archaeal DNA Polymerase B from Domas Hot Spring West Java Suharti, Rukman Hertadi, Fida Madayanti Warganegara, Santi Nurbaiti, Akhmaloka	169
The Effect of Configuration to Interaction Energy Between The Segments of Chitosan and Ascorbic Acid Molecule: Theoretical Study of Drug Release Control Suci Zulaikha Hildayani, Parsaoran Siahaan	172

Analyze of Classification Accaptence Subsidy Food Using Kernel Discriminant Alan Prahutama, Moch. Abdul Mukid	177
Volatility Modelling Using Hybrid Autoregressive Conditional Heteroskedasticity (ARCH) - Support Vector Regression (SVR) Hasbi Yasin, Tarno, Abdul Hoyyi	183
Quality Function Deployment and Fuzzy TOPSIS Methods in Decision Support System for Internet Service Provider Selection Novianto Dwi Prasongko, Bayu Surarso, Rahmat Gernowo	188
Automatic Speech Recognition for Indonesian using Linear Predictive Coding (LPC) and Hidden Markov Model (HMM) Sukmawati Nur Endah, Satriyo Adhy, Sutikno, Rizky Akbar	194
Mathematical Modeling of worm infection on computer in a Network: Case study in the Computer Laboratory, Mathematics Department, Diponegoro University, Indonesia Nurfitriani S., Widowati, Robertus H.	200

Phytochemical Screening and Antibacterial Activity of Leaves Extract Balangla (*Litsea cubeba* (Lour) Pers.) from Malinau, East Borneo

Hetty Manurung^{1,a}, Rudy Agung Nugroho^{1,b} and Elvi Marina^{1,c}

¹Department of Biology, Faculty of Mathematics and Natural Sciences, Mulawarman
University, Samarinda. Indonesia

^a hetty_manroe@gmail.com, ^b rudyagung.nugroho@fmipa.unmul.ac.id,

^c elvimarina92@gmail.com

Abstract. Balangla (*Litsea cubeba*) plant is used as traditional medicine by Dayak Kenyah tribe in East Kalimantan. It contains active compounds that are efficient to treat many human diseases and believed to have an antibacterial activity. The purposes of this study were to determine the phytochemical compounds of the balangla leaves and to investigate the antibacterial activity of ethanol extract of leaves *L.cubeba*. Respectively, various levels of ethanol extract of *L.cubeba* leaves viz: 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% were used to examine its antibacterial activity against bacteria gram positive (*Staphylococcus aureus*) and gram negative (*Escherichia coli*) by using pitting diffusion method. The results indicated that alkaloids, flavonoids, phenols and steroids have been found as phytochemical compounds in the ethanol extracts of *L. cubeba* leaves. Meanwhile, the antibacterial activities of ethanol extracts of *L. cubeba* leaves against the test organisms had been determined and significantly inhibited the growth of *S. aureus* and *E.coli*, forming a wide inhibition zone (15.91 ± 0.950 mm) for *S. aureus* and (16.23 ± 0.416 mm) for *E. coli*. Further, antibacterial activity of (*L.cubeba*) *in-vitro* had been justified on its utility in traditional medicines for the treatment of infections of bacterial origin.

Keywords: Balangla (*Litsea cubeba*), Lauracea, Antibacterial, Phytochemical Screening.

Introduction

Litsea cubeba (Lours.) Pers., locally known as Balangla belongs to the Lauraceae plant family. *L. cubeba* is a 3- to 10 m evergreen tree or shrub widely distributed in Southeastern Asia, Southern China, Japan, and Taiwan [1,2]. In Indonesia, *L.cubeba* distributed in Sumatera, Java, and Borneo. This plant is one of the common medicinal plants in Borneo. *L. cubeba* can be used as a flavoring or herbal medicine. As a flavoring, it gives a unique flavor resembling that of a mixture of pepper, ginger, and citrus. It is popular as a flavor enhancer in foods, cosmetics, and cigarettes [3,4]. Balangla by the people of East Kalimantan especially Dayak Kenyah tribe used as traditional medicine to cure for coughs, colds, migraine headaches, skin diseases, pain relieve, diarrhea, fever, aromatherapy, and even benefit as seasoning (spices) and believed to have an antibacterial activity.

Recent study has been reported that an aqueous EtOH extract of the barks of *L.*

cubeba has yielded five novel isoquinoline alkaloids, namely (+)-*N*-(methoxycarbonyl)-*N*-nordicentrin, (+)-*N*-(methoxycarbonyl)-*N*-norpredicentrine, (+)-*N*-(methoxyl-carbonyl)-*N*-norbulbodione, and (+)-*N*-(methoxycarbonyl)-*N*-norisocorydione, and (+)-8-methoxyl-isolaurenine *N*-oxide, and one known compound, (+)-*N*-(methoxycarbonyl)-*N*-norglaucine [5]. Another study [6] showed that the essential oil in the leaves of *L.cubeba* containing sineol, citronellol, oinen alpha, beta-pinene, and limonene citronellal. Previous studies revealed that essential oil and bioactive compound were shown potential as an anticancer, antioxidant, and antibacterial. Antibacterial compounds usually found in plant parts such as leaves, twigs, bark, and other parts. Antibacterial is usefull to eliminate bacteria potentially harmful bacteria to health and pathogens such as *Staphylococcus aureus* and *Escherichia coli*. *E. coli* generate bladder infections and diarrhea [7] while *S. aureus* can cause purulent infection. *S. aureus* led to various types of diseases such as infections of

the polikel hair, sweat glands, ulcers, skin infections, and wound infections [8]. The purposes of this study were to determine the phytochemical compounds that found in the balangla leaves and to investigate the antibacterial activity of ethanol extract of leaves *Litsea cubeba*.

Material and Methods

Sample Collection

The leaves of *L. cubeba* were collected locally from Gunung Seribu (Can batu) Mahak Baru, district Sungai Boh Malinau, East Borneo, in December 2014. The plants collected were identified botanically in Plant Anatomy and Taxonomy Laboratory, Biology Departement Mulawarman University and deposited in the herbarium for future reference.

Preparation of plant material

The fresh leaves were washed with tap water and then thoroughly cleaned with distilled water and shade dried for a week. Then the dried leaves (1500 g) were grinded to a fine powder by using blender. The powder was taken and macerated with 95% EtOH. They were kept at room temperature for 5 days. Thereafter the mixtures were filtered by using Whatmann filter paper no .1. The supernatant were pooled together, concentrated in rotary evaporator at 40°C. The dried extract (57.283 g) was used directly for the determination of a presence of phytochemicals and antibacterial activity.

Qualitative Analysis of Phytochemicals

Phytochemical examinations were carried out for leaves extract *L. cubeba* as per the standard methods [9].

Test for Alkaloids

Dragendroff's test: A-2 mL of extract was added with 5 mL of chloroform-ammonia 0.005 M, then homogenized and filtered. The filtrate was added with a few drops of 2M sulfuric acid and shaken to form two layers of acids and bases. The layer acid (found on the top layer), added with a few drops of reagent Dragendroff's. Formation of deposits of red-brown color indicated the presence of alkaloids.

Test for Flavonoids

A-2 mL crude extract was added with 5 mL of water, boiled for 5 minutes and filtered. Two mL filtrate was added with 0.05 mg of Mg powder and 1 mL chloride acid, then shaken until homogeneous. A yellow or red colouration indicates the presence of flavonoids.

Test for phenols

Lead acetate test: To 2 mL of the extract, few mL of 1 % lead acetate solution was added. The formation of bluish black precipitate indicated the presence of tannins and phenolic compounds.

Test for saponins

Frothing test: A-2 mL of filtrate was diluted with 5 mL of distilled hot water and the mixture was shaken vigorously and observed for persistent foam which lasted for atleast 10 mins which indicated the presence of saponins.

Test for Terpenoids

Liebermann - Burchard's test: Extract was treated with a few drops of acetic anhydride, boiled and cooled. Concentrated sulfuric acid was added from the sides of the test tube which showed a brown ring at the junction of two layers, and the formation of deep red color indicated the presence of terpenoids.

Test for Antimicrobial Activity

Source of Bacterial

Two microbial isolates were chosen for antimicrobial investigation: Gram positive bacteria (*S. aureus*) and Gram negative bacteria (*E. coli*). Both microbial were purchased from Laboratory of Microbiology, Faculty of Sciences Mulawarman University, Samarinda, Indonesia.

Testing for Antibacterial Activity

The test organisms were sub cultured by streaking them on nutrient agar, followed by incubation for 20 hr at 37 °C. These were subcultured prior to each experiment and were used for antibacterial. The pitting diffusion method of two species of bacteria *Staphylococcus aureus* and *Escherichia coli* was performed in *in-vitro* antibacterial

activity. Medium that was used in the antimicrobial activity was MHA (Mueller Hinton Agar), an inoculum containing 10^7 bacteria/mL was incorporated. After the suspension of bacteria seep into the media, then created pit in the media using the sterile iron cork borer (6 mm diameter). Using sterilized dropping pipettes, different concentrations (10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and 100% /well) of leaves extract *L.cubeba* was carefully added into the wells. The plates were then incubated at 37 °C for 24–48 hr. Chloramphenicol was used as positive control for antibacterial activities while DMSO was used as the negative control. The diameter of the inhibition zone was measured for antibacterial activities testing. Experiments were performed in triplicate, and the results were presented as the mean values of the diameters of the inhibitory zones from three runs. The antimicrobial activity was evaluated by measuring the diameter of inhibition zone [10,11]. The diameters of the inhibitory zones value were used as criteria to judge the antimicrobial activity (strong active: the diameters of the inhibitory zones ≥ 20 mm, active: the diameters of the inhibitory zones 10-20 mm, moderately active: the diameters of the inhibitory zones 5-10 mm, not active: the diameters of the inhibitory zones are invisible or ≤ 5 mm) [12].

Results and Discussion

Phytochemical Analysis

The results of the phytochemical analysis were carried out in extracts of *Litsea cubeba*. The experiment showed the presence of secondary metabolites such as alkaloids, flavonoids, phenols and steroids. The results of the phytochemical analysis of *L.cubeba* are shown in Table 1.

Tabel 1. Qualitative phytochemical analysis of leaves extract *L.cubeba*

No	Parameters	Leaves extract
1	Alkaloid	+
2	Flavonoid	+
3	Phenol	+
4	Saponin	-
5	Steroid	+

(+indicates presence, - indicates absence)

The presence of alkaloids in leaves extract is likely to be responsible for antimicrobial activity effects observed. Alkaloid is one of the phytochemical compounds in *Litsea cubeba* that act as an antibacterial activity [13,14].

Antimicrobial Activity

Antimicrobial activity is used to test whether the leaf extract has capability to control the growth of the bacterial. The zone of inhibition had been obtained for both bacteria *Staphylococcus aureus* and *Escherichia coli*. The result obtained, the zone of inhibition was recorded at ten concentrations of 10,20,30,40,50,60,70,80,90, and 100% in Table 2. In general Table 2 showed that increasing concentrations of leaf extracts enhanced inhibitory zone on both bacteria. The extracts displayed relative antimicrobial activities against bacterial tested with the diameter of inhibition zones ranging between 9.58 ± 0.36 to 15.91 ± 0.95 mm (*S. aureus*) and 9.63 ± 0.32 to 18.63 ± 0.18 mm (*E. coli*).

Tabel 2. Antimicrobial activity of leaves extract of *Litsea cubeba*

Sample extract Conc.(%)	Zone inhibition (mm)	
	<i>Stapylococcus aureus</i>	<i>Escherichia coli</i>
10%	9.58 ± 0.36	9.63 ± 0.32
20%	10.71 ± 0.62	10.65 ± 0.75
30%	11.36 ± 0.41	11.07 ± 0.40
40%	12.40 ± 0.40	12.41 ± 0.18
50%	$12,71 \pm 0.70$	13.62 ± 0.11
60%	13.06 ± 0.29	13.16 ± 0.68
70%	13.63 ± 0.27	14.00 ± 0.37
80%	14.03 ± 0.81	14.56 ± 0.66
90%	13.80 ± 0.69	15.08 ± 0.32
100%	15.91 ± 0.95	15.23 ± 0.41
Chloramphenicol (25%)	18.33 ± 0.10	18.63 ± 0.18

The highest antibacterial activity in both bacteria was found at 100 % concentration and the lowest at 10 % concentration of leaves extract. The results obtained was indicated of the presence of broad spectrum antimicrobial compounds or metabolic toxins in *L.cubeba* leaves extract that could be exploited in treating infections associated with the aforementioned bacterial. Previous

studies on antimicrobial activities of medicinal plants indicated that inhibition zones of 10 mm or greater were taken to represent good activity of such plants [15,16].

Conclusion

Balangla leaf extract can be used as a source of secondary metabolites such as alkaloids, Flavanoids, Phenol, and steroids. The results suggest that the antibacterial activity of leaves extract *L. cubeba* may contribute to prevent some of the diseases caused by *S. aureus* and *E. coli*.

References

- T.T. Liu, T.S. Yang. Antimicrobial impact of the components of essential oil of *Litsea cubeba* from Taiwan and antimicrobial activity of the oil in food systems. *Int. J. Food Microbiol.* 2012, 156, 68-75.
- C.H. Huang, W.J. Huang, S.J. Wang, P.H. Wu, W.B. Litebamine, a phenanthrene alkaloid from the wood of *Litsea cubeba*, inhibits rat smooth muscle cell adhesion and migration on collagen. *Eur. J. Pharmacol.* 2008, 596, 25-31.
- Y. Wang, Z.T. Jiang R. Li. Complexation and molecular microcapsules of *Litsea cubeba* essential oil with β -cyclodextrin and its derivatives. *Eur. Food Res. Technol.* 2009, 228, 865-873.
- M. Luo, L.K. Jiang, G.L. Zou. Acute and genetic toxicity of essential oil extracted from *Litsea cubeba* (Lour.) Pers. *J. Food Prot.* 2005, 68, 581-588.
- W. Zhang, J-F. Hu, Lv. Wen-Wen, Q-C. Zhao, G-B. Shi. Antibacterial, Antifungal and Cytotoxic Isoquinoline Alkaloids from *Litsea cubeba*. *Molecules* 2012, 17, 12950-12960.
- Y.N. Heryati, A.S. Mindawati, Kosasih. Profek Pengembangan Lemo (*Litsea cubeba* L. Persoon) Di Indonesia. *Jurnal Tekno Hutan tanaman. Pusat penelitian Hutan Tanaman.* 2009, 2(1):9-17.
- Books, F.G., Butel, J.S., dan Morse, S.A. 2001. *Mikrobiologi Kedokteran (Edisi Terjemahan)*. Selemba Medika. Jakarta.
- I. Entjang. *Mikrobiologi & Parasitologi untuk Akademi Keperawatan dan Sekolah Tenaga Kesehatan yang Sederajat.* 2003. PT. Citra Aditya Bakti.
- J.B. Harbone. *Phytochemical Methods*. London. Chapman and Hall Ltd, 1973, pp 49-188.
- E.G. Trease, W.C. Evans. *Textbook of Pharmacognosy*, 3 rd edn. Bailliere Tindal, London pp.81-90,268-98,(1989).
- J. Owolabi, E.K.I. Omogbai, O. Obasuyi. Antifungal and antibacterial activities of ethanolic and aqueous extract of *Kigelia africana* (Bignoniaceae) stem bark. *African Journal of Biotechnology*, 2007, 6(14):882-85.
- W.W. Davis, T.R. Stout. *Disc Plate Methods in Microbiological Antibiotic Assays*. *Microbiology.* 1971, 22(4): 659-665.
- T. Feng, Y. Xu, X.N. Cay, Z.Z. Du, X.D. Liu. Antimicrobially active isoquinoline alkaloids from *Litsea cubeba*, *Planta med.* 2009, 75: 76-79.
- W.A. Volk, M.F. Wheeler. *Bacteriology*. Philadelphia. Lippincott, 1973.
- C.A. Bukola, A.A. Onilude. Screening of lactic acid bacteria strains isolated from some nigerian fermented foods for EPA production. *World Applied Science Journal.* 2008, 4(5):741-747.
- N. Ramesh, M.C.Sathyanarayana, H.Lloyd. Abundance of grey junglefowl *Gallus sonneratii* at Theni Forest Division, Western Ghats, India: implications in monitoring and conservation. *2011 International Journal of Galliform Conservation* 2, 14-21.

**Proceedings of
the 5th International
Seminar on New Paradigm
and Innovation on Natural
Sciences and Its Application
(5th ISNPINSA)**



**Faculty of Sciences and Mathematics
Diponegoro University**