

Pathogenicity assay of probiotic-potential bacteria from the Kelabau fish (*Osteochilus melanopleurus*)

by Agustina Agustina


Submission date: 06-Jan-2023 02:07PM (UTC+0700)

Submission ID: 1989116310

File name: Pathogenicity_assay_of_probiotic.pdf (616.87K)

Word count: 4989

Character count: 26249



13

Pathogenicity assay of probiotic-potential bacteria from the Kelabau fish (*Osteochilus melanoleurus*)

^{1,2}Agustina, ³Slamet B. Prayitno, ⁴Agus Sabdono, ¹Gina Saptiani

19

¹ Aquatic Microbiology Laboratory, Faculty of Fisheries and Marine Science, Mulawarman University, Samarinda, Indonesia; ² Doctorate Coastal Resources Management, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia;

³ Aquaculture Study Program, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia; ⁴ Marine Science Study Program, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia. Corresponding author: Agustina, agustina197748@gmail.com

13

Abstract. This study aimed to evaluate the pathogenicity of probiotic-potential bacteria from the gut of Kelabau fish (*Osteochilus melanoleurus*) collected from Melintang Lake, East Kalimantan Province, Indonesia. Common carp (*Cyprinus carpio*) juveniles were injected with 0.1 mL (10^6 CFU mL $^{-1}$) bacterial suspension of BP1, BP3, BP s_2 , BK, *Aeromonas hydrophila*, and phosphate buffer saline, respectively. The injected fish were maintained in 20 L aquarium at a density of 10 individuals/aquarium for 7 days of experiment. Clinical sign, mortality, hematological parameters, histopathology of gills and kidney were observed. The ANOVA test showed that probiotic-potential bacteria from gut of Kelabau fish had a significant effect on the survival rate and hematological parameters of the *C. carpio* juveniles. The infected fish showed changes in clinical signs such as hemorrhage, edema, and ulcer. Histopathological analysis indicated gill hemorrhage, necrosis, hyperplasia, edema and epithelial lifting, kidney necrosis, melanosis and vacuolization in nephron tubular cells. BP s_2 isolate provided the best survival rate of 93% and it is a safe option for *C. carpio* as a probiotic-potential bacteria.

Key Words: survival rate, histopathological analysis, common carp, *Aeromonas hydrophila*.

36

Introduction. Efforts to increase fish production through freshwater fish farming in East Kalimantan Province, especially at Kutai Kartanegara Regency are constant. Kutai Kartanegara Regency was the most important fish producer in East Kalimantan, it delivered 140,133.70 tonnes (BPS 2017), freshwater fish representing 76.28% (106,900.3 tonnes). However, diseases are still a major obstacle in freshwater fish farming in East Kalimantan. The use of local natural ingredients to overcome infectious diseases is an alternative to reduce the use of chemicals and antibiotics. Probiotics are safer ingredients providing benefits like improving the farmed fish health status and being environmentally friendly (Ouwehand et al 2002; Maslowski & Mackay 2010). Screening of potential probiotic bacteria from fish and aquaculture environments in this area has already been carried out and showed positive results (Agustina et al 2010; Agustina et al 2018).

14

Several types of bacteria obtained from the gut of Kelabau fish (*Osteochilus melanoleurus*) showed the ability to inhibit the growth of two pathogenic bacteria aquaculture fish in the Kutai Kartanegara Regency, namely *Aeromonas hydrophila* and *Pseudomonas* sp. (Agustina et al 2018). Previous tests of several types of bacteria were able to inhibit pathogenic bacteria in vitro, both in the form of whole bacterial cells and cellular product components. Products in the form of probiotic potential bacteria can be produced after a series of follow-up tests in vivo. Pathogenicity assay of intestinal bacteria with demonstrated abilities to inhibit pathogenic bacteria in vitro needs to be further carried out, in order to guarantee the safety of probiotics candidate. Therefore, it is necessary to evaluate their non-pathogenicity in the host (Giri et al 2012).

2

Based on the background above mentioned, pathogenicity assay was carried out to four selected bacterial isolates from previous tests, namely BP1, BP3, BP_s2 and BK with common carp (*Cyprinus carpio*) juveniles as experimental animals, due to its local frequency. This test aimed to evaluate the pathogenicity of the four bacterial isolates and obtain non-pathogenic bacterial isolates that can be used as potentially probiotic bacteria. The health status of *C. carpio* juveniles was checked post infection. The observed parameters were clinical signs, survival rate, hematological parameters and histopathology of the gill and kidney.

Material and Method

Experimental fish. Healthy *C. carpio* juveniles were obtained from a common carp hatchery in the Loa Kulu District, Kutai Kartanegara Regency, East Kalimantan Province, Indonesia. The fish was not showing any signs of disease (inspected through a gross examination of the skin, fins and gills of the respective samples) with no previous history of parasitic and bacterial infection, and having an average weight of 2.6 ± 0.2 g. Fish were maintained in aerated freshwater for 5 days in 40 L glass aquarium. All the fishes were fed three times a day to satiation with pellet containing 38% protein during the acclimatization period.

Preparation of bacteria. Isolates BP1, BP3, BP_s2, BK and *A. hydrophila* (AH-1) obtained from a previous test (Agustina et al 2018) were grown for 24 hours in tryptone soy broth media in incubator at 30°C. The harvested cells were washed, resuspended in phosphate buffer saline (PBS). Each bacterial suspension (0.1 mL) contained approximately 10^6 CFU mL⁻¹, as determined by the standard plate count method (Gupta et al 2014).

5

Determination of possible harmful effects of the bacteria on fish. Each bacterial suspension (0.1 mL) of BP1, BP_s2, BP3 and BK were injected intramuscularly into separate groups of 10 *C. carpio*. Another group of 10 fish were injected with phosphate buffer saline as negative control and other 10 fish were injected with 0.1 mL *A. hydrophila* AH-1, as positive control. Once the treatment was performed, each experimental group of fish was reared in 20 L aquarium at a density of 10 fish/aquarium. All experimental fishes were fed three times a day with commercial pellet containing 38% protein for 7 days. Water quality was maintained as follows: temperature 27-29°C, pH 6.4-7.1, dissolved oxygen 6.4-8.16 mg L⁻¹ with 7.67-7.95% oxygen saturation and ammonia of 0.0001-0.0035 mg L⁻¹. Half of the water of each aquarium was renewed daily with aerated tap water. Daily observation was recorded for disease incidence and mortality during a 7 days experimental period.

Experimental design. Six experimental treatments were carried out using completely randomized design (CRD). *C. carpio* juveniles were randomly selected and placed into eighteen aquaria with a stocking density of 10 individuals/aquarium. The experiments were run in triplicates. The treatments consisted of an injection of 0.1 mL PBS as a negative control (T0), an injection of *A. hydrophila* AH-1 as a positive control (T1), and four injections of gut bacteria of *O. melanopleurus*, namely BP1 (T2), BP_s2 (T3), BP3 (T4) and BK (T5).

Survival rate. Survival rate of the experimental fish (*C. carpio*) was carried out by calculating the number of survivors divided by the total number of initial fish, times 100%. Afterwards, each treatment was tabulated and analysed accordingly.

Haematological parameters. Fish blood parameters were observed three days, five days and seven days post injection. The first step was to take blood samples from the caudal vein using a syringe. Assessments of the blood parameters were done for the total red blood cells and total white blood cells following the Blaxhall & Daisley (1973) procedure, hemoglobin content was assessed using the Sahli method with a haemometer

(Wedemeyer & Yasutake 1977), and hematocrit was assessed according to the method described by Anderson & Siwicki (1995).

Clinical sign. The clinical sign of experimental fish at each treatment were observed post injection treatments. The clinical signs of the experimental fish as a result of treatments were compared with each other. The clinical signs observed included the presence of hemorrhage, edema and ulcer.

Histopathology. A histological examination of the internal organs was performed at the end of the experiment. Samples of gills and kidney were fixed in a buffered neutral formalin solution (10%). The specimens were performed according to Roberts (2001). Tissue sections were stained with hematoxylin-eosin (HE) and observed under Olympus CX23 microscope, photographed using an Olympus SZ-14 camera to study the pathological changes.

Data analysis. The experimental data were tabulated using Microsoft Office Excel 2016 and further analyzed with ANOVA (analysis of variance) using the application of SPSS 14.0 at a 95% confidence level. If the ANOVA showed a significantly different result, a Tukey post hoc test was performed. The parameters of clinical sign, histopathology and water quality were descriptively analyzed.

Results

Survival rate. The survival rate of the experimental fish injected by two gut bacteria, treatments (T3 and T4) were significantly different compared to the positive control (T1) ($p<0.05$) (Table 1). The survival rates of the fish which received T2 and T5 were significantly lower than in the case of the fish treated with T3 and T4. The highest survival rate was obtained by treatment T3 (*C. carpio* juveniles injected by BP_{s2} isolate, Table 1).

Table 1
Survival rate of *Cyprinus carpio* juveniles one week post pathogenicity test

Treatment	Cumulative mortality (fish)							Survival rate (%)
	1	2	3	4	5	6	7	
T0	0	0	0	1	1	1	1	96.67±5.77 ^b
T1	7	11	15	17	18	18	18	40.00±10.00 ^a
T2	7	14	17	18	18	18	18	40.00±10.00 ^a
T3	0	1	2	2	2	2	2	93.00±11.55 ^b
T4	1	5	8	9	9	9	9	70.00±10.00 ^b
T5	8	12	15	17	19	19	19	36.67±15.28 ^a

T0 (negative control), T1 (Positive control), T2 (BP1), T3 (BP_{s2}), T4 (BP3), T5 (BK). Data are presented as mean ± SD. The different superscript letter within same column shows a significantly different effect ($p<0.05$).

Total erythrocyte. Total erythrocyte of all treatments tended to decrease post injection except the negative control (To). Total erythrocyte of probiotic-potential bacteria (T2, T3, T4, T5) showed a significant difference ($p<0.05$) compared to the negative control (To) (Table 2). The erythrocyte on treatment T3 (BP_{s2}) was significantly higher ($p<0.05$) compared to the other treatments after three days post injection. However, on day 5 and 7, the negative control (T0) showed the highest total erythrocyte.

Table 2
Total erythrocyte of *Cyprinus carpio* juveniles at the beginning of stocking, day 3, 5 and 7 post bacterial injection

Treatment	Total erythrocyte ($\times 10^6$ cell mm^{-3})			
	Initial	Day 3	Day 5	Day 7
T0	1.33±0.04	1.58±0.03 ^d	1.72±0.04 ^d	2.03±0.02 ^f
T1	1.33±0.04	1.01±0.04 ^c	0.86±0.01 ^b	0.75±0.01 ^b
T2	1.33±0.04	1.10±0.04 ^c	0.88±0.01 ^b	1.48±0.01 ^e
T3	1.33±0.04	1.66±0.05 ^d	0.84±0.05 ^b	0.87±0.01 ^c
T4	1.33±0.04	0.84±0.02 ^b	1.06±0.02 ^c	1.36±0.01 ^d
T5	1.33±0.04	0.65±0.05 ^a	0.46±0.02 ^a [10]	0.61±0.01 ^a

T0 (negative control), T1 (Positive control), T2 (BP1), T3 (BPs2), T4 (BP3), T5 (BK). Data are presented as mean ± SD. The different superscript letter within same column shows a significantly different effect ($p<0.05$).

Total leukocyte. The total leukocyte in the negative control T0, T3 (BPs2) and T5 (BK) increased significantly ($p<0.05$) compared to T1 (positive control) at day 3 after the bacterial injection. T4 (BP3) showed the highest total leucocyte in day 5 and 7, whilst T3 (BPs2) significantly decreased ($p<0.05$) compared to T1 (positive control) in day 7 post bacterial injection. The total leucocyte of the experimental fish is presented in Table 3.

Table 3
Total leucocyte of *Cyprinus carpio* juveniles at the beginning of stocking, day 3, 5 and 7 post bacterial injection

Treatment	Total leucocyte ($\times 10^4$ cell mm^{-3})			
	Initial	Day 3	Day 5	Day 7
T0	8.48±0.09	9.24±0.06 ^c	7.84±0.03 ^b	6.46±0.13 ^a
T1	8.48±0.09	6.54±0.11 ^a	8.71±0.12 ^c	12.15±0.05 ^d
T2	8.48±0.09	7.83±0.10 ^b	5.71±0.18 ^a	7.50±0.01 ^b
T3	8.48±0.09	9.59±0.10 ^d	11.35±0.02 ^d	8.47±0.13 ^c
T4	8.48±0.09	6.49±0.04 ^a	11.79±0.18 ^e	15.83±0.26 ^e
T5	8.48±0.09	9.99±0.01 ^e	11.16±0.36 ^d [10]	15.50±0.46 ^e

T0 (negative control), T1 (Positive control), T2 (BP1), T3 (BPs2), T4 (BP3), T5 (BK). Data are presented as mean ± SD. The different superscript letter within same column shows a significantly different effect ($p<0.05$).

Hemoglobin level. The hemoglobin level of the experimental fish in T3 (BPs2) on day 3 post bacterial injection was significantly different ($p<0.05$) compared to the control treatments (T0) (Table 4). In day 5 post bacterial injection, the hemoglobin level of the experimental fish was not significantly different ($p>0.05$) in all the treatments, except at T0 (negative control). The hemoglobin level of T2 (BP1), T3 (BPs2) and T4 (BP3) in day 7 post bacterial injection increased significantly compared to T1 (positive control) (Table 4).

Table 4
Hemoglobin level of *Cyprinus carpio* juveniles at the beginning of stocking, day 3, 5 and 7 post bacterial injection [33]

Treatment	Hemoglobin level ($g dL^{-1}$)			
	Initial	Day 3	Day 5	Day 7
T0	3.97±0.47	3.92±0.14 ^c	4.83±0.14 ^b	7.00±1.00 ^d
T1	3.97±0.47	2.83±0.14 ^a	2.58±0.38 ^a	2.42±0.14 ^{ab}
T2	3.97±0.47	3.25±0.25 ^{ab}	3.17±0.29 ^a	3.58±0.29 ^{bc}
T3	3.97±0.47	3.33±0.14 ^b	2.67±0.29 ^a	3.42±0.14 ^{bc}
T4	3.97±0.47	2.92±0.14 ^{ab}	2.92±0.14 ^a	4.04±0.29 ^c
T5	3.97±0.47	2.92±0.14 ^{ab}	2.75±0.43 ^a [10]	2.17±0.14 ^a

T0 (negative control), T1 (Positive control), T2 (BP1), T3 (BPs2), T4 (BP3), T5 (BK). Data are presented as mean ± SD. The different superscript letter within same column shows a significantly different effect ($p<0.05$).

Hematocrit. Three days after the treatments, T1 (positive control) had a lower level of hematocrit compared to the other treatments. T4 (BP3) and T5 (BK) had a significantly lower level of hematocrit ($p<0.05$) compared to T0 (negative control) (Table 5). The hematocrit of the experimental fish was not significantly different at five days after the treatments ($p>0.05$). Whereas, seven days after the treatments, the hematocrit level of T3 (BPs2) increased compared to the other treatments, except T0 (negative control), yet it was not significantly different ($p>0.05$) (Table 5).

Table 5
The hematocrit level of *Cyprinus carpio* juveniles at the beginning of stocking, day 3, 5 and 7 post bacteria injection

Treatment	Hematocrit (%)			
	Initial	Day 3	Day 5	Day 7
T0	22.83±0.50	20.33±1.26 ^b	18.5±1.32 ^a	20.83±0.76 ^b
T1	22.83±0.50	16.33±0.76 ^a	15.17±1.26 ^a	17.33±0.76 ^{ab}
T2	22.83±0.50	18.33±1.26 ^{ab}	17.67±1.53 ^a	18.50±0.50 ^{ab}
T3	22.83±0.50	17.67±1.53 ^{ab}	17.83±1.44 ^a	19.50±0.87 ^{ab}
T4	22.83±0.50	16.83±1.26 ^a	16.33±1.26 ^a	17.83±1.04 ^a
T5	22.83±0.50	16.83±0.76 ^a	16.33±1.26 ^a	17.67±1.15 ^a

T0 (negative control), T1 (Positive control), T2 (BP1), T3 (BPs2), T4 (BP3), T5 (BK). Data are presented as mean ± SD. The different superscript letter within same column shows a significantly different effect ($p<0.05$).

Clinical signs. The clinical signs found in the experimental fish injected with *O. melanoleurus* gut bacteria were edema and hemorrhage. In addition to these two signs, fish injected with *A. hydrophila* have also experienced ulcer. In general, the clinical signs were seen 36 hours after treatments. The clinical signs of the experimental fish are presented in Figure 1.




Figure 1. (A) Clinical sign of fish in negative control. (B) Fish injected with gut bacteria of *Osteochilus melanoleurus* (BK) showed edema. (C) Fish injected with gut bacteria of *Osteochilus melanoleurus* (BP1) showed hemorrhage (D) Fish injected with *Aeromonas hydrophila* AH-1 showed ulcer.

Histopathology. The observed histopathological section of gill showed hemorrhage, necrosis, hyperplasia, edema and epithelial lifting, whilst the kidney showed necrosis,

melanism and vacuolization in the nephron tubular cells. The histopathological section of gill and kidney of the experimental fish are presented in Figure 2 and Figure 3.




Figure 2. (A) Gill of *Cyprinus carpio* juvenile injected with *Aeromonas hydrophila* AH-1 showed edema (Ed) and epithelial lifting (El) occurred. (B) Gill of *Cyprinus carpio* juvenile injected with gut bacteria of *Osteochilus melanopleurus* (BK) showed hemorrhage (He), necrosis (Ne) and hyperplasia of epithelial cells (Hp) that cover the surface of secondary lamellae (HE stain 100x).




Figure 3. (A) Kidney of *Cyprinus carpio* juvenile injected with *Aeromonas hydrophila* AH-1 showed haemopoietic cells undergo necrosis (Ne) and vacuolization (Va) in nephron tubular cells. (B) Gill of *Cyprinus carpio* juvenile injected with gut bacteria of *Osteochilus melanopleurus* (BK) showed melanosis (Me) in kidney cells (HE stain 100x).

Discussion. The gut bacteria of *O. melanopleurus* that were injected into *C. carpio* juveniles showed different levels of pathogenicity. The T3 (BPs2) gave the best results on the survival rate (93.00 ± 11.55) compared to the positive control (40.00 ± 10.00). The differential levels of resistance in fish could be determined by estimating their survival rate after being infected with bacteria (Wassom & Kelly 1990). In this study, *C. carpio* juveniles had the highest resistance to T3 (BPs2) compared to other treatments especially the positive control (Table 1). A similar result was obtained by Agustina (2007), the probiotic-potential bacteria from catfish (*Clarias* sp.) gut also demonstrated low pathogenicity to catfish juveniles with a survival rate of 85-90%. Moreover, bacteria

of T2 and T5 were suspected to be pathogenic in *C. carpio* juveniles due to a high mortality after injection. Bacteria that have the potential to cause death, both primary and secondary infections, could not be ignored and were thought to cause systemic infections (Burbank et al 2012).

The total erythrocyte of *C. carpio* juveniles decreased after the treatments until the fifth day and increased on the seventh observation day, except on the positive control. The total erythrocyte in the T2, T3, T4 and T5 treatments were significantly higher than in the positive control on day seven after injection (Table 2). The total erythrocyte in the positive control significantly decreased because *A. hydrophila* had hemolysin that could cause erythrocyte lysis (Thune et al 1993; Shao et al 2004). The total leucocytes fluctuated or increased in the infected fish. The total leucocyte on the seventh day of observation decreased in the negative control, T2 and T3 treatments, whereas T1, T4 and T5 were increased (Table 3). Similar to the total leucocytes in T1, T4 and T5 in this study, Li et al (2016) found that the number of white blood cells of grass carp *Ctenopharyngodon idella* increased significantly at 7, 14, 21 days post infection with *A. hydrophila* in the treatment group.

Hemoglobin is a pigment in the erythrocyte and has a function to bind oxygen to be further distributed throughout the body (Reece et al 2014). The hemoglobin level of *C. carpio* juveniles decreased five days after being injected with *O. melanopleurus* gut bacteria T2-T5 (Table 4). The hematocrit levels after the injection with bacteria were not significantly different compared to the positive control (Table 5). In this experiment, a gradual decrease in the total erythrocyte, hemoglobin and hematocrit levels after the injection indicated a damage of the haematopoietic tissues including kidney. This might cause haemorrhagic septicaemia. This could be seen from the haemorrhages on dorsal body parts as well as sero-sanguinous fluid that filled the peritoneal cavity. Decreased blood red cell values, hemoglobin and hematocrit levels indicated that red blood cells were being destroyed by the leucocytosis activity in erythrocytic anemia with subsequent erythroblastosis (Haney et al 1992). Similar to our findings, significant increase in total leucocytes and decrease in the total erythrocyte compared to the control fish was found in *Puntius sarana* infected with *A. hydrophila* (Das et al 2011).

The dorsal part of a *C. carpio* showed hemorrhage 36 hours after treatment. Edema appeared two days after the injection with *A. hydrophila* (positive control) and some *C. carpio* had ulcers on the injection site (Figure 1). Commonly, fish juvenile gills and kidney in this study showed some changes when observed in histological examination. Fish gills showed hemorrhage, necrosis, hyperplasia, edema and epithelial lifting (Figure 2), while kidney experienced necrosis, melanosis and vacuolization in the nephron tubular cells (Figure 3). Changes of clinical signs in this experiment have some similarities, although the degree of severity was observed in the positive control treatments. Clinical signs and histopathological sections were in line with the results of previous studies. Clinical signs such as abdominal ascites (edema), hemorrhage on skin, hemorrhage in gill filaments, swelling and rounded edges also found in the infected tilapia fish with several types of bacteria from the Enterobacteriaceae family (El-Barbary & Hal 2017). Behera et al (2017) further found that the challenged fish with *Acinetobacter baumannii* had loss of mucus and reddish (hemorrhage) lesion near the pectoral fin, however there was no clinical sign in the gills. The histology of experimentally challenged *Labeo rohita* showed hemorrhages and shrunken glomeruli with densely basophilic nuclei in the kidney. Other species have been studied by Rozi et al (2018), they infected *Osphronemus goramy* with five isolates of *A. hydrophila* from virulent and non-virulent strains, and they showed signs of ulcers, hemorrhagic on the base of the fins, body, mouth, and exophthalmia. The histopathology indicated spleen necrosis, picnosis, and inflammatory cells in the kidney.

Factors that play a role in bacterial pathogenicity were the propagation speed of pathogen and host defense against pathogen. Some bacterial extracellular products such as leucosidine and haemolysin were able to induce lysis of the blood cells, and then the bacteria spread throughout the host body to several target organs. Bacteria also have several types of enzymes in their extracellular products such as casein, gelatinase, amylase, lipase, chitinase, collagenase, elastase, hyaluronidase and proteinase that are

able to break down complex compounds into simpler compounds so that the bacteria can easily enter and damage the host cells (Pelcza & Chan 1986; Li et al 2012). The gill and kidney of *C. carpio* juveniles injected with several bacterial isolates of *O. melanopleurus* gut showed some damage and were thought to be caused by some extracellular products from these bacteria.

Bacteria that demonstrated an antagonistic ability to pathogenic bacteria in vitro test were not all safe or non-pathogenic to fish (Burbank et al 2012). In this pathogenicity assay the bacterial isolate BP_{S2} of *O. melanopleurus* gut bacteria was recommended due to its probiotic potential for further testing in *C. carpio* or other freshwater fish.

Conclusions. From the present study it was concluded that BP_{S2} bacterial isolate from *O. melanopleurus* gut showed the highest survival rate (93%) after being injected to *C. carpio* juveniles. This bacterium is safe and promising as potential-probiotic bacteria for common carp based on pathogenicity assay. Hematological parameters of *C. carpio* injected with BP_{S2} were better compared to the positive control and to other treatments. Histopathology of gills and kidney of experimental fish exhibited similar patterns.

16

Acknowledgements. The authors would like to thank to the Directorate General of Higher Education of the Republic of Indonesia (DIKTI) for The Doctoral Dissertation Research grant (No 042.06.1.401516/2018) for funding this research and to the Aquatic Microbiology Laboratory, Aquaculture Department, Faculty of Fisheries and Marine Sciences, Mulawarman University for facilitating the completion of the present research.

References

- Agustina, 2007 [Screening of probiotic bacteria to control *Aeromonas hydrophila* infection on catfish *Clarias* sp.]. Jurnal Aquacultura Indonesiana 8(3):135-143. [In Indonesian].
- Agustina, Saptiani G., Kusuma A. P., 2010 [Intestine bacterial selection of carp *Cyprinus carpio* against *Aeromonas hydrophila*]. Jurnal Badan Penelitian dan Pengembangan 'Daerah Gerbang Etam'. Kabupaten Kutai Kartanegara Kalimantan Timur 4(1). [In Indonesian].
- Agustina, Prayitno S. B., Sabdono A., Saptiani G., 2018 Antagonistic activity of Kelabau fish (*Osteochilus melanopleurus*) gut bacteria against *Aeromonas hydrophila* and *Pseudomonas* sp. AACL Bioflux 11(6):1859-1868.
- Anderson D. P., Siwicki A. K., 1995 Basic haematology and serology for fish health programs. In: Diseases in Asian aquaculture II. Fish health section. Shariff M., Arthur J. R., Subasinghe J. P. (eds), pp. 185-202, Asian Fisheries Society, Manila.
- Behera B. K., Paria P., Das A., Bhowmick S., Sahoo A. K., Das B. K., 2017 Molecular characterization and pathogenicity of a virulent *Acinetobacter baumannii* associated with mortality of farmed Indian Major Carp *Labeo rohita* (Hamilton 1822). Aquaculture 41:157-162.
- Blaxhall P. C., Daisley K. W., 1973 Routine haematological method for use with fish blood. Journal of Fish Biology 5:771-781.
- Burbank D. R., LaPatra S. E., Fornshell G., Cain K. D., 2012 Isolation of bacterial candidates from the gastrointestinal tract of rainbow trout, *Oncorhynchus mykiss* (Walbaum) and screening for inhibitory activity against *Flavobacterium psychrophilum*. Journal of Fish Diseases 35(11):809-816.
- Das A., Sahoo P. K., Mohanty B. R., Jena J. K., 2011 Pathophysiology of experimental *Aeromonas hydrophila* infection in *Puntius sarana*: Early changes in blood and aspects of the innate immune-related gene expression in survivors. Veterinary Immunology and Immunopathology 142:207-218.
- El-barbary M. I., Hal A. M., 2017 Molecular identification and pathogenicity of *Citrobacter* and *Serratia* species isolated from cultured *Oreochromis niloticus*. Egyptian Journal of Aquatic Research 43:255-263.

- Giri S. S., Sen S. S., Sukumaran V., 2012 Effects of dietary supplementation of potential probiotic *Pseudomonas aeruginosa* VSG-2 on the innate immunity and disease resistance of tropical freshwater fish, *Labeo rohita*. Fish Shellfish Immunology 32:1135-1140.
- Gupta A., Gupta P., Dhawan A., 2014 Dietary supplementation of probiotic affects growth, immune response and disease resistance of *Cyprinus carpio* fry. Fish & Shellfish Immunology 41:113-119.
- Haney D. C., Hursh D. A., Mix M. C., Winton J. R., 1992 Physiological and Hematological changes in chum salmon artificially infected with erythrocytic necrosis virus. Journal of Aquatic Animal Health 4:48-57.
- Li L., Dang Y., Shen Y., Xu X., Huang W., Li J., 2016 Hematological and immunological plasma assay for grass carp (*Ctenopharyngodon idella*) infected with *Aeromonas hydrophila* as an immune model in carp aquaculture. Fish & Shellfish Immunology 55:647-653.
- Li Y., Powell D. A., Shaffer S. A., Rasko D. A., Pelletier M. R., Leszyk J. D., Scott A. J., Masoudi A., Goodlett D. R., Wang X., Raetz C. R. H., Ernst R. K., 2012 LPS remodeling is an evolved survival strategy for bacteria. Proceedings of the National Academy of Sciences of the United States of America 109(22):8716-8721.
- Maslowski K. M., Mackay C. R., 2010 Diet, gut microbiota and immune responses. Nature Immunology 12:5-9.
- Ouwehand A. C., Salminen S., Isolauri E., 2002 Probiotic: a review of beneficial effects. Antonie van Leeuwenhoek 82:279-289.
- Pelcza M. J., Chan E. C. S., 1986 [Elements of microbiology]. Jakarta. UI Press. [In Indonesian].
- Reece J. B., Urry L. A., Cain M. L., Wasserman S. A., Minorsky P. V., Jacson R. B., 2014 Campbell biology. Pearson, New York, US, 1488 p.
- Roberts R. J., 2001 Fish pathology. W. B. Saunders, London, UK, 472 p.
- Rozi, Rahayu K., Daruti D. N., Stella M. S. P., 2018 Study on characterization, pathogenicity and histopathology of disease caused by *Aeromonas hydrophila* in gourami (*Osteogaster gourami*). IOP Conference Series. Earth and Environmental Science 137(1):012003.
- Shao J-z., Liu J., Xiang L-x., 2004 *Aeromonas hydrophila* induces apoptosis in *Carassius auratus* lymphocytes in vitro. Aquaculture 229:11-23.
- Thune R. L., Stanley L. A., Cooper R. K., 1993 Pathogenesis of gram-negative bacterial infections in warmwater fish. Annual Review of Fish Diseases 3:37-68.
- Wassom D. L., Kelly E. A. B., 1990 The role of the major histocompatibility complex in resistance to parasitic infection. Critical Reviews in Immunology 10:31-52.
- Wedemeyer G. A., Yasutake W. T., 1977 Clinical methods for the assessment of the effect on environmental stress on fish health. US Department of the Interior, Fish and Wildlife Service, Technical papers of the US Fish and Wildlife Service No. 89, 18 p.
- *** BPS-Statistics of Kalimantan Timur Province, 2017 [Kalimantan Timur Province in Figures]. Samarinda, 549 p. [In Indonesian].

Received: 17 September 2019. Accepted: 21 October 2019. Published online: 30 October 2019.

Authors:

Agustina, Mulawarman University, Faculty of Fisheries and Marine Science, Aquatic Microbiology Laboratory, Gunung Kelua Campus, Jl. Gunung Tabur, 75123 Samarinda, Indonesia, e-mail: agustina197748@gmail.com
Slamet Budi Prayitno, Diponegoro University, Faculty of Fisheries and Marine Science, Department of Aquaculture, Tembalang Campus, Jl. Prof. Sudarto SH, 50275 Semarang, Indonesia, e-mail: sbudiprayitno@gmail.com

Agus Sabdono, Diponegoro University, Faculty of Fisheries and Marine Science, Department of Marine Science, Tembalang Campus, Jl. Prof. Sudarto SH, 50275 Semarang, Indonesia, e-mail: agus_sabdono@yahoo.com

Gina Saptiani, Mulawarman University, Faculty of Fisheries and Marine Science, Aquatic Microbiology Laboratory, Gunung Kelua Campus, Jl. Gunung Tabur, 75123 Samarinda, Indonesia, e-mail: gina_saptiani@yahoo.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Agustina, Prayitno S. B., Sabdono A., Saptiani G., 2019 Pathogenicity assay of probiotic-potential bacteria from the Kelabau fish (*Osteochilus melanopleurus*). AACL Bioflux 12(5):1994-2003.

Pathogenicity assay of probiotic-potential bacteria from the Kelabau fish (*Osteochilus melanopleurus*)

ORIGINALITY REPORT

24%
SIMILARITY INDEX

15%
INTERNET SOURCES

19%
PUBLICATIONS

5%
STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | www.researchsquare.com
Internet Source | 2% |
| 2 | Submitted to Institut Pertanian Bogor
Student Paper | 2% |
| 3 | A. Das, P.K. Sahoo, B.R. Mohanty, J.K. Jena.
"Pathophysiology of experimental Aeromonas hydrophila infection in Puntius sarana: Early changes in blood and aspects of the innate immune-related gene expression in survivors", Veterinary Immunology and Immunopathology, 2011
Publication | 2% |
| 4 | Sukenda Sukenda, Rahman Rahman, Khairun Nisaa, Dendi Hidayatullah, Apriana Vinasiyam.
"The efficacy of Streptococcus agalactiae vaccine preparations, administered to tilapia broodstock, in preventing streptococcosis in their offspring, via transfer of maternal immunity", Aquaculture International, 2018
Publication | 2% |

- 5 Akhil Gupta, Paromita Gupta, Asha Dhawan. "Dietary supplementation of probiotics affects growth, immune response and disease resistance of Cyprinus carpio fry", Fish & Shellfish Immunology, 2014 2%
- Publication
-
- 6 Cheng Chi, Bing Jiang, Xiao-Bo Yu, Tian-Qiang Liu, Lei Xia, Gao-Xue Wang. "Effects of three strains of intestinal autochthonous bacteria and their extracellular products on the immune response and disease resistance of common carp, Cyprinus carpio", Fish & Shellfish Immunology, 2014 1%
- Publication
-
- 7 www.science.gov 1%
- Internet Source
-
- 8 www.cabi.org 1%
- Internet Source
-
- 9 Rozi, K Rahayu, D N Daruti. " Detection and analysis of hemolysin genes in isolated from Gouramy () by polymerase chain reaction (PCR) ", IOP Conference Series: Earth and Environmental Science, 2018 1%
- Publication
-
- 10 Khosravi, S., H.T.D. Bui, S. Rahimnejad, M. Herault, V. Fournier, J.B. Jeong, and K.-J. Lee. "Effect of dietary hydrolysate 1%

supplementation on growth performance, non-specific immune response and disease resistance of olive flounder (*Paralichthys olivaceus*) challenged with *Edwardsiella tarda*", Aquaculture Nutrition, 2014.

Publication

- | | | |
|----|---|--------|
| 11 | e-journal.unair.ac.id
Internet Source | 1 % |
| 12 | Manal I. El-Barbary, Ahmed M. Hal. "Molecular identification and pathogenicity of <i>Citrobacter</i> and <i>Serratia</i> species isolated from cultured <i>Oreochromis niloticus</i> ", The Egyptian Journal of Aquatic Research, 2017
Publication | 1 % |
| 13 | pui-oktal.unmul.ac.id
Internet Source | 1 % |
| 14 | G. Deepika, P. Sivakumar, M.R Rajan. "Isolation and Characterization of Bacteria from the Gut of Blue Gourami (<i>Trichogaster tricopterus</i>) and its Role on Growth", Journal of Pure and Applied Microbiology, 2019
Publication | <1 % |
| 15 | journals.lww.com
Internet Source | <1 % |
| 16 | insightsociety.org
Internet Source | <1 % |

- 17 Dash, Gyanaranjan, Ram Prakash Raman, K. Pani Prasad, M. Makesh, M.A. Pradeep, and Swatipriyanka Sen. "Evaluation of *Lactobacillus plantarum* as feed supplement on host associated microflora, growth, feed efficiency, carcass biochemical composition and immune response of giant freshwater prawn, *Macrobrachium rosenbergii* (de Man, 1879)", *Aquaculture*, 2014.
Publication <1 %
- 18 Submitted to Federal University of Technology <1 %
Student Paper
- 19 Gina Saptiani, Slamet Budi Prayitno, Sari Anggarawati. "Effect of mangrove leaf extract (*Acanthus ilicifolius*) on non-specific immune status and vibriosis resistance of black tiger shrimps (*Penaeus monodon*) challenged with *Vibrio harveyi*", *Veterinary World*, 2021
Publication <1 %
- 20 eprints.undip.ac.id <1 %
Internet Source
- 21 Submitted to Program Pascasarjana Universitas Negeri Yogyakarta <1 %
Student Paper
- 22 EH Hardi, G Saptiani, IW Kusuma, W Suwinarti, A Sudaryono. "Inhibition of fish bacteria pathogen in tilapia using a concoction three <1 %

of Borneo plant extracts", IOP Conference
Series: Earth and Environmental Science, 2018

Publication

- 23 www.golfalencon.fr <1 %
Internet Source
-
- 24 www.jafs.com.pl <1 %
Internet Source
-
- 25 E Omorogie. "Utilization and nutrient digestibility of mango seeds and palm kernel meal by juvenile *Labeo senegalensis* (Antheriniformes: Cyprinidae)", Aquaculture Research, 9/2001 <1 %
Publication
-
- 26 epdf.pub <1 %
Internet Source
-
- 27 pinnacle.allenpress.com <1 %
Internet Source
-
- 28 www.jparonline.com <1 %
Internet Source
-
- 29 Giri, S.S., V. Sukumaran, S.S. Sen, and P.K. Jena. "Effects of dietary supplementation of potential probiotic *Bacillus subtilis* VSG1 singularly or in combination with *Lactobacillus plantarum* VSG3 or/and *Pseudomonas aeruginosa* VSG2 on the growth, immunity <1 %

and disease resistance of *Labeo rohita*",
Aquaculture Nutrition, 2013.

Publication

-
- 30 eprints.cmfri.org.in <1 %
Internet Source
-
- 31 projects.sare.org <1 %
Internet Source
-
- 32 Phromkunthong, Wutiporn, Nutt Nuntapong, Sunee Wanlem, and Mali Boonyaratpalin. "A study on growth, histopathology and oxidative stress in Asian sea bass on diets with various loadings of melamine and cyanuric acid adulterants", Aquaculture, 2015.
Publication
-
- 33 coek.info <1 %
Internet Source
-
- 34 eprints.untirta.ac.id <1 %
Internet Source
-
- 35 jvr.ut.ac.ir <1 %
Internet Source
-
- 36 suaire.suanet.ac.tz <1 %
Internet Source
-
- 37 www.ijsr.net <1 %
Internet Source
-
- 38 www.ros.hw.ac.uk <1 %
Internet Source

39

"Aquaculture Nutrition", Wiley, 2014

Publication

<1 %

40

Gobi, Narayanan, Balasubramanian
Malaikozhundan, Vijayakumar Sekar,
Sathappan Shanthi, Baskaralingam
Vaseeharan, Rengarajan Jayakumar, and
Abdul Khudus Nazar. "GFP tagged Vibrio
parahaemolyticus Dahv2 infection and the
protective effects of the probiotic Bacillus
licheniformis Dahb1 on the growth, immune
and antioxidant responses in Pangasius
hypophthalmus", Fish & Shellfish
Immunology, 2016.

Publication

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On

Pathogenicity assay of probiotic-potential bacteria from the Kelabau fish (*Osteochilus melanopleurus*)

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
