

Check 8

by Sifri Sifri

Submission date: 08-May-2023 10:19PM (UTC+1000)

Submission ID: 2087492427

File name: -_Determining_-_Faisal_2021_J._Phys._Conf._Ser._1751_012028.pdf (1.03M)

Word count: 2515

Character count: 11376

PAPER • OPEN ACCESS

Determining the Noetherian Property of Generalized Power Series Modules by Using X-Sub-Exact Sequence

To cite this article: A Faisol *et al* 2021 *J. Phys.: Conf. Ser.* **1751** 012028

View the [article online](#) for updates and enhancements.

You may also like

- [Lie algebras of triangular polynomial derivations and an isomorphism criterion for their Lie factor algebras](#)
V. V. Bavula

- [Regular subcategories in bounded derived categories of affine schemes](#)
A. D. Elagin and V. A. Lunts

- [PROJECTIVE MODULES OVER QUANTUM POLYNOMIAL ALGEBRAS](#)
V A Artamonov



The advertisement features a dark blue background on the left with the ECS logo and text, and a photograph of a woman at a laptop on the right. A blue button with white text is overlaid on the image.

ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

243rd Meeting with SOFC-XVIII
Boston, MA • May 28 – June 2, 2023
Accelerate scientific discovery!

Learn More & Register

Determining the Noetherian Property of Generalized Power Series Modules by Using X -Sub-Exact Sequence

A Faisol¹, Fitriani², and Sifriyani³

^{1,2}Department of Mathematics, Universitas Lampung, Bandar Lampung, Indonesia

³Department of Mathematics, Universitas Mulawarman, Samarinda, Indonesia

email: ahmadfaisol@fmipa.unila.ac.id¹, fitriani.1984@fmipa.unila.ac.id², sifriyani@fmipa.unmul.ac.id³

Abstract. The Noetherian property of the generalized power series module can determine in several ways. This paper uses the sub-exact sequence of modules over a ring R to determine this property. This investigation not only determines the Noetherian property of the generalized power series module but also the Noetherian property of its submodule. Furthermore, we give a construction of $R[[S]]$ -homomorphism between the generalized power series modules.

Keyword: noetherian, strictly ordered monoid, generalized power series modules, exact sequence, sub-exact sequence.

1. Introduction

The exact sequence of modules is one of the essential concepts in module theory [1], [2]. In [3], Fitriani et al. introduced a sub-exact sequence of modules. This concept is motivated by the quasi exact sequence established by Davvaz and Parnian-Garamaleky [4]. Furthermore, they use this concept to generalize the generator of modules related to a family of modules over a ring R [5]. Moreover, using a generalization of a linearly independent set of modules [6], they obtained a basis and free modules related to a family of modules [7].

Given ring R , monoid (S, \leq) with a strictly ordered, and a monoid homomorphism ω from S to $\text{End}(R)$. In 2019, Faisol and Fitriani gave some conditions for skew GPSM to be a $T[[S, \omega]]$ -Noetherian module over a ring $R[[S, \omega]]$ [8]. This sufficient condition is a generalization of the previous results [9], which were obtained by applying the properties in [10], generalizing the sufficient conditions in [11], and using the relations specified in [12].

Varadarajan [13] introduce the generalized power series module (GPSM). This module is a module over the generalized power series ring (we call it by GPSR), introduced by Ribenboim [14]. Moreover, the results of Ribenboim construction were generalized by Mazurek and Ziemkowski [15] by utilizing the monoid homomorphism used in the convolution multiplication operation. In addition to constructing GPSM, Varadarajan [16] also provides necessary and sufficient conditions that GPSM is a Noetherian module. In this paper, we give a method to determine the Noetherian property of the generalized power series module. We use the concept of the sub-exact sequence to determine this property. In this way, we also can determine the Noetherian property of its submodules. Moreover, we give a construction of $R[[S]]$ -homomorphism between the generalized power series modules.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

2. The Main Results

Let R be a commutative ring with $1_R \in R$ and S be a monoid with strictly ordered. Let $N_1, N_2,$ and N_3 be three modules over ring R . The set $N_i[[S]]$ consists of all function μ from S to N_i such that the support of f is Artinian and narrow (we denote support of f by $\text{supp}(f)$, that is the set of $s \in S$, where $f(s)$ is not equal to 0), for $i = 1, 2, 3$. We can write the set as follow:

$$N_i[[S]] = \{\mu : S \rightarrow N_i \mid \text{supp}(\mu) \text{ is Artinian and narrow}\},$$

$i = 1, 2, 3$.

Before we give a condition when a submodule $L[[S]]$ of $N_2[[S]]$ is Noetherian $R[[S]]$ -module, we recall that if L is a submodule of N , then $L[[S]]$ is a submodule of $N_2[[S]]$ as a module over $R[[S]]$. Let

$$L[[S]] = \{\mu \in N_2[[S]] \mid \mu(s) \in L, \text{ for all } s \in S\}.$$

The set $L[[S]]$ is a submodule of $N_2[[S]]$.

Let K, L, M be R -modules and X be R -submodules of L . Recall that the triple (K, L, M) is said to be X -sub-exact at L if there exist f and g such that the sequence $K \xrightarrow{f} X \xrightarrow{g} M$ is exact. In the following proposition, we give a condition when a submodule $L[[S]]$ of $N_2[[S]]$ is Noetherian.

Proposition 1. Let R be a commutative ring with $1 \in R$ and (S, \leq) be a monoid with a strictly ordered. Let $N_1, N_2,$ and N_3 are R -modules, and L is a submodule of N_2 over R . If the triple $(N_1[[S]], N_2[[S]], N_3[[S]])$ is $L[[S]]$ -sub-exact as an $R[[S]]$ -module, $N_1[[S]]$ and $N_3[[S]]$ are Noetherian $R[[S]]$ -modules, then $L[[S]]$ is a Noetherian $R[[S]]$ -module.

Proof. Since the triple $(N_1[[S]], N_2[[S]], N_3[[S]])$ is $L[[S]]$ -sub-exact, based on [3], we have the following sequence of a module over $R[[S]]$ is exact.

$$N_1[[S]] \rightarrow L[[S]] \rightarrow N_3[[S]] \tag{1}$$

Since (1) is exact, there are $R[[S]]$ -homomorphism f and g , where f is an $R[[S]]$ -homomorphism from $N_1[[S]]$ to $L[[S]]$, g is an $R[[S]]$ -homomorphism from $L[[S]]$ to $N_3[[S]]$, and $\text{Im}(f) = \text{Ker}(g)$. By hypothesis, $N_1[[S]]$ and $N_3[[S]]$ are Noetherian modules over $R[[S]]$. Hence based on [17], we have $N[[S]]$ is a Noetherian as a module over $R[[S]]$.

Given three R -modules $N_1, N_2,$ and N_3 . Fitriani et al. [3] construct a set $\sigma(N_1, N_2, N_3)$ that consists of all submodules X of N_2 such that the triple (N_1, N_2, N_3) is an X -sub-exact at N_2 , i.e.:

$$\sigma(N_1, N_2, N_3) = \{X \text{ submodule of } N_2 \mid (N_1, N_2, N_3) \text{ is an } X\text{-sub exact at } N_2\}.$$

In this case, we construct the set $\sigma(N_1[[S]], N_2[[S]], N_3[[S]])$ that consist of all submodules X of $N_2[[S]]$ such that the triple of generalized power series modules $(N_1[[S]], N_2[[S]], N_3[[S]])$ is an X -sub exact at $N_2[[S]]$, i.e.: $\sigma(N_1[[S]], N_2[[S]], N_3[[S]]) = \{X \leq N_2[[S]] \mid \text{the triple } (N_1[[S]], N_2[[S]], N_3[[S]]) \text{ is an } X\text{-sub exact at } N_2[[S]]\}$.

As a direct consequence of Proposition 1, we have the following result.

Corollary 1. Let R be a commutative ring with $1 \in R$ and (S, \leq) be a monoid with strictly ordered. Let $M_1, M_2,$ and M_3 are modules over ring R . If $N_1[[S]]$ and $N_3[[S]]$ are Noetherian modules over $R[[S]]$, then a submodule X of N_2 is Noetherian, for every $X \in \sigma(N_1[[S]], N_2[[S]], N_3[[S]])$.

Proof. Let $X \in \sigma(N_1[[S]], N_2[[S]], N_3[[S]])$. We have the following exact sequence of $R[[S]]$ -modules:

$$N_1[[S]] \rightarrow X \rightarrow N_3[[S]]$$

From Proposition 1, we have X is Noether.

In [18], Ziembowski gives a construction of a homomorphism of skew GPSR. Based on his construction, we construct a homomorphism of generalized power series modules in the following proposition.

Proposition 2. Given a commutative ring R with identity element 1. Given a monoid (S, \leq) with a strictly ordered, an endomorphism ω of S such that for every subset Artinian and narrow $T \subseteq S$, $\omega(T)$

is Artinian, narrow, and $h(\omega^{-1}(x)) = h(x)$, for every x is in S , and h is in $R[[S]]$. Let φ be an R -homomorphism from N_2 to N_3 , where N_2, N_3 be R -modules. For $\mu \in N_2[[S]]$, we define:

$$\begin{aligned} \phi: N_2[[S]] &\rightarrow N_3[[S]] \\ \mu &\mapsto \bar{\mu}, \end{aligned}$$

where

$$\bar{\mu}(x) = \begin{cases} \varphi \circ \mu \circ \omega^{-1}(x) & ; \text{if } x \in \omega(S), \\ 0 & ; \text{otherwise.} \end{cases} \dots\dots\dots (1)$$

Then ϕ is an $R[[S]]$ -homomorphism from $N_2[[S]]$ to $N_3[[S]]$.

Proof. Since $\text{supp}(\bar{\mu}) \subseteq \omega(\text{supp}(\mu))$, we have $\bar{\mu} \in N_3[[S]]$. Now, we will show that ϕ is a $R[[S]]$ -homomorphism from $N_2[[S]]$ to $N_3[[S]]$.

a. Let $\mu, \beta \in N_2[[S]]$, and $x \in S$. By (1), we have:

$$\begin{aligned} \overline{\mu + \beta}(x) &= \varphi \circ (\mu + \beta) \circ \omega^{-1}(x) \\ &= \varphi((\mu + \beta) \circ \omega^{-1}(x)) \\ &= \varphi(\mu(\omega^{-1}(x)) + \beta(\omega^{-1}(x))) \\ &= \varphi(\mu(\omega^{-1}(x))) + \varphi(\beta(\omega^{-1}(x))) \\ &= \varphi \circ \mu \circ \omega^{-1}(x) + \varphi \circ \beta \circ \omega^{-1}(x) \\ &= \bar{\mu}(x) + \bar{\beta}(x). \end{aligned}$$

This equation implies that $\overline{\mu + \beta} = \bar{\mu} + \bar{\beta}$, and hence $\phi(\mu + \beta) = \phi(\mu) + \phi(\beta)$, for every $\mu, \beta \in N_2[[S]]$.

b. Let $\mu \in N_2[[S]]$, $h \in R[[S]]$, and $x \in S$. By (1), we get:

$$\begin{aligned} \overline{h\mu}(x) &= \varphi \circ (h\mu) \circ \omega^{-1}(x) \\ &= \varphi(h\mu(\omega^{-1}(x))) \\ &= \varphi(\sum_{s+t=\omega^{-1}(x)} h(s)\mu(t)) \\ &= \sum_{s+t=\omega^{-1}(x)} \varphi(h(s)\mu(t)) \\ &= \sum_{s+t=\omega^{-1}(x)} h(s)\varphi(\mu(t)) \\ &= \sum_{\omega^{-1}(u)+\omega^{-1}(v)=\omega^{-1}(x)} h(\omega^{-1}(u))\varphi(\mu(\omega^{-1}(v))) ; s = \omega^{-1}(u) \text{ dan } t = \omega^{-1}(v) \\ &= \sum_{\omega^{-1}(u)+\omega^{-1}(v)=\omega^{-1}(x)} h(u)\varphi(\mu(\omega^{-1}(v))) ; h(\omega^{-1}(u)) = h(u) \\ &\quad \omega^{-1}(u+v)=\omega^{-1}(x) \\ &= \sum_{\substack{u+v=x \\ u+v=x}} h(u)(\varphi \circ \mu \circ \omega^{-1})(v) \\ &= \sum_{u+v=x} h(u)\bar{\mu}(v) \\ &= h\bar{\mu}(x). \end{aligned}$$

Hence, for every $\mu \in N_2[[S]]$, $h \in R[[S]]$, we have $\phi(h\mu) = \overline{h\mu} = h\bar{\mu} = h\phi(\mu)$.

From a-b, we can conclude that ϕ is an $R[[S]]$ -homomorphism from $N_2[[S]]$ to $N_3[[S]]$.

Given an R -module M , we recall that a submodule N of M is a direct summand of M if there exists $K \leq M$ such that $M = N \oplus K$, i.e., $M = N + K$, and $N \cap K = 0$. In this case, every $m \in M$ can be uniquely written as $m = a + b$, where $a \in N$, and $b \in K$ [17]. Next, we will use the construction of $R[[S]]$ -homomorphism in Proposition 2 to provide the Noetherian property of the GPSM.

Proposition 3. Given a commutative ring R with $1 \in R$ and a monoid (S, \leq) with a strictly ordered. Let N_1, N_2 , and N_3 are R -modules, and $L[[S]]$ is a direct summand of $N_2[[S]]$ as an $R[[S]]$ -module. If $(N_1[[S]], N_2[[S]], N_3[[S]])$ is $L[[S]]$ -sub-exact as an $R[[S]]$ -module, $N_1[[S]]$ and $N_3[[S]]$ are Noetherian $R[[S]]$ -modules, then $N_2[[S]]$ is Noether.

Proof. By hypothesis $(N_1[[S]], N_2[[S]], N_3[[S]])$ is $L[[S]]$ -sub-exact as an $R[[S]]$ -module. Since $N_1[[S]]$ and $N_3[[S]]$ are Noetherian $R[[S]]$ -modules, based on Proposition 1, we get $L[[S]]$ is Noether. Since $L[[S]]$ is a direct summand, there exists a submodule K of $N_2[[S]]$ such that $N_2[[S]] = L[[S]] \oplus K$. Then every $\mu \in N_2[[S]]$ can uniquely write as $\mu = \mu' + \mu''$, where $\mu' \in L[[S]]$, and $\mu'' \in K$. Besides that, the triple $(N_1[[S]], N_2[[S]], N_3[[S]])$ is $L[[S]]$ -sub-exact implies that there are two $R[[S]]$ -homomorphisms f and g such that the following sequence is exact.

$$N_1[[S]] \xrightarrow{f} L[[S]] \xrightarrow{g} N_3[[S]],$$

i.e., $\text{Im}(f) = \text{Ker}(g)$.

Thus, we can define an $R[[S]]$ -homomorphism

$$g': N_2[[S]] \rightarrow N_3[[S]],$$

where $g' = \begin{cases} g(\mu); & \text{if } \mu \in L[[S]]; \\ 0 & ; \text{ otherwise.} \end{cases}$

Hence, we get the following diagram of $R[[S]]$ -module:

$$\begin{array}{ccccc} N_1[[S]] & \xrightarrow{f} & L[[S]] & \xrightarrow{g} & N_3[[S]] \\ & \searrow & \downarrow i & \nearrow & \\ & i \circ f & N_2[[S]] & g' & \end{array}$$

Based on [3], the following sequence of $R[[S]]$ -module is exact.

$$N_1[[S]] \xrightarrow{i \circ f} N_2[[S]] \xrightarrow{g'} N_3[[S]].$$

Since $N_1[[S]]$ and $N_3[[S]]$ are Noetherian, based on Proposition 1, $N_2[[S]]$ is Noetherian.

Conclusion

Based on the results, we can conclude that we can use the concept of a sub-exact sequence of modules over $R[[S]]$ to determine the Noetherian property of generalized power series modules. Besides that, we also can determine the Noetherian property of its submodule.

Acknowledgments

The authors wish to thank the Research Institutions and Community Service of Universitas Lampung for this research's support and funding under the Research Contract No: 1491/UN 26.21/PN/2020.

References

[1] Adkins W A and Weintraub S H 1992 *Algebra: An Approach via Module Theory* (New York: Springer-Verlag)

[2] Dummit S and Foote R 2004 *Abstract Algebra* (USA: John Wiley and Sons, Inc.)

[3] Fitriani, Surodjo B and Wijayanti I E 2016 On sub-exact sequences *Far East J. Math. Sci.* **100**(7) 1055

[4] Davvaz B and Pamian-Garamaleky P 1999 A Note on Exact Sequences *Bull. Malaysian Math. Sci. Soc.* **22** 53

[5] Fitriani, Wijayanti I E and Surodjo B 2018 Generalization of U-Generator and M-Subgenerator Related to Category $\sigma[M]$ *J. Math. Res.* **10**(4) 101

[6] Fitriani, Surodjo B and Wijayanti I E 2017 On X-Sub-Linearly Independent Modules *Journal of Physics: Conference Series* **893**(012008) 1

[7] Fitriani, Wijayanti I E and Surodjo B 2018 A Generalization of Basis and Free Modules Relatives to a Family of R-modules *Journal of Physics: Conference Series* **1097**(012087) 1

[8] Faisol A and Fitriani 2019 The Sufficient Conditions for Skew Generalized Power Series Module $M[[S, \omega]]$ to be $T[[S, \omega]]$ -Noetherian $R[[S, \omega]]$ -module *Al-Jabar J. Pendidik. Mat.* **10**(2) 285

- [9] Faisal A, Surodjo B and Wahyuni S 2019 T[[S]]-Noetherian Property On Generalized Power Series Modules *JP J. Algebr. Number Theory Appl.* **43**(1) 1
- [10] Faisal A, Surodjo B and Wahyuni S 2018 The Impact Of The Monoid Homomorphism On The Structure Of Skew Generalized Power Series Rings *Far East J. Math. Sci.* **103**(7) 1215
- [11] Faisal A, Surodjo B and Wahyuni S 2019 The Sufficient Conditions for $R[X]$ -module $M[X]$ to be $S[X]$ -Noetherian *Eur. J. Math. Sci.* **5**(1) 1
- [12] Faisal A, Surodjo B and Wahyuni S 2019 The Relation between Almost Noetherian Module, Almost Finitely Generated Module and T-Noetherian Module *Journal of Physics: Conference Series* **1306**(01200) 1
- [13] Varadarajan K 2001 Noetherian Generalized Power Series Rings and Modules *Commun. Algebr.* **29**(1) 245
- [14] Ribenboim P 1992 Noetherian Rings Of Generalized Power Series *J. Pure Appl. Algebr.* **79**(3) 293
- [15] Mazurek R and Ziemkowski M 2008 On Von Neumann Regular Rings Of Skew Generalized Power Series *Commun. Algebr.* **36**(5) 1855
- [16] Varadarajan K 2001 Generalized Power Series Modules *Commun. Algebr.* **29**(3) 1281
- [17] Wisbauer R 1991 *Foundations of Module and Ring Theory* (USA: Philadelphia)
- [18] Ziemkowski M 2010 *Right Gaussian Rings and Related Topics* (University of Edinburgh)

Check 8

ORIGINALITY REPORT

7%

SIMILARITY INDEX

7%

INTERNET SOURCES

2%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

teknik.trunojoyo.ac.id

Internet Source

3%

2

ejournal2.undip.ac.id

Internet Source

2%

3

[Submitted to Binus University International](#)

Student Paper

2%

Exclude quotes On

Exclude bibliography On

Exclude matches < 2%

Check 8

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
