

Evaluation of non-timber forest products used as medicinal plants from East Kalimantan (Indonesia) to inhibit α -glucosidase and free radicals

by Rudianto Amirta

Submission date: 05-May-2023 01:56AM (UTC+0700)

Submission ID: 2084332994

File name: limantan_Indonesia_to_inhibit_-glucosidase_and_free_radicals.pdf (1,016.99K)

Word count: 5821

Character count: 32582

Evaluation of non-timber forest products used as medicinal plants from East Kalimantan (Indonesia) to inhibit α -glucosidase and free radicals

RICO RAMADHAN^{1,2,*}, RAMADHAN TOSEPU³, PREECHA PHUWAPRAISRISAN⁴, RUDIANTO AMIRTA⁵, KIETTIPUM PHONTREE⁶, YAZDANIYAR FAJRI HALIMI FIRDAUS², NURLITA ABDULGANI⁷, MUHAMMAD ZAINUL MUTTAQIN⁸, SAPARWADI⁹

¹Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115, East Java, Indonesia. Tel/fax.: +62-31-5936501, *email: rico.ramadhan@fst.unair.ac.id

²Division of Exploration and Synthesis of Bioactive Compounds, University CoE-Research Center for Bio-Molecule Engineering, Universitas Airlangga, Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115, East Java, Indonesia

³Faculty of Public Health, Universitas Halu Oleo, Jl. H.E.A. Mokodompit, Kendari 93232, Sulawesi Tenggara, Indonesia

⁴Center of Excellent in Natural Products, Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

⁵Faculty of Forestry, Universitas Mulawarman, Jl. Gunung Kelua, Samarinda 75123, East Kalimantan, Indonesia

⁶Division of Pharmacognosy and Toxicology, Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen 40002, Thailand

⁷Department of Biology, Faculty of Science and Data Analytics, Institut Teknologi Sepuluh Nopember, Kampus ITS Sukolilo, Surabaya 60111, Indonesia

⁸Generasi Biologi Indonesia (Genbinesia) Foundation, Jl. Swadaya Barat No. 4, Semampir, Gresik 61171, East Java, Indonesia

⁹BIOMA Foundation, Jl. A Wahab Syahrani, Sempaja, Samarinda 75119, East Kalimantan, Indonesia

Manuscript received: 9 July 2022. Revision accepted: 25 October 2022.

Abstract. Ramadhan R, Tosepu R, Phuwapraisrisan P, Amirta R, Phontree K, Firdaus YFH, Abdulgani N, Muttaqin MZ, Saparwadi. 2022. Evaluation of non-timber forest products used as medicinal plants from East Kalimantan (Indonesia) to inhibit α -glucosidase and free radicals. *Biodiversitas* 23: 5551-5558. Non-timber forest products (NTFPs) have essential uses as medicines, food sources, and traditional ceremonies for the local people living in forest areas. Medicinal plants are one of the non-timber forest products used as traditional medicine by local people in East Kalimantan. This study was conducted to search for new natural antidiabetic and antioxidant resources from medicinal plants used traditionally in Mahakam Ulu District, East Kalimantan. Based on ethnopharmacological information, the present study evaluates non-timber forest products as in vitro antidiabetic (rat intestinal α -glucosidase inhibitory activity) and antioxidant (free radical scavenging activity against DPPH, ABTS, and Nitric Oxide) activities. The antidiabetic and antioxidant activities were evaluated using α -glucosidase inhibition assay and free radical scavenging methods. This study demonstrated that methanol extracts of *Flacourtia rukam* Zoll. & Moritz, *Syzygium chloranthum* (Duthie) Merr. & L.M. Perry and *Shorea ulangeran* Burck exhibited strong α -glucosidase inhibitory activity with maltose as a substrate with IC₅₀ values of 0.034 mg/mL, 0.039 mg/mL, and 0.039 mg/mL, respectively. On the other hand, methanol extract of *S. chloranthum* displayed the strongest α -glucosidase inhibitory activity in sucrose as a substrate, similar to the IC₅₀ value of the quercetin as a positive control. Furthermore, the antioxidant test showed that all medicinal plant extracts from East Kalimantan are good sources of natural antioxidants indicated by their IC₅₀ values. The results of this study support the scientific background of the uses of medicinal plant extracts from East Kalimantan as folk medicine.

Keywords: Antidiabetics, antioxidant, biodiversity, medicinal plants, tropical regions

INTRODUCTION

Non-timber forest products (NTFPs) are products other than timber produced in the forests, such as medicinal plants, honey, mushrooms, resins, fruit and nuts, vegetables, and various barks and natural fibers (Mipun et al. 2019). NTFPs have an important role in supporting the livelihood of local communities around forest areas (Shackleton et al. 2018). Medicinal plants as non-timber forest products (NTFPs) have been used to treat several diseases since ancient times (Mbopi et al. 2021) for therapeutic purposes. Medicinal plants are well-known storehouses of various bioactive secondary metabolites, such as phenolics, flavonoids, alkaloids, terpenes, and tannins, that are responsible for their therapeutic activity, such as anti-diabetic, antioxidant, anti-inflammation, wound-healing, and antibacterial (Khan et al. 2022; Moukette et al. 2015). East Kalimantan has vast tropical

forest biodiversity, which includes medicinal plants that Dayak people traditionally used for generations. Several medicinal plants growing in the tropical forests of East Kalimantan have been reported to have biological activities, including antioxidant, antidiabetic, antibacterial, anti-tyrosinase, and anticancer activities (Arung et al. 2012; Sukara et al. 2014; Kusuma et al. 2014; Abuga et al. 2022). Regarding the search for the most potent traditional medicinal plants and phytotherapy, scientific approaches should be conducted to develop new effective, inexpensive, and nontoxic traditional medications.

Diabetes mellitus is a chronic disease characterized by hyperglycemia, primarily due to disproportionate insulin demand. Hyperglycemia causes numerous complications, including diabetic foot ulcers, heart disease, kidney failure, retinopathy, stroke and pregnancy-related complications (Ji et al. 2021). The pathogenesis of type 2 diabetes mellitus is a complex metabolism involving the

overproduction of free radicals and oxidative stress, which contributes to previously mentioned complications (Liu et al. 2014). In curing diabetes, various strategies have been developed to lower blood sugar levels, such as insulin secretagogues, an insulin sensitizer, inhibitors of glucose recapture, and anti-hyperglycemia. Along with insulin treatment, previous studies have shown that inhibiting α -glucosidase is an effective strategy to control the development of diabetes via regulated postprandial blood glucose (Trinh et al. 2016). α -Glucosidase inhibitors such as acarbose, miglitol, and l-deoxynojirimycin are well-known digestive enzyme inhibitors as synthetic drugs for type 2 diabetic patients. Although these drugs can reduce postprandial hyperglycemia and prevent impaired glucose intolerance, they still have side effects related to gastrointestinal problems (Lee et al. 2020). Fortunately, active secondary metabolites in medicinal plants have the potency to treat hyperglycemia and its complications by scavenging free radicals (Akyuz et al. 2022).

In this study, ten selected medicinal plants from Mahakam Ulu were traditionally used for blood sugar level therapy by local communities in Batu Majang Village. This study area has never been ethnobotanically explored; therefore, the present study is considered the first study on the ethnobotanical medicinal plants in this region. This study aims to evaluate the activity of α -glucosidase inhibition and free radical scavenging activity of medicinal plants from East Kalimantan. Therefore, it might be more beneficial than common antidiabetic drugs. To the best of the authors' knowledge, this study is the first report on antidiabetic and antioxidant activity screening of selected medicinal plants from the East Kalimantan tropical forest.

MATERIALS AND METHODS

Study site and plant collections

The study was conducted in Batu Majang Village, Long Bagun Sub-district, Mahakam Ulu District, Indonesia (Figure 1). The study site was chosen because the traditional local communities in this area use traditional medicinal plants to treat diseases, including blood sugar levels. Selected plant parts were collected, and the voucher specimens of ten selected medicinal plants were stored in the Laboratory of Forest Products Chemistry and Renewable Energy, Faculty of Forestry, Universitas Mulawarman. In addition, ten medicinal plants were identified at the Laboratory of Forest Dendrology, Faculty of Forestry, Universitas Mulawarman, Samarinda, Indonesia. These specimens are listed in Table 1.

Chemical reagents and instrument

ABTS (2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) and DPPH (1,1-diphenyl-2-picrylhydrazyl) were procured from Tokyo Chemical Industry Co. Ltd. Potassium persulfate, ascorbic acid, naphthyl ethylenediamine dihydrochloride, sodium nitroprusside, and potassium perchlorate were purchased from Sigma Chemical Co (St. Louis, MO, USA). Rat intestinal acetone powder (Sigma Aldrich) and all other chemicals and solvents were of the highest commercial grade. Absorbance was taken using a microplate reader BIOCROM EZ Read 2000 (BioChrom Ltd, United Kingdom).




Figure 1. Study site in Batu Majang Village (red), Mahakam Ulu District, East Kalimantan, Indonesia

Table 1. Selected medicinal plants from East Kalimantan

Medicinal plants	Part used	Local names	Voucher specimens
<i>Shorea balangeran</i> Burck	Branch & Twig	Kahoi	LBTI-R-30
<i>Syzygium chloranthum</i> (Duthie) Merr. & L.M.Perry	Branch & Twig	Bumbun	LBTI-R-40
<i>Lagerstroemia speciosa</i> (L.) Pers.	Stem	Bungur	LBTI-T-22
<i>Flacourtia rukam</i> Zoll. & Moritz	Stem	Rukam	LBTI-P-08
<i>Gluta velutina</i> Blume	Stem	Serengat	LBTI-S-09
<i>Kleinhovia hospita</i> L.	Branch & Twig	Tahongai	LBTI-R-05
<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	Stem	Sengon Buto	LBTI-S-23
<i>Vatica umbonata</i> Burck	Branch & Twig	Masinstan	LBTI-S-27
<i>Pertusadina eurhyncha</i> (Miq.) Ridsdale	Stem	Kadamba	LBTI-P-33
<i>Dillenia excelsa</i> (Jack) Martelli ex Gilg.	Stem	Tanekara	LBTI-S-14

Extraction of medicinal plants

Plant stems, twigs, and branches were washed with distilled water and dried at room temperature in a well-ventilated room. The dried plant materials were ground into a fine powder using a grinder, then separately extracted in methanol (3 x 400 mL) and macerated at ambient temperature. The mixture was then filtered through filter papers. Next, the filtrate was dried under a vacuum in a rotary evaporator (Rotavapor R100, BUCHI) to obtain dried crude extracts, which were then transferred into amber bottles. After that, crude extracts were stored at 4°C for further bioassay analysis.

30 *in vitro* antidiabetic activity

Rat intestinal α -glucosidase inhibition assay

Rat intestinal α -glucosidase inhibition test *in vitro* following the methods by Ramadhan et al. (2020). In brief, 50 μ L of phosphate buffer (pH 6.9) was mixed with crude enzyme solution, 10 μ L of the sample various concentrations, and a glucose kit before they were added into a 96-well plate and incubated for 10 minutes (maltose), and 40 minutes (sucrose) at 37°C, respectively. After incubation, the absorbance was recorded using a microplate reader BIOCROM EZ Read 2000. The percentage of inhibition (%) was calculated with the following formula:

$$\% \text{ Inhibition} = (\text{Abs control} - \text{Abs sample}) / \text{Abs control} \times 100.$$

IC₅₀ is the concentration required to inhibit 50% of rat intestinal α -glucosidase activity. Quercetin was used as a positive control.

24 Antioxidant properties

DPPH radical scavenging activity

The ability of selected plant extracts to scavenge free radical DPPH was determined based on Khongkara et al. (2020) methods with minor modification. First, the DPPH solution was prepared by dissolving free radical DPPH powder in 100 mL of methanol to obtain a concentration of 0.10 mM. Next, the 0.10 mM DPPH solution was added to 20 μ L of extracts of various concentrations sequentially and then incubated for 30 min in the darkness. After incubation, the absorbance was measured at 517 nm. Finally, the free radical scavenging activity of samples was expressed as a percentage of inhibition (PI) using the following equation (1):

$$\% \text{ Scavenging activity} = (\text{Abs control} - \text{Abs sample}) / \text{Abs control} \times 100.$$

The concentration of samples required to inhibit 50% free radical DPPH is IC₅₀. The test was performed in triplicate for each sample concentration, and ascorbic acid was used as a positive control.

ABTS radical scavenging activity

The ABTS radical scavenging activity assay was carried out using the free radical ABTS method employed by Seyrekoglu et al. (2022) with a slight modification. The free radical ABTS solution was prepared by mixing equal quantities of ABTS solution and potassium persulfate solution, allowing it to react for 24 hours at room temperature. Various concentrations of respective test samples (extract and positive control) were prepared and individually mixed in 10 μ L of ABTS^{•+} solution. After a 30-minute of incubation, the absorbance of the mixture was measured spectrophotometrically at 750 nm. The scavenging activity was calculated using the formula (1).

20 *Nitric oxide radical scavenging activity*

Nitric oxide was generated from sodium nitroprusside in an aqueous solution and examined using the Griess test. The method of Ramadhan et al. (2019) was adopted to examine the scavenging activity of the plant extract against radical nitric oxide. Briefly, 20 μ L of various extract concentrations was mixed with 5 mM sodium nitroprusside, then 100 μ L of Griess reagent was added to the mixture. After a 30-minute incubation, the absorbance of the mixture was measured spectrophotometrically at 540 nm. The percentage of free radical nitric oxide scavenging activity was calculated using the formula (1). Ascorbic acid was used as a positive control.

Data analysis

All data were expressed as the mean \pm SD of three measurements and analyzed with nonlinear regression analysis. The obtained quantitative data were analyzed descriptively. All measurements were conducted in triplicate.

RESULTS AND DISCUSSION

In vitro antidiabetic activity

The present study was carried out to evaluate the antidiabetic and antioxidant potency of 10 selected medicinal plants from the East Kalimantan tropical forest. The antidiabetic activity was carried out using rat intestinal α -glucosidase on maltase and sucrose as substrates. The α -glucosidase enzyme (EC 3.2.1.20, α -D-glucoside glucosylhydrolase) is the main enzyme located in the small intestine vital to cleave maltose and sucrose into glucose to be further absorbed into the blood. One of the most therapeutic approaches to control postprandial blood sugar levels is to inhibit intestinal α -glucosidase, which is highly similar to human α -glucosidase (Pyner et al. 2017).

In this study, we comprehensively investigated the part of plants such as stems, branches, and twigs instead of leaves traditionally used to treat blood sugar levels in the communities. It is the first study that investigated extracts of ten tropical, medicinal plants concerning rat intestinal α -glucosidase inhibition (maltase and sucrose). The inhibitory activity of extracts is presented in Table 2. Rat intestinal enzymes can only hydrolyze maltose and sucrose. The principle of the assay is to inhibit carbohydrate-hydrolyzing enzymes from delaying the absorption of glucose that mimics the metabolism system (Seo et al. 2015). Methanol extracts of *Flacourtia rukam* Zoll. & Moritzi, *Syzygium chloranthum* (Duthie) Merr. & L.M.Perry, and *Shorea balangeran* Burck exhibited strong inhibition activity against maltase with IC₅₀ values of 0.034 mg/mL, 0.039 mg/mL and 0.039 mg/mL, respectively. Meanwhile, quercetin is a positive control has an IC₅₀ value of 0.040 mg/mL. Proença et al. (2017) and Indrianingsih et al. (2015) reported that quercetin could be used as a positive control since it has a high inhibitory effect on α -glucosidase for its aromatic ring with *ortho* -OH position in the flavonoid structure and it is a natural inhibitor with low side effects.

The present study showed the inhibitory activity of *F. rukam*, *S. chloranthum* and *S. balangeran* are comparable

to quercetin as a positive control. Afifi et al. (2021) found that the bark and leaves of *F. rukam* contains phenolic. Thai et al. (2020) reported several isolated secondary metabolites, including rukamtenol, chaulmoogric acid, flacourtin, 3,4,5-trimethoxyphenyl- β -D-glucopyranoside, and daucosterol. Elfita et al. (2019) also reported the stem bark of *F. rukam* contain friedelin, poliothryoside, and β -sitosteryl-3 β -glucopyranoside. The previous studies show the phytochemical presence in the extract of *F. rukam* can be responsible for the biological activity against α -glucosidase. As far as we know, there is no report on the α -glucosidase inhibitory activity of *F. rukam*, *S. chloranthum*, and *S. balangeran*.

Table 2 showed that *Lagerstroemia speciosa* (L.) Pers., *Vatica umbonata* Burck, *Pertusadina eurhyncha* (Miq.) Ridsdale, and *Gluta velutina* Blume have moderate inhibitory activity against maltase. The order of inhibitory activity of the ten medicinal plant extracts against maltase α -glucosidase from the lowest to the strongest was as follows: *Kleinhovia hospita* L. > *Enterolobium cyclocarpum* (Jacq.) Griseb. > *Dillenia excelsa* (Jack) Martelli ex Gilg > *G. velutina* > *V. umbonata* > *P. eurhyncha* > *L. speciosa* > *S. chloranthum* > *S. balangeran*. This finding is in line with Li et al. (2021), who reported that resorcinol derivatives in *Syzygium latilimbum* had the potency to inhibit α -glucosidase. Furthermore, Hu et al. (2021) also stated that resorcinol derivatives isolated from *Syzygium samarangense* had antidiabetic activity by inhibiting α -glucosidase. Moreover, Kissinger et al. (2016) showed that *S. balangeran* had antidiabetic activity by inhibiting α -glucosidase, possibly due to oligostilbenoids that contain the -OH group.

Moreover, the results showed that *S. chloranthum* extract showed good antidiabetic activity against sucrose with IC₅₀ values of 0.042 mg/mL and exhibited quite similar IC₅₀ values with quercetin as a positive control (Table 2.). On the other hand, *F. rukam*, *G. velutina*, *V. umbonata*, *P. eurhyncha* and *D. excelsa* extracts have no inhibition against sucrose.

Table 2. The IC₅₀ values of α -glucosidase inhibitory activity of selected medicinal plants

Medicinal plants	Part used	IC ₅₀ (mg/mL)	
		Maltase	Sucrose
<i>Shorea balangeran</i> Burck	Branch & Twig	0.039 ± 0.02	0.075 ± 0.01
<i>Syzygium chloranthum</i> (Duthie) Merr. & L.M.Perry	Branch & Twig	0.039 ± 0.01	0.042 ± 0.01
<i>Lagerstroemia speciosa</i> (L.) Pers.	Stem	0.051 ± 0.01	0.062 ± 0.01
<i>Flacourtia rukam</i> Zoll. & Moritzi	Stem	0.034 ± 0.01	NI
<i>Gluta velutina</i> Blume	Stem	0.076 ± 0.02	NI
<i>Kleinhovia hospita</i> L.	Branch & Twig	NI ^b	0.161 ± 0.08
<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	Stem	0.139 ± 0.01	0.086 ± 0.02
<i>Vatica umbonata</i> Burck	Branch & Twig	0.060 ± 0.01	NI
<i>Pertusadina eurhyncha</i> (Miq.) Ridsdale	Stem	0.075 ± 0.01	NI
<i>Dillenia excelsa</i> (Jack) Martelli ex Gilg.	Stem	0.111 ± 0.02	NI
Quercetin ^a		0.040 ± 0.01	0.030 ± 0.02

Note: ^a Positive control; ^b No inhibition, inhibitory effects less than 30% at 0.0625 mg/mL; Each value represents mean ± S.D (n=3)

Plant extracts of *E. cyclocarpum*, *S. balangeran*, and *L. speciosa* have inhibitory activity against sucrose with IC₅₀ values of 0.086 mg/mL, 0.075 mg/mL, and 0.062 mg/mL, respectively while quercetin as positive control with IC₅₀ value of 0.030 mg/mL. A previous Kolakul and Sripanidkulchai (2017) study reported that *L. speciosa* contains phenolic groups such as ellagic acid, epicatechin gallate, and gallic acid. In addition, a previous study by Riyanti et al. (2020) reported *Lagerstroemia loudonii* Teijsm. & Binn exhibited inhibitory activity against yeast α -glucosidase. Our results are in agreement with the previous study by Ratnadewi et al. (2020) showed that *L. speciosa* had the potency as an antidiabetic agent. This study is the first report on the biological activities of ten medicinal plants' methanol extracts from East Kalimantan tropical forests against maltase and sucrose. It indicates that medicinal plants from the tropical forest in Kalimantan had the potential as natural antidiabetic agents. However, further in-depth studies are required to analyze and isolate the chemical compounds responsible for their antidiabetic activity.

Antioxidant properties

Recent studies reported that chronic hyperglycemia generates excessive reactive radicals due to glucose autooxidation and contributes to cell damage (Kanwugu et al. 2021). Therefore, three methods were used to determine the antioxidant activity of selected medicinal plant extracts, i.e., DPPH, ABTS, and Nitric oxide scavenging activities. In addition to the antioxidant activity assay, α -glucosidase inhibition activity was also performed to screen ten medicinal plant extracts as antidiabetic and its complication by free radicals.

The DPPH radical scavenging activity of ten medicinal plant extracts and standard ascorbic acid is shown in Table 3. The percentage of inhibition of ten medicinal plants is presented in Figure 2. These results showed that *S. balangeran* and *G. velutina* had a good inhibition percentage of 79.5 % and 83.1 % at 1 mg/mL extract concentration. The greater the inhibitory activity against DPPH free radicals, the higher the antioxidant activity and the lower the 50% inhibition concentration (IC₅₀) (Adouni et al. 2022). Among ten medicinal plant extracts tested, *S. balangeran* and *G. velutina* displayed the highest antioxidant activity with IC₅₀ values of 0.05 mg/mL and 0.04 mg/mL, respectively, which are similar to the IC₅₀ value of ascorbic acid as a positive control (0.05 mg/mL). Marjoni and Zulfisa (2017) stated that antioxidant activity is categorized as strong if the IC₅₀ value is 0.05-0.1 mg/mL and very strong if the IC₅₀ value is less than 0.05 mg/mL. DPPH radical is stabilized by atoms that can donate hydrogen to form a stable reduced DPPH molecule (Arroussi et al. 2022). Subramanian et al. (2013) showed that the methanol extract of *Shorea roxburghii* stem bark has radical scavenging activity against DPPH. In addition, Subramanian et al. (2015) isolated hopeaphenol, a polyphenol group composed of resveratrol units that exhibit free radical scavenging activity against DPPH. This

study showed that *S. chloranthum* and *D. excelsa* had moderate antioxidant activity against DPPH free radical, indicating the potency as natural antioxidant activity.

Furthermore, the antioxidant capacity of selected medicinal plants was also performed against ABTS⁺ free radical. ABTS⁺ free radical solution was generated by reacting with an oxidizing agent, i.e., potassium persulfate, which took approximately 16 hours and neutralized the radicals based on electron transfer (Xiao et al. 2020). Results of the ABTS⁺ assay showed that selected medicinal plant extracts at 1 mg/mL had the potency to scavenge radicals (Figure 2.). The extracts of *S. balangeran*, *S. chloranthum* and *F. rukam* demonstrated the highest scavenging activity against ABTS⁺ free radical, i.e., 90.2%, 88.3%, and 91.4%, respectively. Table 3 shows the SC₅₀ values of ABTS⁺ free radical scavenging activity of selected medicinal plant extracts in the following order: *G. velutina* > *V. umbonata* > *P. eurhyncha* > *E. cyclocarpum* > *L. speciosa* = *D. excelsa* > *K. hospita* > *S. chloranthum* > *S. balangeran* > *F. rukam*. The greater the scavenging activity of the selected medicinal plant extracts against ABTS⁺, the higher their antioxidant activity and the lower their 50% scavenging concentration (SC₅₀). *Flacourtia* species contain phenolic compounds. Bourjot et al. (2012) showed that *F. rukam* stem bark contained phenolic glycosides (Flacourtosides A-F). Sashidhara et al. (2014) also discovered polythryoside from *Flacourtia indica*. It is the first study on the antioxidant and scavenging activity of selected medicinal plants collected from East Kalimantan against ABTS⁺.


Moreover, the antioxidant efficiency of selected medicinal plants was also evaluated against nitric oxide free radicals. Nitric oxide (NO) plays an important role in inflammatory processes. The overproduction of NO leads to tissue damage and acute and chronic inflammation. Chronic inflammation by nitric oxide free radicals is associated with various diseases, such as juvenile diabetes, arthritis, and ulcerative colitis (Venkatachalam and Muthukrishnan 2012). Therefore, nitric oxide free radical inhibitory agents (selected medicinal plants) might be beneficial in treating inflammatory responses. In this study, *S. balangeran*, *F. rukam* and *G. velutina* had similar antioxidant activity against nitric oxide free radicals with IC₅₀ values shown in Table 3. Ascorbic acid used as a positive control for nitric oxide radical scavenger in this assay showed a scavenging percentage of 77.5% with an IC₅₀ value of 0.05 mg/mL. The order of IC₅₀ values of selected medicinal plants from the lowest to the highest is as follows: *S. chloranthum* > *K. hospita* > *L. speciosa* > *S. balangeran* = *F. rukam* = *G. velutina*. The selected medicinal plant extract inhibits nitric oxide free radicals by inhibiting nitrite formation in the radical-generating process through competition with oxygen (Habu and Ibeh 2015).

This study is the first work on ten selected medicinal plants from East Kalimantan that exhibit scavenging activity against several free radicals, such as DPPH, ABTS, and Nitric Oxide.

Table 3. Antioxidant activity of selected medicinal plants against free radicals

	Part used	SC ₅₀ (mg/mL)		
		DPPH	ABTS	Nitric oxide
<i>Shorea balangeran</i> Burck	Branch & Twig	0.05 ± 0.01	0.08 ± 0.01	0.11 ± 0.03
<i>Syzygium chloranthum</i> (Duthie) Merr. & L.M.Perry	Branch & Twig	0.11 ± 0.01	0.21 ± 0.01	0.16 ± 0.01
<i>Lagerstroemia speciosa</i> (L.) Pers.	Stem	0.26 ± 0.05	0.28 ± 0.07	0.12 ± 0.01
<i>Flacourtia rukam</i> Zoll. & Moritz	Stem	0.39 ± 0.01	0.05 ± 0.01	0.11 ± 0.01
<i>Gluta velutina</i> Blume	Stem	0.04 ± 0.01	0.95 ± 0.05	0.11 ± 0.01
<i>Kleinhovia hospita</i> L.	Branch & Twig	0.78 ± 0.41	0.25 ± 0.05	0.13 ± 0.02
<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	Stem	NI ^b	0.43 ± 0.19	NI
<i>Vatica umbonata</i> Burck	Branch & Twig	NI	0.63 ± 0.10	NI
<i>Pertusadina eurhyncha</i> (Miq.) Ridsdale	Stem	0.89 ± 0.02	0.45 ± 0.11	NI
<i>Dillenia excelsa</i> (Jack) Martelli ex Gilg.	Stem	0.13 ± 0.01	0.28 ± 0.07	NI
Ascorbic acid ^a	5	0.05 ± 0.01	0.12 ± 0.01	0.05 ± 0.02

Note: ^a Positive control. ^b No inhibition, inhibitory effects less than 45% at 1 mg/mL. Each value represents the mean ± S.D (n=3)

**Figure 2.** Inhibition percentage of medicinal plant extracts against free radicals (at extract concentration of 1 mg/mL)

In summary, the results in this study indicate that ten selected medicinal plant extracts from East Kalimantan exhibited dual functions as antidiabetic and antioxidant through both rat intestinal α -glucosidase (maltase and sucrase) inhibitory activity and free radical scavenging activity against DPPH, ABTS, and nitric oxide. However, the study area has never been explored ethnobotanically. In this regard, the present study can be considered the first one, which deals with an ethnobotanical study guided by biological activities on medicinal plants in this region. Hence, these medicinal plant extracts might be used as natural agents with great importance as therapeutic agents in preventing the progress of hyperglycemia associated with oxidative stress related to degenerative diseases. In addition, this study provides important findings for the bioassay guide to isolate bioactive secondary metabolites from

selected medicinal plants. Therefore, further studies are needed to investigate individual constituents from some actively selected medicinal plant extracts that may be responsible for anti-diabetes and its complications caused by free radicals.

ACKNOWLEDGEMENTS

R.R is grateful to the Universitas Airlangga, Surabaya, Indonesia, for providing research funds through the Hibah Riset Mandat Kolaborasi Mitra Luar Negeri (797/UN3.15/PT/2021) program. Furthermore, R.R is thankful to the Laboratory of Forest Products Chemistry and Renewable Energy, Faculty of Forestry, Universitas Mulawarman, Samarinda, Indonesia, for its plant deposit.

REFERENCES

- Abuga I, Sulaiman SF, Wahab RA, Ooi KL, Rasad MBSA. 2022. Phytochemical constituents and antibacterial activities of 45 Malay traditional medicinal plants. *J Herb Med* 32: 100496. DOI: 10.1016/j.hermed.2021.100496.
- Adouni K, Bendif H, Zouaoui O, Kraujalis P, Flamini G, Venskutonis PR, Achour L. 2022. Antioxidant activity of extracts obtained by high-pressure extraction procedures from *Asparagus stipularis* Forssk. *S Afr J Bot* 146: 789-793. DOI: 10.1016/j.sajb.2021.12.027.
- Afifi NI, Moawad AS, Zaki MA, Rateb ME, Rashed MH, Saleh IG, Hetta MH, Mohammed RM. 2021. Four new phenolic and antiparasitic secondary metabolites from *Flacourtia rukam* Zoll. & Mortizi. *Nat Prod Res* 2021. DOI: 10.1080/14786419.2021.1875462.
- Akyuz M, Yabo-Dambagi L, Kilic T, Cakir A. 2022. Antidiabetic, neuroprotective and antioxidant potentials of different parts of *Pistacia terebinthus* fruits. *S Afr J Bot* 147: 443-456. DOI: 10.1016/j.sajb.2022.01.040.
- Arroussi J, Ouerfelli M, Smaoui A, Ahmed HB, Kaâb LBB. 2022. Antioxidant activity of seven plant extracts collected from Tunisia and their allelopathic potential on *Lactuca sativa* L. and *Phalaris minor* L. *S Afr J Bot* 148: 135-143. DOI: 10.1016/j.sajb.2022.04.029.
- Arung ET, Kuspradini H, Kusuma IW, Shimizu K, Kondo R. 2012. Validation of *Eupatorium triplinerve* Vahl leaves, a skin care herb from East Kalimantan, using a melanin biosynthesis assay. *J Acupunct Meridian Stud* 5: 87-92. DOI: 10.1016/j.jams.2012.01.003.
- Bourjot M, Leysen P, Eydoux C, Guillemot JC, Canard B, Rasoanaivo P, Guéritte F, Litaudon M. 2012. Flacourtosides A-F, phenolic glycosides isolated from *Flacourtia ramontchi*. *J Nat Prod* 15: 752-758. DOI: 10.1021/np300059n.
- Elfita M, Yohandini H, Julinar, Yasrina, Miranti. 2019. Chemical Constituents from Stem Bark of *Flacourtia rukam* Zoll. & Morand Their Antioxidant Activities. *Sains Malays* 48: 1899-1906. DOI: 10.17576/jsm-2019-4809-10.
- Habu JB, Ibeh BO. 2015. In vitro antioxidant capacity and free radical scavenging evaluation of active metabolite constituents of *Newbouldia laevis* ethanolic leaf extract. *Biol Res* 48: 1-10. DOI: 10.1186/s40659-015-0007-x.
- Hu YK, Wang L, Wang JH, Li MJ, Li F, Yang J, Zhao Y. 2021. Resorcinol derivatives with α -glucosidase inhibitory activities from *Syzygium samarangense*. *Nat Prod Res* 35: 5948-5953. DOI: 10.1080/14786419.2020.1805606.
- Indrianingsih AW, Tachibana S, Itoh K. 2015. In vitro evaluation of antioxidant and α -glucosidase inhibitory assay of several tropical and subtropical plants. *Procedia Environ Sci* 28: 639-648. DOI: 10.1016/j.proenv.2015.07.075.
- Ji Y, Liu D, Jin Y, Zhao J, Zhao J, Li H, Li L, Zhang H, Wang H. 2021. In vitro and in vivo inhibitory effect of anthocyanin-rich bilberry extract on α -glucosidase and α -amylase. *LWT* 145: 111484. DOI: 10.1016/j.lwt.2021.111484.
- Kanwugu ON, Glukhareva TV, Danilova IG, Kovaleva EG. Natural antioxidant in diabetes treatment and management: prospect of astaxanthin. *Crit Rev Food Sci Nutr* 2021. DOI: 10.1080/10408398.2021.1881434.
- Khan I, Ullah Z, Shad AA, Fahim M, Öztürk M. In vitro antioxidant, anticholinesterase inhibitory, and antimicrobial activity studies of *Terminalia chebula* (Retz) and *Terminalia arjuna* (Roxb). *S Afr J Bot* 146: 395-400. DOI: 10.1016/j.sajb.2021.11.016.
- Kissinger, Yamani A, Pitri RMN. 2016. α -Glucosidase inhibitory activity of *Shorea belangeri* of kerangas forest. *Res. J. Med. Plant* 10: 356-361. DOI: 10.3923/rjmp.2016.356.361.
- Kolakul P, Sripanidkulchai B. 2017. Phytochemicals and anti-aging potentials of the extracts from *Lagerstroemia speciosa* and *Lagerstroemia floribunda*. *Ind Crops Prod* 109: 707-716. DOI: 10.1016/j.indcrop.2017.09.026.
- Kusuma IW, Murdiyanto, Arung ET, Syafrizal, Kim Y. 2014. Antimicrobial and antioxidant properties of medicinal plants used by the Bentian tribe from Indonesia. *Food Sci Hum Wellness* 3: 191-196. DOI: 10.1016/j.fshw.2014.12.004.
- Lee SK, Ryu SH, Turk A, Yeon SW, Jo YH, Han YK, Hwang BY, Lee KY, Lee MK. 2020. Characterization of α -glucosidase inhibitory constituents of the fruiting body of lion's mane mushroom (*Hericium ericaceus*). *J. Ethnopharmacol* 262: 113197. DOI: 10.1016/j.jep.2020.113197.
- Li YY, Hu YK, Wang JH, Huang YP, Zeng GJ, Yang Y, Zhao Y. 2021. α -Glucosidase inhibitors from *Syzygium latilimum*. *Chem Nat Compd* 57: 551-552. DOI: 10.1007/s10600-021-03412-z.
- Liu X, Zhu L, Tan J, Zhou X, Xiao L, Yang X, Wang B. 2014. Glucosidase inhibitory activity and antioxidant activity of flavonoid compound and triterpenoid compound from *Agrimonia pilosa* Ledeb. *BMC Complement Altern Med* 14: 1-10. DOI: 10.1186/1472-6882-14-12.
- Marjoni MR and Zulhisa A. 2017. Antioxidant activity of methanol extract/fractions of Senggani leaves (*Melastoma candidum* D. Don). *Pharm Anal Acta* 8: 1-6. DOI: 10.4172/2153-2435.1000557.
- Mbopi PY, Fozeng HDS, Nguekeu YMM, Bitchagno GTM, Ngantchouko CBN, Awouafack MD, Opatz T, Ngouela SA, Morita H, Tene M. 2021. Chemical constituents, total phenolic content, antioxidant activity and bactericidal effect of *Dicliptera verticillate* (Acanthaceae). *S Afr J Bot* 142: 216-221. DOI: 10.1016/j.sajb.2021.07.001.
- Mipun P, Bhat NA, Borah D, Kumar Y. Non-timber forest products and their contribution to healthcare and livelihood security among the Karbi tribe in Northeast India. *Ecol Process* 8: 1-21. DOI: 10.1186/s13717-019-0194-4.
- Moukette BM, Pieme CA, Nji,ou JR, Biapa CPN, Ngogang JY. 2015. In vitro antioxidant properties, free radicals scavenging activities of extracts and polyphenol composition of a non-timber forest product used as spice: *Monodora myristica*. *Biol Res* 48 (1): 1-17. DOI: 10.1186/s40659-015-0003-1.
- Proença C, Freitas M, Ribeiro D, Oliveira EFT, Sousa JLC, Tome SM, Ramos MJ, Silva AMS, Fernandes PA, Fernandes E. 2017. α -Glucosidase inhibition by flavonoids: An in vitro and in silico structure-activity relationship study. *J Enzyme Inhib Med Chem* 32: 1216-1228. DOI: 10.1080/14756366.2017.1368503.
- Pyner A, Nyambe-Silavwe H, Williamson G. 2017. Inhibition of human and rat sucrase and maltase activities to assess antiglycemic potential: Optimization of the assay using acarbose and polyphenols. *J Agric Food Chem* 65: 8643-8651. DOI: 10.1021/acs.jafc.7b03678.
- Ramadhan R, Kristanti AN, Amirta R, Kusuma IW, Phuwapraisrisan P, Haqiqi MT, Saparwadi. 2019. Screening for potential antidiabetes and antioxidant activities of selected plants from East Kalimantan, Indonesia. *Biodiversitas* 20: 1820-1826. DOI: 10.13057/biodiv/d200705.
- Ramadhan R, Phuwapraisrisan P, Kusuma IW, Amirta R. 2020. Ethnopharmacological evaluation of selected East Kalimantan flora for diabetes therapy: The isolation of lupane triterpenoids as α -glucosidase inhibitors from *Ceripogon gal* (Perr). C.B. Robb. *Rasayan J Chem* 13: 1727-1734. DOI: 10.31788/RJC.2020.1335559.
- Ratnadewi AAI, Wahyudi LD, Rochman J, Susilowati, Nugraha AS, Siswoyo TA. 2020. Revealing antidiabetic potency of medicinal plants of Meru Betiri Nasional Park, Jember - Indonesia. *Arab J Chem* 13: 1831-1836. DOI: 10.1016/j.arabj.2018.01.017.
- Riyanti S, Dewi PS, Windyaswari AS, Azizah SAN. 2020. Alpha-glucosidase inhibitory activities of bungur (*Lagerstroemia loudonii* Teijsm. & Binn.) leaves and fruits. *IOP Conf Ser Earth Environ Sci* 462: 012042. DOI: 10.1088/1755-1315/462/1/012042.
- Sashidhara KV, Singh SP, Varshney S, Beg M, Gaikwad AN. 2014. Poliothryoside and its derivatives as novel insulin sensitizers potentially driving AMPK activation and inhibiting adipogenesis. *Eur J Med Chem* 86: 570-577. DOI: 10.1016/j.ejmech.2014.09.015.
- Seo KH, Ra JE, Lee SJ, Lee JH, Kim SR, Lee JH, Seo WD. 2015. Anti-hyperglycemic activity of polyphenols isolated from bamyard millet (*Echinochloa utilis* L) and their role inhibiting α -glucosidase. *J Korean Soc Appl Biol Chem* 58: 571-579. DOI: 10.1007/s13765-015-0070-6.
- Seyrekoglu F, Temiz H, Eser F, Yildirim C. 2022. Comparison of antioxidant activities and major constituents of three *Hypericum* species (*H. perforatum*, *H. scabrum* and *H. origanifolium*) from Turkey. *S Afr J Bot* 146: 723-727. DOI: 10.1016/j.sajb.2021.12.012.
- Shackleton CM, Ticktin T, Cunningham AB. 2018. Nontimber forest products as ecological and biocultural keystone species. *Ecol* 23: 22. DOI: 10.5751/ES-10469-230422.
- Subramanian R, Raj V, Manigandan K, Elangovan. 2015. Antioxidant activity of hopeaphenol isolated from *Shorea roxburghii* stem bark extract. *J Taibah Univ Sci* 9: 237-244. DOI: 10.1016/j.jtusc.2014.11.004.
- Subramanian R, Subramanian P, Raj V. 2013. Antioxidant activity of the stem bark of *Shorea roxburghii* and its silver reducing power. *Springerplus* 2: 1-11. DOI: 10.1186/2193-1801-2-28.

- Sukara E. 2014. Tropical forest biodiversity to provide food, health and energy solution of the rapid growth of modern society. *Procedia Environ Sci* 20: 803-808. DOI: 10.1016/j.proenv.2014.03.097.
- Thai CT, Duong TH, Trung NT, Phan DC, Nguyen QV, Nguyen TL, Wanatabe K, Do TT, Huynh TN, Tran H. 2020. Rukamtenol, a new spiro compound isolated from *Flacourtia rukam* Zoll. & Mortizi growing in Vietnam. *Nat Prod Res* 36: 1966-1972. DOI: 10.1080/14786419.2020.1839451.
- Trinh BTD, Staerk D, Jäger AK. 2016. Screening for potential α -glucosidase and α -amylase inhibitory constituents from selected Vietnamese plants used to treat type 2 diabetes. *J Ethnopharmacol* 186: 189-195. DOI: 10.1016/j.jep.2016.03.060.
- Venkatachalam U, Muthukrishnan S. 2012. Free radical scavenging activity of ethanolic extract of *Desmodium gangeticum*. *J Acute Med* 2: 36-42. DOI: 10.1016/j.jacme.2012.04.002.
- Xiao F, Xu T, Lu B, Liu R. 2020. Guidelines for antioxidant assays for food components. *Food Front* 1: 60-69. DOI: 10.1002/fft2.10.

Evaluation of non-timber forest products used as medicinal plants from East Kalimantan (Indonesia) to inhibit α -glucosidase and free radicals

ORIGINALITY REPORT

13%

SIMILARITY INDEX

9%

INTERNET SOURCES

13%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1

oak.go.kr

Internet Source

1%

2

Bibi, Tahira, Mushtaq Ahmad, Rsool Bakhsh Tareen, Niaz Mohammad Tareen, Rukhsana Jabeen, Saeed-Ur Rehman, Shazia Sultana, Muhammad Zafar, and Ghulam Yaseen.

"Ethnobotany of medicinal plants in district Mastung of Balochistan province-Pakistan", *Journal of Ethnopharmacology*, 2014.

Publication

1%

3

ojs.openagrar.de

Internet Source

1%

4

Van Bon Nguyen, Quang Vinh Nguyen, Anh Dzung Nguyen, San-Lang Wang. "Screening and evaluation of α -glucosidase inhibitors from indigenous medicinal plants in Dak Lak Province, Vietnam", *Research on Chemical Intermediates*, 2016

Publication

1%

5	docksci.com Internet Source	1 %
6	www.internationalscholarsjournals.com Internet Source	<1 %
7	chimie-biologie.ubm.ro Internet Source	<1 %
8	rjptonline.org Internet Source	<1 %
9	"Herbal Medicine in India", Springer Science and Business Media LLC, 2020 Publication	<1 %
10	Submitted to University of Johannesburg Student Paper	<1 %
11	pure.ewha.ac.kr Internet Source	<1 %
12	www.journal-jams.org Internet Source	<1 %
13	Fatemeh Sarmadi, Parinaz Kazemi, Pouria Tirgar, Samaneh Fayazi et al. "Using natural honey as an anti-oxidant and thermodynamically efficient cryoprotectant in embryo vitrification", Cryobiology, 2019 Publication	<1 %
14	KR. Nagulendran, S. Velavan, R. Mahesh, V. Hazeena Begum. " Antioxidant Activity and	<1 %

Total Polyphenolic Content of Rhizomes ", E- Journal of Chemistry, 2007

Publication

15

Sristicheta Misra, Krishna Bihari Pandeya,
Ashok Kumar Tiwari, Amtul Zehra Ali et al.
"Antihyperglycemic, α -glucosidase inhibitory
and DPPH free radical scavenging activity of
5-bromosalicylaldehyde and schiff bases",
Medicinal Chemistry Research, 2010

Publication

16

jjbs.hu.edu.jo

Internet Source

17

Meriam El Aboubi, Douae Ben Hdech, Samir
Bikri, Asmae Benayad, Anouar El Magri,
Youssef Aboussaleh, El Mahjoub Aouane. "
Chemical composition of essential oils of peel
from three Moroccan regions and their
antioxidant, anti-inflammatory, antidiabetic
and dermatoprotective properties ", Journal of
Herbmed Pharmacology, 2022

Publication

18

Namdeo Gaonkar, Rakhee Dangi.
"Comparative evaluation of antioxidant,
antiglycation and α -glucosidase inhibitory
potential of some indigenous medicinal
Trigonella species", Vegetos, 2022

Publication

<1 %

<1 %

<1 %

<1 %

- 19 Yanglin Ji, Dong Liu, Yan jin, Juan Zhao, Jiang Zhao, Heyu Li, Liwei Li, Hua Zhang, Hao Wang. "In vitro and in vivo inhibitory effect of anthocyanin-rich bilberry extract on α -glucosidase and α -amylase", LWT, 2021
Publication <1 %
-
- 20 ijpsr.com
Internet Source <1 %
-
- 21 Salehi, Ata, V. Anil Kumar, Sharopov et al. "Antidiabetic Potential of Medicinal Plants and Their Active Components", Biomolecules, 2019
Publication <1 %
-
- 22 Submitted to Universiti Putra Malaysia
Student Paper <1 %
-
- 23 E. Hamad, M. Hassan, I. Ashoush. "QUALITY CHARACTERISTICS AND ANTIOXIDANT ACTIVITY OF STIRRED YOGHURT FORTIFIED WITH DESERT TRUFFLE", Journal of Food and Dairy Sciences, 2016
Publication <1 %
-
- 24 Submitted to International Islamic University Malaysia
Student Paper <1 %
-
- 25 Kai Hu, Dongyan Chen, Zhida Sun. "Structures, physicochemical properties, and hypoglycemic activities of soluble dietary <1 %

fibers from white and black glutinous rice bran: a comparative study", Food Research International, 2022

Publication

26

Submitted to University of Wales Institute, Cardiff

Student Paper

<1 %

27

biolres.biomedcentral.com

Internet Source

<1 %

28

onlinelibrary.wiley.com

Internet Source

<1 %

29

Beauty E Omoruyi, Graeme Bradley, Anthony J Afolayan. "Antioxidant and phytochemical properties of *Carpobrotus edulis* (L.) bolus leaf used for the management of common infections in HIV/AIDS patients in Eastern Cape Province", BMC Complementary and Alternative Medicine, 2012

Publication

<1 %

30

Dommati Anand Kumar, Sanga Venkata Anusha, Swathi Oruganti, Manaswini Deshpande, Amtul Zehra, Ashok Kumar Tiwari. "Raw versus cooked vegetable juice", Nutrafoods, 2015

Publication

<1 %

31

Sara El-Leithy, Tahany Mazal, Samah Aamer. "Performance and Yield Stability of some

<1 %

Egyptian Rice (*Oryza sativa* L.) Cultivars under Different Environments", Journal of Agricultural Chemistry and Biotechnology, 2021

Publication

32

Ahmed Nokhala, Qamar Uddin Ahmed, Mohammed S. M. Saleh, Tanzina Sharmin Nipun et al. "Characterization of α -glucosidase inhibitory activity of *Tetracera scandens* leaves by Fourier transform infrared spectroscopy-based metabolomics", *Advances in Traditional Medicine*, 2019

<1 %

Publication

33

Hasnah M. Sirat, Mohd Fazlin Rezali, Zanariah Ujang. " Isolation and Identification of Radical Scavenging and Tyrosinase Inhibition of Polyphenols from L. ", *Journal of Agricultural and Food Chemistry*, 2010

<1 %

Publication

34

Leila Kebal, Katarzyna Pokajewicz, Noureddine Djebli, Nadjat Mostefa, Anna Poliwoda, Piotr P. Wieczorek. "HPLC-DAD profile of phenolic compounds and In vitro antioxidant activity of *Ficus carica* L. fruits from two Algerian varieties", *Biomedicine & Pharmacotherapy*, 2022

<1 %

Publication

35

eprints.lums.ac.ir

Internet Source

<1 %

36 pubag.nal.usda.gov
Internet Source

<1 %

37 www.frontiersin.org
Internet Source

<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On