

The potential of Macaranga plants as skincare cosmetic ingredients: A review

by Rudianto Amirta

Submission date: 10-May-2023 01:24PM (UTC+0700)

Submission ID: 2089270374

File name: f_Macaranga_plants_as_skincare_cosmetic_ingredients_A_review.pdf (982.49K)

Word count: 8864

Character count: 55563



The potential of *Macaranga* plants as skincare cosmetic ingredients: A review

Enih Rosamah¹, Muhammad Taufiq Haqiqi¹, Agmi Sinta Putri¹, Harlinda Kuspradini¹, Irawan Wijaya Kusuma¹, Rudianto Amirta¹, Yuliansyah Yuliansyah¹, Wiwin Suwinarti¹, Swandari Paramita², Rico Ramadhan³, Didi Tarmadi⁴, Maya Ismayati⁵, Harits Atika Ariyanta⁵, Widya Fatriasari⁵, Aswandi Aswandi⁵, Cut Rizlani Kholibrina⁵, Nur Izyan Wan Azelee⁶, Yong-ung Kim⁷, Enos Tangke Arung^{1*}

¹Faculty of Forestry, Mulawarman University, Samarinda, Indonesia.

²Faculty of Medicine, Mulawarman University, Samarinda, Indonesia.

³Department of Chemistry, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia.

⁴Research Center for Applied Zoology, National Research and Innovation Agency, Bogor, Indonesia.

⁵Research Center for Biomass and Bioproducts, National Research and Innovation Agency, Bogor, Indonesia.

⁶Institute of Bioproduct Development (IBD), Universiti Teknologi Malaysia, UTM Skudai, Johor, Malaysia.

⁷Department of Pharmaceutical Engineering, College of Cosmetics and Pharmaceuticals, Daegu Haany University, Gyeongsangbuk-do, South Korea.

ARTICLE INFO

Received on: 04/12/2022

Accepted on: 11/01/2023

Available Online: XX

Key words:

Cosmetics, fast-growing plant, industrial product, *Macaranga*, skincare.

ABSTRACT

The application of bioactive ingredients extracted from plants utilized as additives in various cosmetic products has gained popularity, since they are safe, with low adverse effects when applied properly, and are environmentally friendly. Moreover, the awareness of healthier cosmetic products has rapidly increased so that the exploration of screened plants with appropriate properties has attracted significant attention worldwide. This review discusses the potential of one of the fastest-growing tropical plants, *Macaranga*, applied in skincare cosmetics. Their interesting characteristics, such as being anti-inflammatory, antioxidant, and antimicrobial, and their tyrosinase inhibitory effect have been comprehensively summarized. Various scientific literature works have further proven ethnopharmacological studies that explore the traditional use of *Macaranga* species by local people in the tropics for medicinal and skincare purposes. Therefore, we believe this will allow *Macaranga* to become a promising material in the future for large-scale industrial skincare cosmetics.

INTRODUCTION

The largest organ on the outside of the human body, the skin, is a barrier to shield interior organs from dehydration, microbial infection, and UV exposure. Besides these important functions, it is well known that a better skin appearance is also responsible for an individual's awareness of looking more

beautiful. Hence, various companies have attempted to pursue improving their innovative products in skincare cosmetics. The business sector in this area has been recognized as one of the most promising ones, with an annual revenue worth of billions of dollars (Kouassi *et al.*, 2022). Currently, a phenomenon in healthy lifestyles contributes to shifting from synthetic skincare cosmetics to green ones, since synthetic materials reportedly have harmful side effects, such as low absorption ability and allergic reactions (irritation) (Morais *et al.*, 2021). A previous study also indicated their negative influences on the environment (Amberg and Fogarassi, 2019). On the other hand, green technology offers several advantages, including safety, nontoxicity, having no adverse effects when used correctly, and biodegradability. These

*Corresponding Author

Enos Tangke Arung, Faculty of Forestry,

Mulawarman University, Samarinda, Indonesia.

E-mail: tangkearung@yahoo.com

products are generally fabricated by incorporating bioactive compounds obtained from plants. Therefore, this trend motivates scientists worldwide to explore plant materials with appropriate properties, particularly those used in skincare cosmetics.

Euphorbiaceae is considered to be one of the largest families among flowering plants, with 218 genera and 5,735 species (Zixi *et al.*, 2016). According to ethnobotanical studies, most plants in this family are frequently utilized in local medicine and skincare (Zahidin *et al.*, 2017). *Macaranga* is a genus of Euphorbiaceae, which consists of more than 300 species, and it is widely distributed across tropical and subtropical countries in Africa, Asia, Australasia, and the Pacific (Huonga *et al.*, 2019). Nevertheless, its distribution center has been recorded in tropical Asia (Syah and Ghisalberti, 2015). This fast-growing tree species, also known as the highest genus in the Euphorbiaceae, can be found in gaps between the forest canopy, disturbed forests, and open areas (Tanjung *et al.*, 2018). Due to the change in light conditions after logging activities or forest fires, some *Macaranga* species are usually found with high dominance as pioneer species. The presence of *Macaranga* can be a reliable bioindicator of deforestation (Zapanta *et al.*, 2019). Its rapid growth has been recorded to significantly influence the composition of the forest vegetation community during the secondary succession stage (Susanto *et al.*, 2018). In Indonesia, its wide distribution has been found in Sumatera and on Kalimantan Island (Tanjung *et al.*, 2018). Considering its ability to cover land quickly and attract wildlife, it is also becoming popular as a species to promote revegetation after coal mining in this country (Amirta and Candra, 2016). The leaves of several species from this genus are often utilized in traditional medicine and have the potential for use as antipyretics, antitussives, and anti-inflammatories (Pailee *et al.*, 2015). Due to its good adaptation to various types of land, even with low available nutrient status, combined with its interesting properties, further domestication of *Macaranga* will allow the use of these plant species in various applications, including sustainable raw materials for additives in skincare production.

Nowadays, the practice of using direct herbal products for skincare purposes has been declining, since various modern skincare products containing bioactive ingredients with simpler usage have been introduced to the market (Milito *et al.*, 2021). As a result, valuable information about crude extraction and isolated compounds, including those obtained from *Macaranga*, will need to be collected, as several authors have reported its remarkable bioactivity (Di *et al.*, 2020; Hashim *et al.*, 2022; Minarti *et al.*, 2021). This review presents a comprehensive summary of the chemical and pharmacological studies on *Macaranga* spp. in terms of their suitability for use as skincare cosmetic products. Furthermore, the range of their potential application in large-scale industrial skincare products is also considered.

METHODOLOGY

The current review was conducted based on a number of publications, specifically, plant extracts and bioactive constituents of various *Macaranga* plants, which can be used as skincare cosmetic ingredients. Other potential applications of *Macaranga*'s bioactive constituents are also presented. There is no limitation on the year of publication. Exhaustive searches were performed on four electronic databases: PubMed, ScienceDirect, Scopus,

and Google Scholar. Various keywords, such as *Macaranga* plant, *Macaranga* genus and species, chemical compounds of *Macaranga*, bioactivity of *Macaranga*, anti-inflammatory activity, antioxidant activity, antimicrobial activity, and antityrosinase activity, were employed to find appropriate data and articles.

Bioactive constituents of *Macaranga* plants

An earlier work summarized the chemical investigations of 25 species of *Macaranga* (Magadula, 2014). In this study, the phytochemical characterization has been updated to include 40 species. They are identified as *M. adantha*, *M. allorobinsonii*, *M. alnifolia*, *M. balansae*, *M. barteri*, *M. bicolor*, *M. conglomerata*, *M. conifera*, *M. constricta*, *M. deheiculata*, *M. denticulata*, *M. gigantea*, *M. gigantifolia*, *M. hemsleyana*, *M. heynei*, *M. hosei*, *M. hurifolia*, *M. hypoleuca*, *M. indica*, *M. javanica*, *M. kurzii*, *M. lowii*, *M. magna*, *M. mappa*, *M. monandra*, *M. peltata*, *M. pleiostemona*, *M. pruinosa*, *M. pustulata*, *M. recurvata*, *M. rhizinoides*, *M. rubiginosa*, *M. sampsonii*, *M. schweinfurthii*, *M. siamensis*, *M. sinensis*, *M. tanarius*, *M. trichocarpa*, *M. triloba*, and *M. vedeliana*. This list reveals, however, that only a small percentage of species (less than 15%) have been further chemically investigated, despite the fact that there are more than 300 species of *Macaranga* available in the world.

Each plant species is distinctive regarding its bioactive constituents. Based on the previous study by Mai *et al.* (2020), it has been discovered that some *Macaranga* species are rich in phenolic compounds, particularly flavonoid and stilbene derivatives. Yang *et al.* (2015a, 2015b) reported the presence of two novel stilbenes, denticulatains A and B, and flavonoid derivatives, including denticulatain C, D, and E from *M. denticulata* fronds. C-methylated and isoprenylated chalcone derivatives, denticalcones A, B, and C, were also discovered in related species. According to Pailee *et al.* (2015), macasiamenenes A-U, macasiamenin A, macasiamenone A, and macasiamenols A and B, were isolated from *M. siamensis* leaves and twigs. Furthermore, secondary metabolites in the class of tannins, terpenes, coumarins, steroids, and many other compounds were also detected. The most prevalent of these is scopoletin, a coumarin that is present in six different species of *Macaranga*: *M. barteri*, *M. conglomerata*, *M. denticulata*, *M. kurzii*, *M. magna*, and *M. triloba*. The number of compounds isolated from *M. tanarius* has been reported as the highest, with 98 compounds collected from its leaves, stem bark, and fruits. Mostly, the compounds in *Macaranga* are stilbenes and flavonoids, as shown in Figure 1. A list of all summarized constituents is presented in Table 1.

Application of *Macaranga* in cosmetics

Anti-inflammatory activity

Inflammation is the response of body to any kind of disturbance caused by irritation, injury, or bacteria. Anti-inflammatory products work by blocking the body response to this skin disorder by inhibiting the effects of certain enzymes that contribute to swelling and inflammation (Attiq *et al.*, 2018). In terms of both skin appearance and health, dealing with inflammation is a major issue. Preventing the impact of skin inflammation is essential for a youthful appearance and avoiding the development of chronic or acute skin diseases




Figure 1. Structure of several stilbenes and flavonoid derivatives isolated from *Macaranga tanarius*.

(Hoang *et al.*, 2021). In general, anti-inflammatory skincare products contain antioxidants, which help the body fight damaging free radicals. Antioxidants prevent collagen breakdown and skin DNA damage (Lin *et al.*, 2018).

An earlier study found that dichloromethane extracts of *M. siamensis* leaves and twigs had a strong anti-inflammatory effect, with Macasiamenene F isolated from these extracts reducing tumor necrosis factor (TNF) alpha by 20% after 24 hours (Leláková *et al.*, 2020). *M. barteri* methanol extract had a strong anti-inflammatory effect with potential compounds such as macabarlerin, 3-O-methylellagic acid, 4-O-b-D-xylopyranoside, ellagic acid, 3-O-methylellagic acid, and gallic acid (Ngoumfo *et al.*, 2008). Moreover, an assay on rats with carrageenan-induced edema showed that hydroalcoholic *M. barteri* bark extract reduced inflammation and skin hyperalgesia. This report also confirmed the traditional application of *M. barteri*'s bark to treat pain and inflammation, acting as an analgesic (Asante-Kwatia *et al.*, 2019). *M. tanarius* demonstrated high inhibition of protein (albumin) denaturation with IC_{50} values of 0.26–1.02 mm with some bioactive

compounds such as nymphaeol A, nymphaeol B, nymphaeol C, isonymphaeol B, and 3'-geranyl-naringenin (Shahinozzaman *et al.*, 2021). Nymphaeol B of this species strongly inhibits acetylcholinesterase (Ache) at 50 $\mu\text{g/ml}$ (Amir Rawa *et al.*, 2022). Another study reported that *M. hurifolia* extract from Nigeria has an inflammatory effect (69.6%) at a dose of 300 mg/kg (Segun *et al.*, 2019a).

Antioxidant activity

Natural antioxidants are used in the cosmetics industry for their capacity to reduce oxidative stress on the skin and protect products from oxidative deterioration (He *et al.*, 2021; Hoang *et al.*, 2021). Due to the natural aging process as well as extrinsic causes including UV radiation, air pollution, and pathogenic microorganisms, oxidative stress is the main factor accelerating skin aging (Farage *et al.*, 2008; Rees, 2004). Antioxidant molecules prevent radical chain reactions, which inhibit reactive oxidants' formation; as a result, they can also be used to cure cancer. In cosmetic products, including serums and creams, antioxidants can be employed to stabilize ingredients and prevent the rancidity of

Table 1. Current status of Bioactive Compounds Isolated from *Macaranga* species.

No.	Plant species	Location	Plant part	Constituents	Sources
1	<i>M. adenantha</i>	China	Twigs	Macadenanthin A, macadenanthin B, macadenanthin C, glyasperin A, broussonflavonol F, and macarangin	Yang <i>et al.</i> (2015c)
2	<i>M. allorobinsonii</i>	Indonesia	Leaves	Gallic acid, and methyl gallate	Darmawan <i>et al.</i> (2012)
3	<i>M. alnifolia</i>	Madagascar	Fruit	Alnifoliol, bonanniol A, diplacol, bonnanione A, and mphaeol A	20 Yoder <i>et al.</i> (2007)
			Leaves	schweinfurthin E, schweinfurthin F, schweinfurthin G, and schweinfurthin H	
4	<i>M. balansae</i>	Vietnam	Fruit	4'-methyl-8-prenyltaxifolin, 6,8-diprenyl-4'-methyl-naringenin, and 4'-deprenyl-4-methoxymap	Mai <i>et al.</i> (2020)
5	<i>M. barteri</i>	Cameroon	Stem bark	Macabarterin, scopoletin, 3-O-methylgallate, 4-O- β -D-xylopyranoside, Ellagic acid, 3-O-Methylgallate, Gallic acid, and Methyl gallate	Ngoumfo <i>et al.</i> (2008)
		Nigeria	Leaves	Benzamide, 2,2-Bis(4-nitrobenzyl)-1-phenylbutane-1,3-dione, 2-Acetophenone, 1,2,3-Trimethylbenzene, 1,2-Diacetoxypropane, 3-Hydroxy-1-phenyl-4-hexen-1-one, Methyl vinyl carbinol, 2-Isobutyl-3-methylfuran, acetyl tetrahydrofuran, isopropenyl dodecanoate, 2-hydroxy-2-methyl propanoic acid, undecane, Caryophyllene, aromadendrene, a-humulene, 3-Hydroxy-4(3H)-quinolinone, Caryophylla-2,6-dien-5-one, Germacrene B, Germacrene D, nerolidol, Spathulenol, tetradecane, b-elemene, N-acetylbenzamide, a-Linalool-2-penten-4-one, Caryophyllene oxide, N-phenylacrylamide, 2,12-Epoxy caryophyll-5-ene, 4-Methyl-1,3-oxathiolane, Methyl ethynyl carbinol, n-octadecane, Neophytadiene, Citronellyl propionate, trans-Phytol, 4-Methylpentanoic acid, n-Hexadecanoic acid, 2-Hexyl-2-decenal, Ethyl ester hexadecanoic acid, n-docosane, Citronellyl butyrate, cyclodecene, 2,5,8-Heptadecatrien-1-ol, dodecanedialdehyde, 9,12,15-Octadecatrien-1-ol, Ethyl Hept-6-enoate, 2-Keto-butyric-acid, Pentadecanal, 1-Aza-4-thiapentane, 2-Ethoxy carbonyloxy-1-phenyl-2-nonene, mappain, vedelianin, schweinfurthin G, macabartebene A, macabartebene B, and macabartebene C	
6	<i>M. bicolor</i>	Philippines	Leaves	(2S)-5,7-dihydroxy-4'-methoxy-8-(3''-methylbut-2''-enyl) flavanone, (2S)-5,7,4'-trihydroxy-8-(3''-methylbut-2''-enyl) flavanone, and (2S)-5,7,4'-trihydroxy-8-(3''-8''-dimethylocta-2'',7''-dienyl) flavanone	Versian <i>et al.</i> (2011)
7	<i>M. conglomerata</i>	Kenya	Leaves	Conglomeratin, macarangin, quercetin, 3,3',4'-trimethoxyellagic acid, and 3,3'	
			Stem bark	-dimethoxy ellagic acid 3-acetyllauritic acid, 2a-hydroxyauritic acid-3-p-hydroxybenzoate, and scopoletin	
8	<i>M. confiera</i>	Indonesia and China	Leaves	5-Hydroxy-4'-methoxy-2'',2''-dimethylpyrano-(7,8:6'',5'') flavanone, 5,4'-Dihydroxy-[2''-(1-hydroxy-1-methylethyl) dihydrofuran]-(7,8:5'',4'') flavanone, 5,7-Dihydroxy-4'-methoxy-8-(3-methylbut-2-enyl) flavanone, Lonchocarpol A, sophoraflavanone B, 5,7-Dihydroxy-4'-methoxy-8-(2-hydroxy-3-methylbut-3-enyl) flavanone, tomentosanol D, lupinifolinol, isolicoflavonol, (2S)-5,7,3'-trihydroxy-4'-methoxy-8-(3''-methylbut-2''-enyl) flavanone, (2S)-5,7,3',5'-tetrahydroxy-6,8-(3'',8''-dimethylbut-2'',7''-dienyl) flavanone, and 20-epibryonolic acid	20 Jang <i>et al.</i> (2004); Versian <i>et al.</i> (2011)
9	<i>M. constricta</i>	Malaysia	Leaves	Taraxerone, taraxerol, and β -amyrin	Salleh <i>et al.</i> (2017)
10	<i>M. deheiculata</i>	China	Fronds	Deheiculatin A, Deheiculatin B, Deheiculatin C, Deheiculatin D, Deheiculatin E, Deheiculatin F, Deheiculatin G, Deheiculatin H, Deheiculatin I, Deheiculatin J, Deheiculatin K, and Deheiculatin L	Qi <i>et al.</i> (2017)

Continued

No.	Plant species	Location	Plant part	Constituents	Sources
		Vietnam	Fruit	6-(8"-hydroxy-3",8"-trimethyl-oct-2"-enyl)-kaempferol, 3'-dehydroxy-solophenol C, denticulatain D, bonannione A, diplacol, macarangin, 3-O-methylmacarangin, 5,7,3',4'-tetrahydroxy-3-methoxy-6-geranylflavone, 3,5,7,3',4'-pentahydroxy-6-geranylflavonol, macarhizinoidin A, robipseudin A, denticulatain E, 8-dimethylallylisakuranetin, 4'-deprenylmappain, deheculatin H, poilaneic acid, derivative phaseic acid, taraxerone, and epitaraxerol	24 Le <i>et al.</i> (2021)
11	<i>M. denticulata</i>	China	Fronds	Denticulatains A, Denticulatains B, Denticulatain C, Denticulatain D, Denticulatain E, 5-hydroxy-2-(4-hydroxyphenyl)-7-methoxy-6-(3-methylbut-2-enyl) chroman-4-one, sophoraflavanone A, bonanniol A, diplacol, 5,7,3',4'-tetrahydroxy 6-geranylflavonol, 5,7,3',4'-tetrahydroxy-3-methoxy-6-geranylflavone, 3 β -hydroxy-7 α -e, thoxy-24 β -ethylcholest 5-ene, (24R)-6 β -hydroxy-24-ethylcholest-4-en-3-one, epitaraxerol, α -tocopherolquinone, boehmenan, denticalcones A, denticalcones C, denticalcones C, (2E)-1-(5,7-dihydroxy-2,2,6-trimethyl-2H-benzopyran-8-yl)-3-(4-methoxyphenyl)-2-propen-1-one, (2E)-1-(5,7-dihydroxy-2,2-dimethyl-2H-benzopyran-8-yl)-3-phenyl-2-propen-1-one, laxichalcone, macarangin, bonanniol A, bonannione A, and cirsitakoaside	Yang et al. (2015a); Yang et al. (2015b); Zhang et al. (2016); Di et al. (2020)
12	<i>M. gigantea</i>	Thailand	Leaves	Macdenticalcone, scopoletin, macarangin, 3-O-methyl-macarangin, denticulflavonol, sophoraflavanone B	Sutthivaiyakit (2002); Lei <i>et al.</i> (2016)
13	<i>M. gigantifolia</i>	Indonesia	Leaves	Macagigantin, Glycerin A, and Apigenin	Tanjung et al. (2009)
14	<i>M. hemsleyana</i>	Indonesia	Leaves	5,7-dihydroxycoumarin	Darmawan <i>et al.</i> (2012)
15	<i>M. heynei</i>	China	Stem bark	3 β -O-acetyl aleuritolic acid, canophyllol, β -Sitosterol, Stigmast-4-en-3-one, and Stigmast-4-en-3,6-dione	Wang <i>et al.</i> (2008)
16	<i>M. hosei</i>	Malaysia and Indonesia	Leaves	Laevifolins A, laevifolins B, Malayheyneiin A, Malayheyneiin B, Malayheyneiin C, Malayheyneiin D, laevifolin A, laevifolin B, macarubiginosin C, eserine, and Trolox	Kamarozaman <i>et al.</i> (2018, 2019)
17	<i>M. hurifolia</i>	Cameroon	Fruit	Lupenone, β -sitostenone, 5-hydroxy-7,4'-dimethoxyflavone, 5-hydroxy-6,7,4'-trimethoxyflavone, 4'-O-methyl-8-isoprenyl eriodictyol, and 6-isoprenyl eriodictyol	Salleh <i>et al.</i> (2017); Mariana <i>et al.</i> (2018a)
18	<i>M. hypoleuca</i>	Indonesia	Leaves	Macafolias A and macafolias B	Pagna et al. (2022)
			Sap	Dibromo chloropropane, levoglucosan, 2-metoksi-5-propeny, guanosine, Limonene, Lacton acid, trans-5,6-Epoksikan, d-Nerolidol, n-pentacosane, 1-Siklo leikosan, terakseron, squalene, Alpha-Beta-D-mannoside, and pentatriacontane	
			Stem	Terakseron and Alpha-Beta-D-mannoside	Gusmailina and Saputra (2021)
			Stem bark	2-propenal, isoeugenol, 1-(4-Hidroksi-3-MET), 4-Metil-2,5-Dimetoksibenzaldehid, Asam homovanillic, acetosyringone, and guaiaicol 3-(p-hidroksi-m-metoksifenil)-2-propenal, phenol, 3-allyl-2-metoksi-, Eugenol, 2-Propanon 1-(4-hidroksi), dimethoxybenzaldehyde, guanosine, Dekanon, and Sikloheksileikosan	

Continued

No.	Plant species	Location	Plant part	Constituents	Sources
19	<i>M. indica</i>	India and Vietnam	Leaves	Macaflavone I, Macaflavone II, macarindicins I, macarindicins II, macarindicins III, macarindicins IV, brousoflavonol F, vedelianin, schweinfurthin E, vitexin, 2"-rhamnosyl vitexin, isovitexin, (6R,7E,9R)-9-hydroxy-megastigman-4,7-dien-3-one-9-O-b-D-glucopyranoside, (6S,9R)-roseoside, (6S,9S)-roseoside, bridelionoside B, macarindicins D, macarindicins E, macarindicins F, macadenathin B, glyasperin A, kaempferol, quercetin, quercitrin, (+)-isolariciresinol, (-)-woonenoside XI, and (+)-lyoniresinol 4-O-β-D-glucopyranoside	Sultana and Ilyas (1986); Vu <i>et al.</i> (2021); Huonga <i>et al.</i> (2019)
20	<i>M. javanica</i>	China	Twigs	Macarindicins A, macarindicins B, macarindicins C, 6-Farnesyl-3',4',5,7-tetrahydroxyflavanone, isolicoflavonol, glyasperin A, brousoflavonol F, broussonol D, macarangin, and ellagic acid	Yang <i>et al.</i> (2015b)
21	<i>M. kurzii</i>	Indonesia	Leaves	Macajavanicin A, Macajavanicin B, Macajavanicin C, laevifolins A, and laevifolins B	Ilmiawati <i>et al.</i> (2015)
22	<i>M. lowii</i>	China	Twig	Kurzphenol A, kurzphenol B, kurzphenol C, 3,5-dihydroxy-4-(3-methyl-2-butenyl)bibenzyl, 6,8-diprenylgalangin, licoflavonol, icaritin, 3,4'-di-O-methyl-8-(3-methyl-2-butenyl) kaempferol, 5,7-dihydroxy-6,8-diprenylflavanone, isosakuranetin, 8-prenylnaringenin, glepidotin B, atractyloidin, acetylratryloidinol, blumenol A, scopoletin, and salicylic acid	Yang <i>et al.</i> (2014)
23	<i>M. magna</i>	Vietnam	Leaves	Macakurzin A, Macakurzin B, Macakurzin C, 5,7-dihydroxy-6-prenylflavanone, Glabranin, Izalpinin, Glepidotin A, 8-Prenylgalangin, Galangin, furanokurzin, Cis-3,5-dimethoxystilbene, Trans-3,5-dimethoxystilbene, and Trans-3,5-dimethoxy-2-prenylstilbene	Thanh <i>et al.</i> (2012)
24	<i>M. mappa</i>	Indonesia	Leaves	macalowiin, 4'-O-methyl-8-isoprenylnaringenin, 4'-O-methyl-5,7,4'-trihydroxyflavone	Agustina <i>et al.</i> (2012)
25	<i>M. monandra</i>	Indonesia	Leaves	scopoletin, apigenin, and gallic acid	Minarti <i>et al.</i> (2021)
26	<i>M. peltata</i>	Hawaii	Leaves	Mappain	Van Der Kaaden <i>et al.</i> (2001)
27	<i>M. pleiostemona</i>	Cameroon	Stem bark	Kolavenic acid and 2-Oxo-kolavenic acid	Salah <i>et al.</i> (2003)
28	<i>M. pruinosa</i>	India	Stem bark	β-Sitosterol, bergenin, 8,10-di-O-methylether, and Tri-O-methyl ether	Ramaiah <i>et al.</i> (1979)
29	<i>M. pruinosa</i>	Papua New Guinea	Leaves	Macaranga flavanone A, Macaranga flavanone B, euchrestalflavanone A, and Bonannione A	Schutz <i>et al.</i> (1995)
30	<i>M. recurvata</i>	Indonesia	Leaves	Macapruinosin B, Macapruinosin C, Papyriflavonol A, nymphaeol C, Macapruinosin D, Macapruinosin E, Macapruinosin F, and Macapruinosin A	Syah and Ghisalberti (2015)
31	<i>M. pustulata</i>	China	Twig	Deheiculatins M, deheiculatins N, deheiculatins O, deheiculatin I, deheiculatin C, deheiculatin B, poilaneic acid, and 20-hydroxy entcembrene	Syah and Ghisalberti (2015)
32	<i>M. recurvata</i>	Indonesia	Leaves	Macarecurvatin A, macarecurvatin B, Di isoprenyl aromadendrin, Glyasperin A, brousoflavonol F, flavestins K, flavestins B, flavestins G, 4-O-methyl-8-isoprenylnaringenin, and 8-isoprenyl-5,7-dihydroxyflavanone	Tanjung <i>et al.</i> (2012); Tjahjandarie <i>et al.</i> (2019)
33	<i>M. rhizinoides</i>	Indonesia	Leaves	Macarhizinoidin A, Macarhizinoidin B, and Methyl 4-isoprenyloxycinnamate	Tanjung <i>et al.</i> (2012)
34	<i>M. rubiginosa</i>	Indonesia	Leaves	Laevifolins A, laevifolins B, macarubiginosins A, macarubiginosins B, and macarubiginosins C	Tanjung <i>et al.</i> (2017)
35	<i>M. sampsonii</i>	China	Leaves	Macaranone A, macaranone B, macaranone C, and macaranone D	Li <i>et al.</i> (2009)
36	<i>M. schweinfurthii</i>	Cameroon	Leaves	Schweinfurthin A, schweinfurthin B, schweinfurthin C, schweinfurthin I, and schweinfurthin J	Beutler <i>et al.</i> (1998); Klausmeyer <i>et al.</i> (2010)

Continued

No.	Plant species	Location	Plant part	Constituents	Sources
35	<i>M. siamensis</i>	Thailand	Fronds	Macasiamenenes A, macasiamenenes B, macasiamenenes C, macasiamenenes D, macasiamenenes E, macasiamenenes F, macasiamenenes G, macasiamenenes H, macasiamenenes I, macasiamenenes J, macasiamenenes K, macasiamenenes L, macasiamenenes M, macasiamenenes N, macasiamenenes O, macasiamenenes P, macasiamenenes Q, macasiamenenes R, macasiamenenes S, macasiamenenes T, macasiamenenes U, macasiamenin A, macasiamenone A, macasiamenols A, and macasiamenols B	Pailee <i>et al.</i> (2015)
36	<i>M. sinensis</i>	China	Leaves	3-Desgalloylterchebin, macaranin A, Macaranin B, Macaranin C, Macarinin A, macarinin B, Macarinin C, tergallic acid bislactone, 3,6-O-(R)-hexahydroxydiphenoyl (HHDP)-D-glucose, tercatatin, mallorepanin, putranjivain A, and chlorogenic acid	Lin <i>et al.</i> (1990)
37	<i>M. tanarius</i>	Taiwan, Thailand, Japan, Indonesia, China, and Malaysia	Leaves	Tanariflavanone A, tanariflavanone B, (-)-nymphaeol C, tanariflavanone C, Tanariflavanone D, nymphaeol A, nymphaeol B, nymphaeol C, Macaflavanone A, macaflavanone B, macaflavanone C, macaflavanone D, Macaflavanone E, Macaflavanone F, Macaflavanone G, Mallotinic acid, Corilagin, macatannin A, Macatannin B, Chebulagic acid, 1(β)-O-galloylglucose, 4-O-galloylglucose, 6-O-galloylglucose, 2,3-di-O-galloylglucose, 1(β),2,6-tri-O-galloylglucose, 2,4,6-tri-O-galloylglucose, 1(β),3,4,6-tetra-O-galloylglucose, 1(β),2,3,4,6-penta-O-galloylglucose, 1(β),2,4,6-tetra-O-galloylglucose, 3-O-galloyl(-)-shikimic acid, 5-O-galloyl(-)-shikimic acid, 4-O-galloylquinic acid, 3,4-di-O-galloylquinic acid, 3,6-(S)-hexahydroxydiphenoyl (HHDP)-D-glucopyranose, punicafolin, furosin, terchebin, geraniin, mallotusinic acid, repandusinic acid A, 1,4-di-O-galloyl-α-D-glucopyranose, 3,4-di-O-galloyl-D-glucopyranose, galloylpunicafolin, galloylgeraniin, (-)-O-galloyl-3-O-brevifolincarboxyl-β-D-glucopyranose, macaranginin, tanarinin, (+)-pinoresinol 4-O-[6''-O-galloyl]-β-D-glucopyranoside, macarangioside E, Macarangioside F, mallophenol B, Macarangioside A, Macarangioside B, Macarangioside C, Macarangioside D, laurosides E, Methyl brevifolin carboxylate, Macarangiosides A, Macarangiosides B, macarangiosides C, macarangiosides D, macarangiosides E, macarangiosides F, macarangiosides A, macarangiosides B, macarangiosides C, macarangiosides D, macarangiosides E, and macarangiosides F	Tseng <i>et al.</i> (2001); Phormmart <i>et al.</i> (2005); Kawakami <i>et al.</i> (2008); Gunawan-Puteri and Kawabata (2010); Lin <i>et al.</i> (1990); Matsunami <i>et al.</i> (2009); Yoshimura <i>et al.</i> (2017); Matsunami and Otsuka (2018); Marlina <i>et al.</i> (2018b); Rawa <i>et al.</i> (2022)
		Hong Kong	Stem bark	macarangenol, kolavenol, blumenol A, blumenol B, annuonone E, and lathyrene	Hui <i>et al.</i> (1971)
		Vietnam and Japan	Fruit	Vedelianin, schweinfurthins E, schweinfurthins F, schweinfurthins G, schweinfurthins K, schweinfurthins L, schweinfurthins M, schweinfurthins N, schweinfurthins O, schweinfurthins P, schweinfurthins Q, mappain, methylmappain, nymphaeol B, isonymphaeol B, nymphaeol A, 3'-geranyl-naringenin, nymphaeol C, propolin C, propolin D, propolin F, propolin G, propolin H	Peresse <i>et al.</i> (2017); Huong <i>et al.</i> (2020); Kumazawa <i>et al.</i> (2014); Nam <i>et al.</i> (2021); Natsume <i>et al.</i> (2021); Lee <i>et al.</i> (2019)
38	<i>M. trichocarpa</i>	Indonesia	Leaves	Macatrichocarpin A, macatrichocarpin B, macatrichocarpin C, macatrichocarpin D, 4'-O-methylmacagigantatin, macatrichocarpin F, macatrichocarpin G, Macatrichocarpin H, licoflavanon, flavokawain C, and helichrysetin	Syah and Ghisalberti (2010); Tanjung <i>et al.</i> (2018)
		Indonesia and Vietnam	Leaves	3,7,3',4'-tetramethylquercetin, 3,7,3'-trimethylquercetin, 3,7-dimethylquercetin, 2'-hydroxy-macarangaflavanone A, 4',7-hydroxy-8-methylflavan, scopoletin, 3-epi-taraxerol, and Taraxerol	Jang <i>et al.</i> (2004); Dinh <i>et al.</i> (2006)
39	<i>M. triloba</i>	Malaysia	Flowers	6-Prenyl-3'-methoxy-eriodictyol, nymphaeol B, nymphaeol C, 6-Farnesyl-3',4',5,7-tetrahydroxyflavanone, 4,5-dihydro-5'-α-hydroxy-4'-α-methoxy-6a,12a-dehydro-α-toxicaro, (+)-Clovan-2β,9α-diol, Ferulic acid, abscisic acid, 1β,6α-dihydroxy-4(15)-eudesmene, 3β-hydroxy-24-ethylcholest-5-en-7-one, loliolide, tanarifuranonol, 5,7-dihydroxycoumarin, and malaysianone A	Zakaria <i>et al.</i> (2010, 2012)
40	<i>M. vedeliana</i>	New Caledonia	Leaves	Macarangin	Hnawia <i>et al.</i> (1990)

lipids (Leopoldini *et al.*, 2011). Antioxidants inhibit lipid oxidation by reacting with lipids and peroxy radicals and converting them into more stable nonradical products (Lin *et al.*, 2018; Petruk *et al.*, 2018). Furthermore, antioxidants also help in overcoming inflammation.

Ogundajo and Ashafa (2019) reported that the methanol extract of a popular traditional medicinal plant in west Africa, *M. barteri*, demonstrated a high antiradical effect against 1,1-diphenyl-2-picrylhydrazyl (DPPH) and nitric oxide (NO) with values of 0.47 and 1.68 mg/ml, respectively. *M. tanarius* fruit extracts exhibited strong antioxidant activity due to the presence of prenylflavonoids in the seed, pericarp, glandular trichome, and leaf (Kumazawa *et al.*, 2014; Kumazawa *et al.*, 2008). Chien *et al.* (2022) also reported that *M. tanarius* new and mature fruit extracts exhibited stronger free radical-scavenging activity and possessed lower IC50 than Taiwanese green propolis extract. Ethyl acetate extract of *M. triloba* leaves from Central Kalimantan, Indonesia, exhibited a strong ability to neutralize free radicals DPPH (Ardany *et al.*, 2018). Kamarozaman *et al.* (2019) reported that new dihydrostilbenes isolated from *M. heynei* showed some antioxidant activity.

Antimicrobial activity

In cosmetic goods, antimicrobial compounds are used to both inhibit the growth of unfavorable microorganisms in skincare products and increase the shelf life of products (Nowak *et al.*, 2021). The skin is the largest organ in the body, and it is exposed to the environment, making it an ideal location for bacteria, viruses, or fungi to grow. In order to protect consumers and stabilize shelf life, especially against potentially harmful bacteria, the formulation of cosmetic products (water based vs. oil based) has considered multifunctional antimicrobial ingredients (Hoang *et al.*, 2021). This is in line with rising consumer awareness and demands for clean beauty and avoiding artificial ingredients, especially preservatives. It is realized that the use of conventional antimicrobial agents, which are usually loaded with preservatives, has disturbed the balance of skin microflora, so the skin is vulnerable to exposure to harmful microorganisms (Nowak *et al.*, 2021).

Methanolic extract of *M. barteri* leaves grown in Nigeria has been reported to inhibit *Pseudomonas aeruginosa*, *Enterococci faecalis*, and *Cryptococcus neoformans* (>80% inhibition (Ogbole *et al.*, 2018). Purayil *et al.* (2019) used a combination solvent, chloroform and water (1:10), to extract from the leaves of *M. peltata*, which resulted in a clearance zone of the well diffusion method against *Staphylococcus aureus*. Panda *et al.* (2017) tested acetone, water, and ethanol solvents to extract from *M. peltata* leaves, which had antimicrobial activity against *Escherichia coli*, *S. aureus*, and *Candida albicans*. As described in another report, both the leaves and stem bark of *M. peltata* also had positive activity against four bacterial strains: *E. coli*, *P. aeruginosa*, *Bacillus subtilis*, and *S. aureus* (Verma *et al.*, 2009).

Bradacs *et al.* (2010) reported that the inner bark extract of *Macaranga dioica*, commonly used as traditional medicine in the South Pacific Archipelago of Vanuatu, exhibited moderate activity against *C. albicans*. Ogundajo *et al.* (2017) extracted hexane, ethyl acetate, and methanol from *M. barteri* leaves. They found that methanolic extract had the best antibacterial activity against

S. aureus, *Bacillus pumilus*, *Streptococcus faecalis*, *Listeria* sp., *P. aeruginosa*, *Plesiomonas shigelloides*, *Aeromonas hydrophila*, *Shigella sonnei*, *Salmonella typhi*, *Salmonella typhimurium*, *E. coli*, *Proteus vulgaris*, *Enterobacter faecalis*, and *Klebsiella pneumoniae*. The authors also discovered that a hexane extract of the plant was effective against fungi such as *Candida neoformans*, *Trichophyton mucoides*, and *Candida albicans*. Lee *et al.* (2019) isolated five propolins from *M. tanarius*. Among them, propolin D showed the highest inhibition of biofilm formation by strains of *S. aureus*, *Staphylococcus epidermidis*, and *C. albicans*, with MICs of 10–50 µg/ml. Isolated prenylated kaempferol and conglomeratin, from *M. conglomerata*, reportedly showed significant permeation of *P. aeruginosa* (MIC = 7.8 mg/ml) and moderate activity against *S. aureus*, *E. coli*, and *Klebsiella pneumoniae* (MIC = 62.5 mg/ml) (Hashim *et al.*, 2022).

Tyrosinase inhibitory effect

Antimelanogenesis activity is one of the most important criteria for determining the suitability of plant compounds for use in skincare agents. It was previously reported by Arung *et al.* (2019) that they isolated glyasperin A, a prenylated flavonoid isolated from *M. pruinosa* that greatly inhibited melanin in B16 melanoma. Another report from Mazlan *et al.* (2013) demonstrated that they had extracted *M. denticulata*, *M. pruinosa*, and *M. gigantea* using methanol to assess their tyrosinase inhibitory activity. Among them, methanolic extract of *M. denticulata* bark showed the highest inhibition (68.7%). Methanol extract from the leaves and stem bark of *M. hurifolia* has been tested to show their application in tyrosinase inhibition, in which they found appropriate activity with values of 159.42 [mg Kojic acid equivalent (KAE)/g] and 160.95 (mg KAE/g), respectively (Sadeer *et al.*, 2019). Lim *et al.* (2009) screened the bioactivity of methanolic extracts of *M. gigantea*, *M. pruinosa*, *M. tanarius*, and *M. triloba*, which resulted in the best tyrosinase inhibition activity obtained by *M. pruinosa* among all the *Macaranga* tested. KAE values and quercetin equivalent were 6.8 and 20.7 mg/g, respectively.

Limitations of natural ingredient in cosmetics

Macaranga extract contains antioxidant, anti-inflammatory, and antimicrobial compounds. Some of these compounds have potential as topical cosmetic preparations. However, it is acknowledged that not all of these natural ingredients are safe because they may be associated with carcinogenic, mutagenic, and reprotoxic chemicals (Hoang *et al.*, 2021). As a result, their application as skincare products that protect the integrity of cosmetics and the skin at the same time must be considered wisely. Furthermore, despite their promising effects, validation with clinical results is required. Due to a lack of clinical data and the limited relevance of data, some of the practical significance of the effects mentioned has not been adequately proven. However, these data provide interesting points for dermatologists to consider in the application of natural ingredients in relation to topical therapies (Hoang *et al.*, 2021).

Although uncommon, skin contact with cosmetics containing plant extracts can result in allergic responses, contact dermatitis, erythema multiforme, and xanthomatous reactions (Hoang *et al.*, 2021). It is possible, due to the lack of separation techniques, that many plant extracts have not been investigated

for their compounds (Zorzi *et al.*, 2016). Antioxidants may also have a variety of adverse effects, including acute toxicity, skin and eye irritation, skin sensitization, and photosensitization (Hoang *et al.*, 2021; Mujtaba *et al.*, 2021). In light of this, the application of natural ingredients requires that they are applied in safe concentrations; for instance, the application of essential oils is only recommended at 0.1–0.8%, or diluting with carrier oils (Guzmán and Lucia, 2021; Vostinaru *et al.*, 2020). The use of some natural extracts, such as polyphenols, in cosmetics is constrained due to their low stability and sensitivity to heat and light (Hoang *et al.*, 2021). Due to this condition, some cosmetic products that contain furocoumarins, compounds that are also present in *Macaranga* extract, should only be applied at night. Sun exposure has phototoxicity effects, causing skin damage.

Natural antioxidants are susceptible to deterioration, and their bioavailability is limited by low absorption. The use of active phytochemicals with the application of nanotechnology in cosmetics has drawn a lot of attention since it can boost absorption by the skin (Salvioni *et al.*, 2021). Various types of nanoemulsions, nanoparticles, liposomes, niosomes, and dendrimers have influenced the formulation of cosmetic products (Musthaba *et al.*, 2009). According to test results, the application of nano formulations can minimize hematological toxic effects, boost bioavailability, and lessen other side effects, such as alopecia, nausea, vomiting, diarrhea, exhaustion, and skin rashes (Hoang *et al.*, 2021; Saklani and Kutty, 2008; Salvioni *et al.*, 2021; Zorzi *et al.*, 2016). Recently, major concerns regarding its safety have been raised, and much more exploration is needed to determine its efficacy in delivering active ingredients into the skin. New regulations established by the European Union have passed amendments in its (new regulations) cosmetics directory for safer nanocosmetics to enter the market, safeguarding the beauty and health of consumers (Hoang *et al.*, 2021; Musthaba *et al.*, 2009). Natural antioxidants are more expensive than synthetic ones in cosmetics, despite the fact that these molecules are safer than synthetic antioxidants.

Other potential uses of *Macaranga* plants

Hendra *et al.* (2017) reported that the extract of *M. tanarius* exhibited positive antihyperlipidemic and hepatoprotective effects. Some active compounds isolated from ethyl acetate extract of *M. tanarius* leaves, i.e., mallotinic acid, corilagin, chebulagic acid, macatannins A and B, can be developed for diabetes treatment agents (Gunawan-Puteri and Kawabata, 2010). Another *Macaranga* species, *M. hurifolia* extract from Nigeria, showed potential for anti-inflammatory and antidiabetic effects (Segun *et al.*, 2019b). Ehile *et al.* (2018) reported that the aqueous extract of *M. barkeri* exhibited a gastric antiulcer effect at doses ranging from 62.5 to 500 mg/kg b.w. This extract has been reported to be safe for oral medicine due to major toxic anthropometric and hematological effects not being present in tested rats, despite the fact that biochemical and histological studies are needed to ensure safe use of the extract (Ehile *et al.*, 2018). Furthermore, as reported earlier, *Macaranga* spp. contains rich prenylated flavonoids (Shahinozzaman *et al.*, 2021). These exhibited several potential effects: antiviral activity (Feng *et al.*, 2010), antiallergic activity (Quan *et al.*, 2008), and larvicidal activity against the fourth-instar larvae of *Aedes albopictus* and *Culex pipiens quinquefasciatus* (Niu *et al.*, 2010). Muhaimin *et al.* (2019) reported that ethanolic extract of *M. gigantea* exhibited

antiplasmodial activity and can be developed as an alternative agent for controlling malaria. The study also confirmed that *M. denticulata* leaf extract has antinociceptive activity. At doses of 200 and 400 mg/kg, it demonstrated dose-dependent and statistically significant antinociceptive activity in acetic acid and formalin tests. Furthermore, among six major *M. denticulata* compounds, *Macaranga* had the best fitness score of 5.81 with the COX-1 enzyme (Hasanat *et al.*, 2017).

CONCLUSIONS

A comprehensive review of the potential of *Macaranga* plants as skincare cosmetics ingredients has been conducted. *Macaranga* plants contain bioactive compounds which have anti-inflammatory, antioxidant, antibacterial, and tyrosinase inhibitory activity. Therefore, *Macaranga* plants have the potential to be used as skincare cosmetic ingredients.

AUTHORS' CONTRIBUTIONS

Conceptualization is done by E.T.A. and M.T.H.; methodology is carried out by E.T.A., M.T.H., and A.S.P.; investigation is done by E.T.A., I.W.K., H.K., R.A., and E.R.; writing original draft preparation is conducted by M.T.H., I.W.K., H.K., R.A., E.R., Y.Y., W.S., S.P., R.R., D.T., M.I., H.A.A., W.F., A.A., C.R.K., N.I.W.A., Y.K., and E.T.A.; writing review and editing is done by E.T.A., W.F., D.T., and H.A.A.; visualization is carried out by M.T.H., D.T., W.D., H.A.A., and E.T.A.; project administration is done by E.T.A. and W.F.; funding acquisition is done by E.T.A. All authors have read and agreed to the published version of the manuscript.

FUNDING

This research received funding from the Deputy of Research and Innovation, National Research and Innovation Agency (BRIN) for “Pusat Kolaboratif Riset Kosmetik Berteknologi Nano Berbasis Bio-massa” in the Fiscal Year 2022 (grant number: 398/IL/FR/3/2022).

CONFLICTS OF INTEREST

The authors have no relevant financial or nonfinancial interests to disclose.

ETHICAL APPROVAL

This study does not involve experiments on animals or human subjects.

DATA AVAILABILITY

All data generated and analyzed are included within this review article.

PUBLISHER'S NOTE

This journal remains neutral with regard to jurisdictional claims in published institutional affiliations.

REFERENCES

- Agustina W, Juliawaty L, Hakim E, Syah Y. Flavonoids from *Macaranga lowii*. ITB J Sci, 2012; 44.
- Amberg N, Fogarassy C. Green consumer behavior in the cosmetics market. Resources, 2019; 8:137.
- Amir Rawa MS, Nurul Azman NA, Mohamad S, Nogawa T, Wahab HA. In vitro and in silico anti-acetylcholinesterase activity from *Macaranga tanarius* and *syzygium jambos*. Molecules, 2022; 27(9):2648.

- Amirta R, Candra KP. Comparative characterization of *Macaranga* species collected from secondary forests in East Kalimantan for biorefinery of unutilized fast growing wood. *Biodivers J Biol Divers*, 2016; 17:116–23.
- Ardany SD, Mulia DS, Rosawanti P. Antioxidant activity of ethyl acetate fraction of *Macaranga triloba* leaves from central Kalimantan. *Asian J Pharm Clin Res*, 2018; 11(3):39–42.
- Arung ET, Sinamabela JR, Rosamah E, Kusuma IW, Kuspradini H, Alam AE, Amen Y, Tanaka H, Satria D, Shimizu K, Ishikawa H. Antioxidant and antimelanogenesis activities of glyasperin A from *Macaranga pruinosa* leaves. *Nat Prod Commun*, 2019; 14(7):1934578X19867192.
- Asante-Kwatia E, Jibira Y, Mensah A, Osei-Sarfoh D. *Macaranga barteri* stem bark extract exerts anti-inflammatory and anti-hyperalgesia activity in murine models. *Discover Phytomed*, 2019; 6(3):130–7.
- Attiq A, Jalil J, Husain K, Ahmad W. Raging the war against inflammation with natural products. *Front Pharmacol*, 2018; 9:976.
- Beutler JA, Shoemaker RH, Johnson T, Boyd MR. Cytotoxic geranyl stilbenes from *Macaranga schweinfurthii*. *J Nat Prod*, 1998; 61(12):1509–12.
- Bradacs G, Maes L, Heilmann J. In vitro cytotoxic, antiprotazoal and antimicrobial activities of medicinal plants from Vanuatu. *Phytother Res*, 2010; 24(6):800–9.
- Chien YH, Yu YH, Ye SR, Chen YW. Antibacterial and antioxidant activity of the fruit of *Macaranga tanarius*, the plant origin of Taiwanese green propolis. *Antioxidants*, 2022; 11(7):1242.
- Darmawan A, Kosela S, Kardono LBS, Syah Y. Scopoletin a coumarin derivative compound isolated from *Macaranga gigantifolia*. *Merr J Appl Pharm Sci*, 2012; 2:17–25.
- Di Q, Zhu H, Pu D, Zhao X, Li X, Ma X, Xiao W, Chen W. The natural compound Cirsitikaoside enhances antiviral innate responses against vesicular stomatitis virus in vitro and in vivo. *Int Immunopharmacol*, 2020; 86:106783.
- Dinh V, Zhang HP, Duc NM, Tuu NV, Qin GW. A new geranyl flavanone from *Macaranga triloba*. *J Asian Nat Prod Res*, 2006; 8(1–2):155–8.
- Ehile E, Goze N, Kouakou KL, Yapo A, Ehile E. Acute toxicity and gastric anti-ulcer activity of an aqueous extract of the leaves of *Macaranga barteri* Mill. Arg (*Euphorbiaceae*) on rat models. *J Med Plants Res*, 2018; 12:96105.
- Farage MA, Miller KW, Elsner P, Maibach HI. Intrinsic and extrinsic factors in skin ageing: a review. *Int J Cosmetic Sci*, 2008; 30(2):87–95.
- Feng T, Wang RR, Cai XH, Zheng YT, Luo XD. Anti-human immunodeficiency virus-1 constituents of the bark of *Poncirus trifoliata*. *Chem Pharm Bull*, 2010; 58(7):971–5.
- Gunawan-Puteri MDPT, Kawabata J. Novel α -glucosidase inhibitors from *Macaranga tanarius* leaves. *Food Chem*, 2010; 123(2):384–9.
- Gusmailina R, Saputra N. Identification on chemical organic compounds of *Macaranga hypoleuca* by using Py-GCMS. *IOP Conf Ser Earth Environ Sci*, 2021; 914:012065.
- Guzmán E, Lucia A. Essential oils and their individual components in cosmetic products. *Cosmetics*, 2021; 8(4):114.
- Hasanat A, Chowdhury TA, Kabir MSH, Chowdhury MS, Chy MNU, Barua J, Chakrabarty N, Paul A. Antinociceptive activity of *Macaranga denticulata* Muell. Arg. (Family: *Euphorbiaceae*): in vivo and in silico studies. *Medicines (Basel)*, 2017; 4(4):88.
- Hashim I, Onyari JM, Omosa LK, Maru SM, Nchiozem-Ngnitedem VA, Karpoomath R. Conglomeratin: a new antibacterial flavonol derivative from *Macaranga conglomerata* Brenan (*Euphorbiaceae*). *Nat Prod Res*, 2022; 36:1–9.
- He H, Li A, Li S, Tang J, Li L, Xiong L. Natural components in sunscreens: topical formulations with sun protection factor (SPF). *Biomed Pharmacother*, 2021; 134:111161.
- Hendra P, Jamil O, Maharani D, Suhadi M, Putri C, Fenty, Julianus J. Antihyperlipidemic and hepatoprotective studies on leaves of *Macaranga tanarius*. *Asian J Pharmaceutical Clin Res*, 2017; 10:239–41.
- Hnawia E, Thoison O, Guéritte-Voegelein F, Bourret D, Sévenet T. A geranyl substituted flavonol from *Macaranga vedeliana*. *Phytochemistry*, 1990; 29(7):2367–8.
- Hoang HT, Moon JY, Lee, YC. Natural antioxidants from plant extracts in skincare cosmetics: recent applications, challenges and perspectives. *Cosmetics*, 2021; 8(4):106.
- Hui WH, Ng KK, Fukamiya N, Koreeda M, Nakanishi K. Isolation and structure of macarangelol, a diterpene ketol from *Macaranga tanarius*. *Phytochemistry*, 1971; 10(7):1617–20.
- Huong DTM, Linh NT, Van TTT, Litaudon M, Roussi F, Van Nam V, Van Cuong P. Stilbenes from *Macaranga tanarius* (Euphorbiaceae) growing in Vietnam. *Vietnam J Chem*, 2020; 58(3):338–42.
- Huanga DTM, Vu LTN, Anh L, Cuc NT, Nhiem NX, Tai BH, Van Kiem P, Litaudon M, Dang Thach T, Van Minh C, Pham VC. Cytotoxic prenylated flavonoids from the leaves of *Macaranga indica*. *Phytochem Lett*, 2019; 34:39–42.
- Ilmiawati A, Hakim EH, Syah YM. Prenylated 9,10-dihydrophenanthrenes from *Macaranga javanica*. *Zeitschrift für Naturforschung B*, 2015; 70(9):659–63.
- Jang DS, Cuendet M, Pawlus AD, Kardono LB, Kawanishi K, Farnsworth NR, Fong HH, Pezzuto JM, Kinghorn AD. Potential cancer chemopreventive constituents of the leaves of *Macaranga triloba*. *Phytochemistry*, 2004; 65(3):345–50.
- Kamarozaman A, Ahmat N, Rahman NF, Yen KH. Prenylated dihydrostilbenes from *Macaranga heynei* (Euphorbiaceae). *Malaysian J Anal Sci*, 2018; 22:258–63.
- Kamarozaman AS, Ahmat N, Isa SNM, Hafiz ZZ, Adenan MI, Yusof MIM, Azmin NFN, Latip J. New dihydrostilbenes from *Macaranga heynei* IM. Johnson, biological activities and structure-activity relationship. *Phytochem Lett*, 2019; 30:174–80.
- Kawakami S, Harinantenaina L, Matsunami K, Otsuka H, Shirazate T, Takeda Y. Macaflavanones A-G, prenylated flavanones from the leaves of *Macaranga tanarius*. *J Nat Prod*, 2008; 71(11):1872–6.
- Keeyari Purayil S, Chew A, Paulraj P, Pattammadath S, Mohamed J, Selvarani J, P R, Ravibalan T, Petchi I, Samrot A. Evaluation of antioxidant and antimicrobial activity of some plants collected from Malaysia. *J Pure Appl Microbiol*, 2019; 13:2363–73.
- Klausmeyer P, Van QN, Jato J, McCloud TG, Beutler JA. Schweinfurthins I and J from *Macaranga schweinfurthii*. *J Nat Prod*, 2010; 73(3):479–481.
- Kouassi MC, Grisel M, Gore E. Multifunctional active ingredient-based delivery systems for skincare formulations: a review. *Colloids Surf B Biointerfaces*, 2022:112676.
- Kumazawa S, Murase M, Momose N, Fukumoto S. Analysis of antioxidant prenylflavonoids in different parts of *Macaranga tanarius*, the plant origin of Okinawan propolis. *Asian Pac J Trop Med*, 2014; 7(1):16–20.
- Kumazawa S, Nakamura J, Murase M, Miyagawa M, Ahn MR, Fukumoto S. Plant origin of Okinawan propolis: honeybee behavior observation and phytochemical analysis. *Naturwissenschaften*, 2008; 95(8):781–6.
- Le TNV, Truong BN, Le TP, Litaudon M, Tran DT, Chau VM, Mai HDT, Pham VC. Cytotoxic phenolic compounds isolated from the fruits of *Macaranga denticulata*. *Nat Prod Res*, 2021; 35(11): 861–1868.
- Lee JH, Kim YG, Khadke SK, Yamano A, Woo JT, Lee, J. Antimicrobial and antibiofilm activities of prenylated flavanones from *Macaranga tanarius*. *Phytomedicine*, 2019; 63:153033.
- Lei C, Zhang LB, Yang J, Gao LX, Li JY, Li J, Hou AJ. Macdenticalcone, a unique polycyclic dimeric chalcone from *Macaranga denticulata*. *Tetrahedron Letters*, 2016; 57(49):5475–8.
- Leláková V, Béraud-Dufour S, Hošek J, Šmejkal K, Prachywarakorn V, Pailee P, Widmann C, Václavík J, Coppola T, Mazella J, Blondeau N, Heurteaux C. Therapeutic potential of prenylated stilbenoid macasiamenene F through its anti-inflammatory and cytoprotective effects on LPS-challenged monocytes and microglia. *J Ethnopharmacol*, 2020; 263:113147.
- Leopoldini M, Russo N, Toscano M. The molecular basis of working mechanism of natural polyphenolic antioxidants. *Food Chem*, 2011; 125(2):288–306.
- Li X, Xu L, Wu P, Xie H, Huang Z, Ye W, Wei X. Prenylflavonols from the leaves of *Macaranga sampsonii*. *Chem Pharm Bull (Tokyo)*, 2009; 57(5):495–8.

- Lim TY, Lim YY, Yule CM. Evaluation of antioxidant, antibacterial and anti-tyrosinase activities of four *Macaranga* species. *Food Chem*, 2009; 114(2):594–99.
- Lin JH, Ishimatsu M, Tanaka T, Nonaka GI, Nishioka I. Tannins related compounds. XCVI. Structures of macaranins and macarinins, hydrolyzable tannins possessing macaranoyl and tergalloyl ester groups, from the leaves of *Macaranga sinensis* (BAILL.) MUELL. ARG. *Chem Pharm Bull*, 1990; 38:1844–51.
- Lin TK, Zhong L, Santiago JL. Anti-inflammatory and skin barrier repair effects of topical application of some plant oils. *Int J Mol Sci*, 2018; 19(1):70.
- Magadula J. Phytochemistry and pharmacology of the genus *Macaranga*: a review. *J Med Pl Res*, 2014; 8(12):489–503.
- Mai HDT, Toan TP, Huu GT, Le TN, Oanh VTK, Hang NTM, Thu HT, Chau VM, Litaudon M, Pham VC. New flavonoid and stilbene derivatives from the fruits of *Macaranga balansae*. *Nat Prod Res*, 2020; 34(19):277–86.
- Marliana E, Astuti W, Kosala K, Hairani R, Tjahjandarie T, Tanjung M. Chemical composition and anticancer activity of *Macaranga hosei* Leav. *Asian J Chem*, 2018a; 30:795–8.
- Marliana E, Hairani R, Tjahjandarie TS, Tanjung M. Plasmodial activity of flavonoids from *Macaranga tanarius* leaves. *IOP Conf Ser Earth Environ Sci*, 2018b; 144:012011.
- Matsunami K, Otsuka H. Okinawan subtropical plants as a promising resource for novel chemical treasury. *Chem Pharm Bull (Tokyo)*, 2018; 66(5):519–26.
- Matsunami K, Otsuka H, Kondo K, Shinzato T, Kawahara M, Yamaguchi K, Takeda Y. Absolute configuration of (+)-pinoreosin, (+)-O-galloyl- β -D-glucopyranoside, macarangiosides E, and F isolated from the leaves of *Macaranga tanarius*. *Phytochemistry*, 2009; 70(10):1277–85.
- Mazlan NA, Mediani A, Abas F, Ahmad S, Shaari K, Khamis S, Lajis NH. Antioxidant, antityrosinase, anticholinesterase, and nitric oxide inhibition activities of three Malaysian *Macaranga* species. *SciWorld J*, 2013; 2013:312741.
- Milito A, Castellano I, Damiani E. From Sea to Skin: is there a future for natural photoprotectants? *Mar Drugs*, 2021; 19(7):379.
- Minarti Cahyana AH, Darmawan A. Isolation, characterization and antidiabetic activity of phenolic compounds isolated from *Macaranga magna* turn. *Rasayan J Chem*, 2021; 14(4):2420–7.
- Morais T, Cotas J, Pacheco D, Pereira L. Seaweeds compounds: an eco-sustainable source of cosmetic ingredients? *Cosmetics*, 2021; 8.
- Muhaimin M, Yusnaidar Y, Wilda S, Madyawati L, Riski Dwimalida P, Adita U, Anis Yohana C, Andreas Yoga A, Josephine Giabeth S. Antiplasmodial activity of ethanolic extract of *Macaranga gigantea* L. and its major constituent. *Pharmacognosy J*, 2019; 11(6):11.
- Mujtaba SF, Masih AP, Alqasmi I, Alsulimani A, Khan FH, Alkhou S. Oxidative-stress-induced cellular toxicity and glycoxidation of molecules by cosmetic products under sunlight exposure. *Antioxidants*, 2021; 10(7):1508.
- Musthafa S, Ahmad S, Ahuja A, Ali J, Baboota S. Nano approaches to enhance pharmacokinetic and pharmacodynamic activity of plant origin drugs. *Curr Nanoscience*, 2009; 5(3):344–52.
- Nam SH, Yamano A, Kim JA, Lim J, Baek SH, Kim JE, Kwon S, Saito Y, Teruya T, Choi SY, Kim YK, Bae YC, Shin HI, Woo JT, Park EK. Prenylflavonoids isolated from *Macaranga tanarius* stimulate ontoblast differentiation of human dental pulp stem cells and tooth root formation via the mitogen-activated protein kinase and protein kinase B pathways. *Endod J*, 2021; 54(7):1142–54.
- Natsume N, Yonezawa T, Saito Y, Woo JT, Teruya T. Prenylflavonoids from fruit of *Macaranga tanarius* promote glucose uptake via AMPK activation in L6 myotubes. *J Nat Med*, 2021; 75(4):813–23.
- Ngoumfo RM, Ngounou GE, Tchamadeu CV, Qadi MI, Mbazoa B, Begum A, Ngninzeke FN, Lontsi D, Choudhary MI. Inhibitory effect of macabarterin, a polyoxygenated ellagitannin from *Macaranga barkeri*, on human neutrophil respiratory burst activity. *J Nat Prod*, 2008; 71(11):1906–10.
- Niu HM, Zeng DQ, Long CL, Peng YH, Wang YH, Luo JF, Wang HS, Shi YN, Tang, GH, Zhao FW. Clrodane diterpenoids and prenylated flavonoids from *Dodonaea viscosa*. *J Asian Nat Prod Res*, 2010; 12(1):7–14.
- Nowak K, Jabłońska E, Ratajczak-Wrona W. Controversy around parabens: alternative strategies for preservative use in cosmetics and personal care products. *Environment Res*, 2021; 198:110488.
- Ogbole OO, Segun PA, Fasinu PS. Antimicrobial and antiprotozoal activities of twenty-four Nigerian medicinal plant extracts. *S Afr J Bot*, 2018; 117:240–6.
- Ogundajo A, Okeleye B, Ashafa AO. Chemical constituents, in vitro antimicrobial and cytotoxic potentials of the extracts from *Macaranga barkeri* Mull-Arg. *Asian Pac J Trop Biomed*, 2017; 7(7):654–9.
- Ogundajo AL, Tom Ashafa AO. Chemical profiling, antioxidant and carbohydrate-metabolising enzymes inhibitory potential of fractions from the leaves of *Macaranga barkeri* Mull-Arg. *Trans R Soc S Afr*, 2019; 74(1):27–39.
- Pagna JIM, Awazi T, Mbarga PE, Mbekou IMK, Mkounga Fotie J, Frese M, Fabrice FB, Lenta BN, Sewald N, Nkengfack EA. Antibacterial flavonoids from the fruits of *Macaranga hurifolia*. *J Asian Nat Prod Res*, 2022: 1–1.
- Pailee P, Sangpetsiripan S, Mahidol C, Ruchirawat S, Phiyawarakorn V. Cytotoxic and cancer chemopreventive properties of prenylated stilbenoids from *Macaranga siamensis*. *Tetrahedron*, 2015; 71(34):5562–71.
- Panda SK, Padhi L, Leyssen P, Liu M, Neyts J, Luyten W. Antimicrobial, anthelmintic, and antiviral activity of plants traditionally used for treating infectious disease in the simlipal biosphere reserve, Odisha, India. *Front Pharmacol*, 2017; 8:658.
- Péresse T, Jézéquel G, Allard PM, Pham VC, Huang DTM, Blanchard F, Bignon J, Lévaïque H, Wolfender JL, Litaudon M, Roussi F. Cytotoxic prenylated stilbenes isolated from *Macaranga tanarius*. *J Nat Prod*, 2017; 80(10):2684–91.
- Petruk G, Del Giudice R, Rigano MM, Monti DM. Antioxidants from plants protect against skin photoaging. *Oxid Med Cell Longev*, 2018; 2018:145415.
- Phommart S, Sutthivaiyakit P, Chimnoi N, Ruchirawat S, Sutthivaiyakit S. Constituents of the leaves of *Macaranga tanarius*. *J Nat Prod*, 2005; 18(6):927–30.
- Qi WY, Shen Y, Wu Y, Leng Y, Gao K, Yue JM. Deheiculatins A-L, 20-oxygenated cembranoids from *Macaranga deheiculata*. *Phytochemistry*, 2017; 136:101–7.
- Quan W, Lee H, Noh C, Um ABH, Oak MH, Kim KM. Anti-allergic prenylated flavonoids from the roots of *sophora flavescens*. *Planta Med*, 2008; 74:168–70.
- Ramaiah PA, Row LR, Reddy DS, Anjaneyulu ASR, Ward RS, Pelter A. Isolation and characterisation of bergenin derivatives from *Macaranga peltata*. *J Chem Soc Perkin Trans 1*, 1979; (0):2313–6.
- Rees J. The genetics of sun sensitivity in humans. *Am. J. Hum. Genet.*, 2005; 75(5):739–51.
- Sadeer, N, Llorent-Martínez E, Bene K, Mahomoodally F, Mollica A, Ibrahim S, Stefanucci A, Ruiz Riaguas A, Fernández de Ardoval ML, Zengin G. Chemical profiling, antioxidant, enzyme inhibitory and molecular modelling studies on the leaves and stem bark extracts of three African medicinal plants. *J Pharm Biomed Anal*, 2019; 174:19–33.
- Saklani A, Kutty SK. Plant-derived compounds in clinical trials. *Drug Disc Today*, 2008; 13(3-4):161–71.
- Salah MA, Bedir E, Toyang NJ, Khan IA, Harries MD, Wedge JE. Antifungal clrodane diterpenes from *Macaranga monandra* (L) Muell. et Arg. (Euphorbiaceae). *J Agric Food Chem*, 2003; 51(26):7607–10.
- Salleh WMNH, Razak N, Farediah A. Phytochemicals and biological activities of *Macaranga hosei* and *Macaranga constricta* (Euphorbiaceae). *Marmara Pharm J*, 2017; 21:881–8.
- Salvioni L, Morelli L, Ochoa E, Labra M, Fiandra L, Palugan L, Prosperi D, Colombo M. The emerging role of nanotechnology in skincare. *Adv Colloid Interface Sci*, 2021; 293:102437.

- Schütz BA, Wright AD, Rali T, Sticher O. Prenylated flavanones from leaves of *Macaranga pleiostemona*. *Phytochemistry*, 1995; 40(4):1273–7.
- Segun P, Gbadebo M, Adebawale M, Olufolabo K, Fred-Jaiyesimi A. Investigation of the anti-inflammatory and hypoglycaemic effects of *Macaranga hurifolia* beille (euphorbiaceae) extract on wistar albino rats. *ACTA Pharm Sci*, 2019a; 57:93.
- Segun PA, Ogbale OO, Akinleye TE, Faleye TOC, Adeniji AJ. In vitro anti-enteroviral activity of stilbenoids isolated from the leaves of *Macaranga barteri*. *Nat Prod Res*, 2021; 35(11):1909–13.
- Segun PA, Ogbale OO, Ismail FMD, Nahar L, Evans AR, Ajaiyeoba EO, Sarker SD. Bioassay-guided isolation and structure elucidation of cytotoxic stilbenes and flavonols from the leaves of *Macaranga barteri*. *Fitoterapia*, 2019b; 134:151–7.
- Shahinozzaman M, Obanda DN, Tawata S. Chemical composition and pharmacological properties of *Macaranga*-type Pacific propolis: a review. *Phytother Res*, 2021; 35(1):07–22.
- Sultana S, Ilyas M. Chromenoflavones from *Macaranga indica*. *Phytochemistry*, 1986; 25(4):953–4.
- Susanto D, Kusuma R, Amirta R. Nutrient distribution on soil and aboveground biomass of *Macaranga gigantea* five years after planting. *Asian J Forest*, 2018; 2:12–9.
- Sutthivaiyakit S, Unganont S, Sutthivaiyakit P, Suksamrarn A. Diterpenylated and prenylated flavonoids from *Macaranga denticulata*. *Tetrahedron*, 2002; 58(18):3619–22.
- Syah YM, Ghisalberti EL. Flavanone derivatives from *Macaranga tanarius*. *Biochemical Systematics and Ecology*, 2015; 62:151–4.
- Syah YM, Ghisalberti EL. Phenolic derivatives with an irregular sesquiterpenyl side chain from *Macaranga pruinosa*. *Nat Prod Commun*, 2010; 5(2):219–2.
- Tanjung M, Hakim EH, Elfahmi Latip J, Syah YM. Dihydroflavonol and flavonol derivatives from *Macaranga recurvata*. *Nat Prod Commun*, 2012; 7(10):1309–10.
- Tanjung M, Hakim EH, Mujahidin D, Hanafi M, Syah YM. Macagigantins, a farnesylated flavonol from *Macaranga gigantea*. *J Asian Nat Prod Res*, 2009; 11(11):929–32.
- Tanjung M, Hakim EH, Syah YM. Prenylated dihydrostilbenes from *Macaranga rubiginosa*. *Chem Nat Compd*, 2017; 53(2):215–8.
- Tanjung M, Juliawaty LD, Hakim EH, Syah YM. Flavonoid and stilbene derivatives from *Macaranga trichocarpa*. *Fitoterapia*, 2018; 126:74–7.
- Tjahjandarie T, Tanjung M, Saputri R, Nadar P, Aldin M, EvaMarliana, Permadi A, Flavestin K. An isoprenylated stilbene from the leaves of *Macaranga recurvata* gage. *Nat Prod Sci*, 2019; 25:244.
- Trinh Thi Thanh V, Doan Thi Mai H, Pham VC, Litaudon M, Dumontet V, Guéritte F, Nguyen VH, Chau VM. Acetylcholinesterase inhibitors from the leaves of *Macaranga kurzii*. *J Nat Prod*, 2012; 75(11):2012–5.
- Tseng MH, Chou CH, Chen YM, Kuo YH. Allelopathic prenylflavanones from the fallen leaves of *Macaranga tanarius*. *J Nat Prod*, 2001; 64(6):827–8.
- Van Der Kaaden JE, Hemscheidt TK, Mooberry SL. Mappain, a new cytotoxic prenylated stilbene from *Macaranga mappa*. *J Nat Prod*, 2001; 64(1):103–5.
- Verma M, Raj V, Hariharapura R, Rao J, Udupa N. Screening of plant *Macaranga peltata* for its antioxidant, antimicrobial and cytotoxicity activity. *Int Conference Biomed Pharm Eng Singapore*, 2009.
- Versiani M, Ratnayake R, Henrich C, Bates S, McMahon J, Gustafson K. flavonoids from eight tropical plant species that inhibit the multidrug resistance transporter ABCG2. *J Nat Prod*, 2011; 74:262–6.
- Vostinaru O, Heghes S, Filip L. Essential oils—bioactive compounds. *New perspectives and applications*. IntechOpen, Paris, France, 2020.
- Vu LTN, Anh LT, Cuc NT, Nhiem NX, Tai BH, Van Kiem P, Litaudon M, Thach TD, Van Minh C, Mai HDT, Van Cuong P. Prenylated flavonoids and other constituents from *Macaranga indica*. *Nat Prod Res*, 2021; 35(13):2123–0.
- Wang TS, Liu BJ, Hua SY, Li TL, Chen GY. Study on the chemical constituents of liposoluble steroidal and triterpenoid compounds from the stem and bark of *Macaranga hemsleyana*. *Zhong Yao Cai*, 2008; 31(3):372–4.
- Yang DS, Li ZL, Wang X, Yan H, Yang YP, Luo HR, Liu KC, Xiao WL, Li XL. Denticulatains A and B: unique stilbene-diterpene heterodimers from *Macaranga denticulata*. *RSC Advances*, 2015a; 5(18):13886–90.
- Yang DS, Peng WB, Yang YP, Liu KC Li, X.-L, Xiao WL. Cytotoxic prenylated flavonoids from *Macaranga indica*. *Fitoterapia*, 2015b; 103.
- Yang DS, Wang SM, Peng WB, Yang YP, Liu KC, Li XL, Xiao WL. Minor prenylated flavonoids from the twigs of *Macaranga adenantha* and their cytotoxic activity. *Nat Prod Bioprospect*, 2015c; 5(2):105–9.
- Yang DS, Wei JG, Peng WB, Wang SM, Sun C, Yang YP, Liu KC, Li XL. Cytotoxic prenylated bibenzyls and flavonoids from *Macaranga kurzii*. *Fitoterapia*, 2014; 99:261–6.
- Yoder BJ, Cao S, Norris A, Miller JS, Ratovoson F, Razafitsalama J, Andriantsiferana R, Rasamison VE, Kingston DG. Antiproliferative prenylated stilbenes and flavonoids from *Macaranga alnifolia* from the Madagascar rainforest. *J Nat Prod*, 2007; 70(3):342–6.
- Yoshimura K, Hosoya T, Fujinami M, Ohta T, Kumazawa S. Nymphaeol-C a prenylflavonoid from *Macaranga tanarius*, suppresses the expression of fibroblast growth factor 18. *Phytomedicine*, 2017; 36:238–42.
- Zahidin NS, Saidin S, Zulkifli RM, Muhamad II, Ya'akob H, Nur H. A review of *Acalypha indica* L. (Euphorbiaceae) as traditional medicinal plant and its therapeutic potential. *J Ethnopharmacol*, 2017; 207:146–73.
- Zakaria I, Ahmat N, Ahmad R, Mohd Jaafar F, Ghani N, Ghani S, Khamis. Flavanones from the Flower of *Macaranga triloba*. *World Appl Sci J*, 2010; 9:1003–7.
- Zakaria I, Ahmat N, Jaafar FM, Widyawaruyanti A. Flavonoids with antiplasmodial and cytotoxic activities of *Macaranga triloba*. *Fitoterapia*, 2012; 83(5):968–72.
- Zhang LB, Lei C, Gao LX, Li JY, Li J, Hou AJ. Isoprenylated flavonoids with PTP1B inhibition from *Macaranga denticulata*. *Nat Prod Bioprospect*, 2016; 6(1):25–30.
- Zapanta BR, Achondo MJMM, Raganas AFM, Fritzie A, Delima AGD, Mantiquilla JA, Salvaña FRP. Species richness of trees in disturbed habitats within a protected area and its implications for conservation: the case of Mt. Apo natural park, Mindanao Island, Philippines. *Biodivers J Biol Divers*, 2019; 20: 2081–91.
- Zixi W, Bainian S, Jin P, Deng P, Chen J, Sun F. A new species of *Macaranga* from the middle miocene of Fujian, China and its significance. *Hist Biol*, 2016; 29: 1–2.
- Zorzi GK, Caregnato F, Moreira JCF, Teixeira HF, Carvalho ELS. Antioxidant effect of nanoemulsions containing extract of *Achyrocline satureioides* (Lam) D.C.—Asteraceae. *AAPS Pharm Sci Tech*, 2016; 17(4):844–50.

How to cite this article:

Rosamah E, R, Haqiqi MT, Putri AS, Kuspradini H, Kusuma IW, Amirta Yuliansyah Y, Suwinarti W, Paramita S, Ramadhan R, Tarmadi D, Ismayati M, Ariyanta HA, Fatrisari W, Aswandi A, Kholibrina CR, Azelee NIW, Kim YU, Arung ET. The potential of *Macaranga* plants as skincare cosmetic ingredients: A review. *J Appl Pharm Sci*, 2023. <https://doi.org/10.7324/JAPS.2023.77745>

The potential of Macaranga plants as skincare cosmetic ingredients: A review

ORIGINALITY REPORT

11 %
SIMILARITY INDEX

9 %
INTERNET SOURCES

9 %
PUBLICATIONS

2 %
STUDENT PAPERS

PRIMARY SOURCES

1 erepository.uonbi.ac.ke 2 %
Internet Source

2 pharmacology.ejournals.ca 1 %
Internet Source

3 academictree.org 1 %
Internet Source

4 pubmed.ncbi.nlm.nih.gov 1 %
Internet Source

5 www.eurekaselect.com 1 %
Internet Source

6 www.wjgnet.com 1 %
Internet Source

7 www.coursehero.com 1 %
Internet Source

8 www.jetir.org <1 %
Internet Source

9 pub.uni-bielefeld.de <1 %
Internet Source

10

iopscience.iop.org

Internet Source

<1 %

11

link.springer.com

Internet Source

<1 %

12

Rodrigue Keumoe, Jean Garba Koffi, Darline Dize, Patrick Valère Tsouh Fokou et al. "Identification of 3,3'-O-dimethylellagic acid and apigenin as the main antiplasmodial constituents of *Endodesmia calophylloides* Benth and *Hymenostegia afzelii* (Oliver.) Harms", *BMC Complementary Medicine and Therapies*, 2021

Publication

<1 %

13

Louisa Gomes, Pedro Monteiro, João Cotas, Ana M. M. Gonçalves, Chantal Fernandes, Teresa Gonçalves, Leonel Pereira. "Seaweeds' pigments and phenolic compounds with antimicrobial potential", *Biomolecular Concepts*, 2022

Publication

<1 %

14

ia601607.us.archive.org

Internet Source

<1 %

15

www.jourlib.org

Internet Source

<1 %

16

synapse.koreamed.org

Internet Source

<1 %

17 Neneng Windayani, Lia Juliawaty, Euis Hakim, Komar Ruslan, Yana Syah. "An Acetylated Triterpene Glycoside and Lignans from *Phyllanthus myrtifolius* and their Antibacterial Activities", *The Natural Products Journal*, 2015
Publication

18 www.degruyter.com
Internet Source

19 Veronika Leláková, Sophie Béraud-Dufour, Jan Hošek, Karel Šmejkal et al. "Therapeutic potential of prenylated stilbenoid macasiamenene F through its anti-inflammatory and cytoprotective effects on LPS-challenged monocytes and microglia", *Journal of Ethnopharmacology*, 2020
Publication

20 Ishtiaque Ahammad, Md. Rafiul Islam Sarker, Akib Mahmud Khan, Sohedul Islam, Mahmud Hossain. "Virtual Screening to Identify Novel Inhibitors of Pan ERBB Family of Proteins from Natural Products with Known Anti-tumorigenic Properties", *International Journal of Peptide Research and Therapeutics*, 2019
Publication

21 Suporn Phommart, Pakawadee Sutthivaiyakit, Nitirat Chimnoi, Somsak Ruchirawat, Somyote Sutthivaiyakit. " Constituents of the Leaves of ", *Journal of Natural Products*, 2005

22

E Marliana, R Ruga, R Hairani, T S Tjahjandarie, M Tanjung. " Antioxidant activity of flavonoid constituents from the leaves of ", *Journal of Physics: Conference Series*, 2019

Publication

23

Submitted to Higher Education Commission Pakistan

Student Paper

24

Submitted to Imperial College of Science, Technology and Medicine

Student Paper

25

Arindam Gangopadhyay, Syamantak Chakraborty, Shyamal K. Jash, Dilip Gorai. "Cytotoxicity of natural flavones and flavonols against different cancer cells", *Journal of the Iranian Chemical Society*, 2022

Publication

26

Fatma Afifi-Yazar, Violet Kasabri, Rana Abu-Dahab. " Medicinal Plants from Jordan in the Treatment of Diabetes: Traditional Uses vs. and Evaluations – Part 2 ", *Planta Medica*, 2011

Publication

27

Maliheh Entezari, Danial Hashemi, Afshin Taheriazam, Amirhossein Zabolian et al. "AMPK signaling in diabetes mellitus, insulin resistance and diabetic complications: A pre-

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

clinical and clinical investigation", Biomedicine & Pharmacotherapy, 2022

Publication

28

Peter A. Segun, Omonike O. Ogbole, Toluwanimi E. Akinleye, Temitope O. C. Faleye, Adekunle J. Adeniji. " anti-enteroviral activity of stilbenoids isolated from the leaves of ", Natural Product Research, 2019

Publication

<1 %

29

Teodora D. Balangcod, Kryssa D. Balangcod. "Chapter 242-1 Macaranga magna Turrrill Euphorbiaceae", Springer Science and Business Media LLC, 2021

Publication

<1 %

30

cebp.aacrjournals.org

Internet Source

<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On