

# AJAS

*by* Rudy Agung

---

**Submission date:** 21-Mar-2023 11:34AM (UTC-0700)

**Submission ID:** 2042866188

**File name:** Manuscript\_turnitin.docx (91.69K)

**Word count:** 4679

**Character count:** 24585



## Effect of Water Hyacinth Leaf Flour (*Eichhornia crassipes*) Fermented by *Aspergillus niger* on the Growth, Survival Rate and Blood Profile of Sangkuriang Catfish (*Clarias gariepinus*)

### ABSTRACT

Catfish sangkuriang (*Clarias gariepinus*) is one of the aquaculture commodities with great potential to be developed in Indonesia to support the demand for feed. Efforts to reduce dependence on imported feed raw materials for fish feed is continued. Water hyacinth (*Eichhornia crassipes*) leaf which contains phytochemical compounds such as flavonoids, tannins, steroids and saponins is an alternatif raw material for fish pellet component. This study aimed to determine the addition of fermented water hyacinth leaf meal in fish feed on the growth, survival rate, and blood profile of sangkuriang catfish. In total 150 fish (Average initial weight  $5.45 \pm 0.02$  g) were randomly divided into triplicates of five groups, containing 10 fishes per group. Group one is a control group, P1 (10% of non fermented water hyacinth leaf meal), while P2, P3 and P4 were treatment groups of fish fed with 10, 15, and 20% fermented water hyacinth leaf meal by *Aspergillus niger*. At the end of the 42 days of treatment, growth, survival rate, and blood profile of all fish were determined. The results showed that the addition of 10% water hyacinth leaf meal fermented by *Aspergillus niger* had an effect on the growth of final weight, body weight gain and spesific growth rate ( $p < 0.05$ ) in comparison to all groups. However, the addition of any concentration of water hyacinth leaf meal did not affect on the feed conversion ratio and survival rate value. Further, the addition of water hyacinth leaf meal had a significant effect ( $p < 0.05$ ) on the erythrocytes, leukocytes and hemoglobin of fish, but did not show a significant effect on the values of hematocrit and platelets.

**Keywords:** *Clarias gariepinus*, *Eichhornia crassipes*; Growth; Survival Rate; Blood Profile

### INTRODUCTION

Catfish (*Clarias gariepinus*) is one of the freshwater fish commodities that has a great potential to be cultured. This fish has been cultivated commercially and is growing rapidly on limited land Production (Sorongan *et al*, 2019). In addition, catfish can be kept with high stocking densities in line with current market demand (Simanjuntak *et al*, 2020). In addition, catfish production has increased by an average of 9.23% per year (KKP, 2021).

The culture of catfish in the community has encouraged aquaculture business actors to continue to strive for maximum production (Audila & Zulfahmi, 2021). However, this is inversely proporsional to the availability of catfish which is still not matched by the availability of feed that affects production costs (Prasekti & Irfan, 2021). Efforts to reduce dependence on imports of feed raw materials are to find substitutes for alternative materials

whose quality is cheap, easy to obtain, and has a high nutritional content (Amin *et al*, 2020). One alternative that can be used as a raw material for feed is water hyacinth (*Eichhornia crassipes*).

Water hyacinth is an aquatic plant commonly referred to as a weed and is able to reproduce very quickly. Water hyacinth is easy to adapt to its environment and is able to compete with other organisms. The rapid growth of water hyacinth can reach 3% per day. Their rapid growth can greatly disrupt activities in the waters and can result in silting and narrowing of rivers, lakes, and swamps which causes reduced oxygen content, decreased nutrients and water evaporation (Kurniawan *et al*, 2020).

The use of water hyacinth in feed ingredients has several weaknesses, including high hemicellulose content and protein content that is difficult to digest which it can interfere

with digestibility and palatability or level of preference (Rahayu, 2020). The nutritional content of water hyacinth in the form of dry matter has a crude protein content of 6.31% and crude fiber 26.61% (Putra *et al.*, 2020). Therefore, it is necessary to use an easy fermentation process using *Aspergillus niger* (Setiawan, 2013).

*Aspergillus niger* is one of the microbial species that does not produce mycotoxins compared to other microbes (Maryanty *et al.*, 2010). In addition, *Aspergillus niger* can produce enzymes that can help the digestive process such as cellulase, amylase, protease, phytase and mananase which make it easier for livestock feed to be digested (Lestari *et al.*, 2021).

Past study by Raudah *et al.*, (2018) used *Aspergillus niger* as a fermenter on the leaf plant of Lamtoro Gung (*Leucaena leucocephala*) in the feed of Siamese catfish (*Pangasius hypophthalmus*). Meanwhile, Pane *et al.*, (2018) added water hyacinth leaf meal in the manufacture of feed with concentration of 10% which is the optimum concentration on the growth and survival or survival of sangkuriang catfish fry (*Clarias gariepinus*). Further, Syaraswati *et al.*, (2022) performed an addition of 10% water hyacinth leaf meal fermented by *Aspergillus niger* in the feed which contains flavonoid compounds, resulting high survival rates for the growth of Nile tilapia (*Oreochromis niloticus*) fry. In addition, the effects of ginger powder (*Zingiber officinale*) 3.75 g/kg feed containing flavonoid compounds affected on the blood profile of catfish (*Pangasius sp*) (Fajriyani *et al.*, 2017).

Though several studies have been performed to determine the effects of the water hyacinth meal on the feed of fish, limited study on the effects of fermented water hyacinth meal on the sangkuriang catfish. Present research was conducted to determine the effects of fermented water hyacinth leaf meal in feed on the growth, survival, and blood profile of sangkuriang catfish.

## MATERIALS AND METHODS

### Fish preparation

In total of 150 sangkuriang catfish with an initial average weight of  $5.45 \pm 0.02$  g, 45 days old was provided from local farmer at Samarinda. All fish was fed a commercial pellet feed (Hi-Pro-Vite 781-2) Made in Indonesia, during acclimatization (7 Days).

After acclimatization, all fish were randomly stocked into triplicates of five tarpaulin ponds, containing 10 fish per pond. Ponds were labelled as (K) control/commercial pellet group, (P1) water hyacinth leaf meal without fermentation + Pellet 90%; (P2) fermented water hyacinth leaf meal 10% + Pellet 90%; (P3) fermented water hyacinth leaf meal 15% + Pellet 85%, and (P4) fermented water hyacinth leaf meal 20% + Pellet 80% (Pane *et al.*, 2018).

### Water Hyacinth Leaf Meal Fermentation

Water hyacinth leaves were cut, washed and finely chopped. The water hyacinth leaves were dried under the sun for 4 days. After that, the water hyacinth leaves were milled and sifted. The water hyacinth leaf meal was ready to be fermented.

The water hyacinth leaf meal was added with water at ratio 2:1.5 (Raudah *et al.*, 2018). The mixture was steamed for 15 minutes to kill pathogens that can interfere fermentation process. Further, the mixture was cooled and inoculated with *Aspergillus niger* powder at a dose of 2%, put into a plastic that had been perforated to obtain aerobic conditions. The fermentation process was done for 72 h at room temperature. After 72 h, the fermented product was re-steamed for 15 minutes to stop the fermentation process. The fermented product was dried and kept into the plastic bag and store in 4 C until its used. The fermented water hyacinth leaf meal was also measured for fiber analysis. Based on our results, the fiber content

of water hyacinth leaf meal before fermentation was  $38.05 \pm 0.43\%$ , while after fermentation was  $7.78 \pm 1.87\%$

The dry fermented water hyacinth leaf meal was mixed with commercial feed with the ratio as describe previously. After added with 1000 mL of distilled water, all the raw materials were mixed, then the dough was re-pellet using pelletizer. The pellet was then dried using an oven at  $60^\circ\text{C}$  or for 17 hours.

### Fish Rearing

Fish were reared for 42 days, fed with commercial feed and fermented water hyacinth leaf meal two times per day at satiation rate. ponds siphoning was carried 2 times a week to remove faeces and food residue and maintained water quality.

### Fish Parameters Measurement

Fish biomass determination was carried out every 2 weeks. Measurement of fish biomass was performed by taking the total number of fish in the pond for each replication using a net. The fish were weighed using a digital scale. Fish was also determined for Body Weight Gain (BWG), Specific growth rate (SGR), Feed Conversion ratio (FCR), and Survival Rate (SR) using equation below:

$$\text{BWG} = \text{Wt} - \text{Wo} \quad (\text{Suci et al., 2020})$$

4  
 BWG = Body weight gain (g)  
 Wt = Fish weight at the end of the research (g)  
 Wo = Fish weight at the beginning of the research (g)

$$\text{SGR} = \frac{[\ln \text{Wt} - \ln \text{Wo}]}{t} \times 100\% \quad (\text{Setiawati et al., 2013})$$

7  
 SGR = Spesific growth rate (%)  
 Wt = Fish weight at the end of the research (g)

Wo = Fish weight at the beginning of the research (g)  
 t = Experiment duration (days)

$$\text{FCR} = \frac{F}{\text{Wt} - \text{Wo}} \quad (\text{Anis et al., 2019})$$

FCR = Feed Conversion Ratio (g)  
 F = Total of Feed Consumed (g)  
 Wt = Fish Weight at the end of the research (g)  
 Wo = Fish weight at the beginning of the research (g)

$$\text{SR}(\%) = \frac{\text{Nt}}{\text{No}} \times 100\% \quad (\text{Tamba et al., 2020})$$

SR = Survival Rate (%)  
 Nt = Total Fish at the end of the research  
 No = Total Fish at the beginning of the research

### Fish Blood Sampling

Fish blood sampling was done by taking out the fish blood through the dorsal aorta, placing a syringe between the anus and anal fin. The collected blood was put into a tube containing 0.5 mL of 0.2% EDTA.

### Hematology Analysis

Hematology profile analysis of collected fish blood was carried out to determine the health condition of fish using a hematology analyzer. The blood profiles data were White Blood Cell Count (WBC), Red Blood Cell Count (RBC), platelets/platelets (Plt), hemoglobin (Hb) and hematocrit (HCT). Observation of fish blood profile was performed at the end of the final day of the study.

### Water Quality Measurement

Water quality measurements such as temperature, pH, DO (*Dissolved Oxygen*), TAN

(Total Ammonia Nitrogen), and nitrite were checked once a week.

### Data Analysis

The result data such as BWG, FCR, SR, and blood profiles were analyzed using SPSS statistic 22 and presented as Mean  $\pm$  standard error. Meanwhile water quality data, namely pH, DO (Dissolved Oxygen), TAN (Total Ammonia Nitrogen), and Nitrite were measured once a week. To determine significantly difference, Duncan Multiple Range Test with a significance level of  $p < 0.05$  was applied.

### Results

#### Growth and Survival

It can be seen that the growth of catfish with 10% fermented water hyacinth leaf meal (P2) has a significant effect ( $p < 0.05$ ) on the final weight, BWG, SGR of fish in comparison to

other treatment groups. However, all fish groups including control, showed no significant effect on the FCR and SR (Table 1).

#### Blood Profile Parameters

The blood profile of catfish showed that the number of erythrocytes, hematocrit and platelets were not significantly different between the control group and the unfermented treatment group ( $p > 0.05$ ). However, the number of leukocytes and hemoglobin in the blood of catfish fed fermented water hyacinth leaf meal groups had a significant effect in comparison to fish both the control group and the unfermented treatment group (Table 2).

#### Water Quality Sangkuriang Catfish

The water quality in the rearing pond was still within the range below the water quality standard for catfish and is categorized as suitable for use as a rearing place (Table 3).

**Table 1.** Growth of Sangkuriang Catfish (*Clarias gariepinus*) fed various levels of fermented and unfermented Water hyacinth (*Eichhornia crassipes*) leaf meal for 42 days.

Parameters	K	P1	P2	P3	P4
Initial Weight (g)	53.09 $\pm$ 1.80 <sup>a</sup>	51.37 $\pm$ 0.07 <sup>a</sup>	50.37 $\pm$ 2.07 <sup>a</sup>	51.56 $\pm$ 2.64 <sup>a</sup>	51.14 $\pm$ 2.33 <sup>a</sup>
Final Weight (g)	138.28 $\pm$ 6.49 <sup>b</sup>	94.20 $\pm$ 8.10 <sup>a</sup>	167.01 $\pm$ 2.85 <sup>c</sup>	126.74 $\pm$ 6.31 <sup>b</sup>	122.13 $\pm$ 7.20 <sup>b</sup>
BWG (g)	85.62 $\pm$ 4.71 <sup>b</sup>	42.83 $\pm$ 8.17 <sup>a</sup>	116.63 $\pm$ 1.25 <sup>c</sup>	75.17 $\pm$ 5.20 <sup>b</sup>	70.98 $\pm$ 5.50 <sup>b</sup>
SGR (%)	10.57 $\pm$ 0.13 <sup>a</sup>	8.86 $\pm$ 0.42 <sup>b</sup>	11.33 $\pm$ 0.26 <sup>c</sup>	10.27 $\pm$ 0.16 <sup>a</sup>	10.13 $\pm$ 0.16 <sup>a</sup>
FCR	3.06 $\pm$ 0.27 <sup>a</sup>	4.09 $\pm$ 0.34 <sup>a</sup>	2.44 $\pm$ 0.58 <sup>a</sup>	3.01 $\pm$ 0.45 <sup>a</sup>	3.23 $\pm$ 0.06 <sup>a</sup>
SR (%)	80.00 $\pm$ 0.00 <sup>a</sup>	80.00 $\pm$ 5.77 <sup>a</sup>	86.66 $\pm$ 3.33 <sup>a</sup>	86.66 $\pm$ 3.33 <sup>a</sup>	86.66 $\pm$ 8.81 <sup>a</sup>

**Note:** K (Control), P1 (Treatment with the addition of 10% water hyacinth leaf meal without fermentation), P2-P4 (Treatment with the addition of 10; 15, and 20% of fermented water hyacinth leaf meal. All data shows as mean  $\pm$  standard error. The mean followed by different superscript letters (a, b, c) in the same row indicates a significant difference ( $p < 0.05$ ). BWG: Body Weight Gain; SGR: Specific Growth Rate; FCR: Feed Conversion Ratio, SR: Survival Rate

**Table 2.** Blood Profile of Sangkuriang Catfish (*Clarias gariepinus*) fed various levels of fermented and unfermented Water hyacinth (*Eichhornia crassipes*) leaf meal for 42 days.

Parameters	Control	P1	P2	P3	P4
Leukosit ( $10^3/\mu\text{L}$ )	0.95 $\pm$ 0.03 <sup>a</sup>	0.42 $\pm$ 0.04 <sup>a</sup>	5.69 $\pm$ 0.41 <sup>b</sup>	7.20 $\pm$ 2.64 <sup>b</sup>	6.35 $\pm$ 0.41 <sup>b</sup>
Eritrosit ( $10^6/\mu\text{L}$ )	0.25 $\pm$ 0.19 <sup>abc</sup>	0.04 $\pm$ 0.04 <sup>b</sup>	0.76 $\pm$ 0.09 <sup>c</sup>	0.55 $\pm$ 0.03 <sup>a</sup>	0.76 $\pm$ 0.09 <sup>c</sup>
Hemoglobin (g/dL)	3.96 $\pm$ 0.23 <sup>a</sup>	2.86 $\pm$ 0.03 <sup>b</sup>	2.46 $\pm$ 0.03 <sup>c</sup>	2.30 $\pm$ 0.17 <sup>c</sup>	2.46 $\pm$ 0.03 <sup>c</sup>
Hematokrit (%)	3.73 $\pm$ 3.08 <sup>a</sup>	0.46 $\pm$ 0.03 <sup>a</sup>	10.76 $\pm$ 1.02 <sup>a</sup>	8.20 $\pm$ 1.03 <sup>a</sup>	10.56 $\pm$ 1.02 <sup>a</sup>
Platelet ( $10^3/\mu\text{L}$ )	21.00 $\pm$ 15.01 <sup>a</sup>	20.66 $\pm$ 0.33 <sup>a</sup>	4.33 $\pm$ 0.33 <sup>a</sup>	7.00 $\pm$ 1.73 <sup>a</sup>	5.33 $\pm$ 1.45 <sup>a</sup>

**Note:** K (Control), P1 (Treatment with the addition of 10% water hyacinth leaf meal without fermentation), P2-P4 (Treatment with the addition of 10; 15, and 20% of fermented water hyacinth leaf meal. All data shows as mean  $\pm$  standard error.



error. The mean followed by different superscript letters (a, b, c) in the same row indicates a significant difference ( $p < 0.05$ ). BWG: Body Weight Gain; SGR: Specific Growth Rate ; FCR: Feed Conversion Ratio, SR: Survival Rate

**Table 3.** Water Quality parameters during the study

Parameters	Control	P1	P2	P3	P4	*BSN
Temperature (°C)	27.41	27.38	27.41	27.38	27.38	25-30
pH	7.57	7.50	7.57	7.50	7.50	6.5-8.5
DO (mg/L)	6.00	6.90	6.00	6.90	6.90	>4.0
TAN (mg/L)	0.1	0.1	0.1	0.1	0.1	<1
Nitrite (mg/L)	0.01	0.01	0 - 0.01	0.01	0.01	<1

**Note:** K (Control), P1 (Treatment with the addition of 10% water hyacinth leaf meal without fermentation), P2-P4 (Treatment with the addition of 10; 15, and 20% of fermented water hyacinth leaf meal. \*(BSN) National Standardization Agency Quality Standard, 2014. Tolerance limit for the maintenance of sangkuriang catfish.

## Discussion

### Growth Parameters

Present study showed that, the highest BWG (116.63 g) and SGR (11.33%) were found in fish fed 10% of fermented water hyacinth leaf meal groups. The addition of water hyacinth leaf meal with a concentration of 10% in the feed was able to improve the growth performance of catfish. This is in line with the past study performed by Wijaya *et al.*, (2015) stated that water hyacinth leaves contain several plant phytochemical compounds such as flavonoids, tannins, steroids and saponins that are beneficial in growth. According to Zhai & Liu (2013), fish fed flavonoid compounds can increase the average final weight and BWG of the fish.

Further, Indariyanti & Rakhmawati (2013) revealed that *Aspergillus niger* produces cellulase and xylanase enzymes which will remodel cellulose into glucose and eventually become glucose. This is in accordance with the research of Ikhwanuddin *et al.*, (2018) fermented bran by *Aspergillus niger* as a feed raw material can increase the value of protein digestibility and growth in tilapia (*Oreochromis niloticus*) compared to unfermented feed.

Meanwhile, present study found that there was no significant difference ( $p > 0.05$ ) on the FCR between control and all treatment groups. Widyastuti *et al.*, (2010) stated that the smaller

the FCR value, the higher the quality of feed and vice versa, while Marzuqi & Anjusary (2013) added that the digestibility of fish itself is influenced by several factors, namely the age of the fish, the nutritional level of the feed and the water temperature.

Furthermore, the results of the SR during the 42-day study showed no significantly different and SR was in the range between 80.00 and 86.66%. This range is supported by previous study by Mulyani *et al.*, (2014) stating that the survival rate of more than 50% is classified as good, 30-50% is classified as moderate and less than 30% is classified as not good. The current study is also in concomitant with Syaraswati *et al.*, (2022) and Kurniawan *et al.*, (2022) study, revealed that the use of fermented or unfermented water hyacinth leaf meal did not affect on the survival rate of nilem (*Osteochilus basselti*) fish and baung fish (*Hemibagrus nemurus*).

### Blood Profiles

The number of erythrocyte of fish fed fermented water hyacinth leaf meal was higher than the control group and the unfermented treatment group. The increase in the number of erythrocytes was due to the flavonoid compounds which is found in water hyacinth leaves that might be able to improve the health of catfish. These results are in line with the

research of Zissalwa *et al.*, (2020), which tested the effect of mangrove leaf extract on the erythrocyte profile parameters of jambal siam (*Pangasius hypophthalmus*) fish. It is found that feeding of mangrove leaf extract containing flavonoid compounds can increase the number of erythrocytes. Putri *et al.*, (2013) also stated that the number of erythrocytes describe the condition of the fish's body which is indicate as the fish's body defense against pathogenic bacteria.

Meanwhile, the hematocrit parameter indicated no significantly different. Present results are in contrast with the research of Kanani *et al.*, (2014), which tested the effect of ginger extract supplementation on the hematological parameters of *Huso huso* fish. It is revealed that feeding from plants containing flavonoid compounds can increase the percentage of hematocrit in fish.

On the other hand, the number of hemoglobin showed fish group fed with fermented water hyacinth leaf meal was lower than the fish both in control group and the unfermented treatment group. This is presumably due to the presence of saponin compounds in water hyacinth leaf meal (Lata & Venapani, 2010). Current result is in accordance with the previous study of Lestari *et al.*, (2018) stated that the effect of adding cinnamon leaf powder (*Cinnamomum burmannii*) to the fish feed affected on the blood profile (hemoglobin level) of tilapia (*Oreochromis niloticus*) infected with *Streptococcus agalactiae*. It is also known that hemoglobin level can be decreased due to the presence of saponin compounds found in cinnamon leaf meal. Further, a low number of hemoglobin causes metabolism to be hampered and the energy produced becomes less (Hardi *et al.*, 2011).

Based on Table 2 the platelet of fish fed fermented water hyacinth leaf meal was lower than either the control group or the unfermented treatment group. These results are supported by past research of Puspitowati *et al.*,

(2022), mentioned that the leukocyte of jambal siam fish (*Pangasianodon hypophthalmus*) had decreased after fed with fermented papaya leaf solution (*Carica papaya*). Santoso *et al.*, (2013) revealed that platelets play a role in the blood clotting process and function to prevent loss of body fluids due to infection on the body surface. Platelets are produced to help the blood clotting process to prevent more bleeding. The increase in platelet levels in fish blood is an indicator that fish are in the process of healing wounds (Salim *et al.*, 2016).

In addition, water quality measurement during the study showed a suitable temperature (27.38-27.41 °C), pH (7.50 - 7.57), DO (6.00 - 6.90 mg/L), TAN (0.1 mg/L) and nitrite (0.01 mg/L) for the growth of catfish. According to Pratama *et al.*, (2016) the optimal temperature for catfish is around 20 - 30 °C. Meanwhile, the pH, DO, TAN, and nitrite obtained in the rearing pond for 42 days ranged was in the range tolerance as suggested by the National Standardization Agency (2014), which is 6.5-8.5 (pH), > 4 mg/L (DO), < 1.0 mg/L (TAN), < 1.0 mg/L (Nitrite).

## Conclusions

Based on the results of the study, it can be concluded that, adding fermented 10% water hyacinth leaf meal in the feed has beneficial to improve the growth of sangkuriang catfish. The fermented water hyacinth leaf flour in feed shows a significant effect on the number of erythrocytes, leukocytes, and hemoglobin.

## References

- Amin, M., Taqwa, F. H., Yulisman, Y., Mukti, R. C., Rarassari, M. A., & Antika, R. M. 2020. The Effectiveness of Utilization of Local Raw Materials as Fish Feed on Increasing Productivity of Cultivation of Catfish (*Clarias sp.*) in Sakatiga Village, Indralaya District, Ogan Ilir Regency, South Sumatra. *Journal of Aquaculture and Fish Health*, 9(3), 222.  
<https://doi.org/10.20473/jafh.v9i3.17969>

- Anis, M., Y. & Dyah, H. 2019. Provision of Commercial Feed with the Addition of EM4 (Effective Microorganism 4) to Increase the Growth Rate of Catfish (*Clarias sp.*). Journal of Biological Research and Its Applications, 1 (1), 1 - 8.
- Audila, A., & Zulfahmi, I. S. 2021. PF800 Type Feeding Technique With Paste & Stocking System on Growth Rate of Sangkuriang Catfish Seed (*Clarias gariepinus*). Journal of Biological Sciences and Applied Biology, 1(1).
- Fajriyani, A., Hastuti, S., & Sarjito. 2017. The Effect of Ginger Powder on Feed on Blood Profile, Growth and Survival of Catfish (*Pangasius sp.*). Journal of Aquaculture Management and Technology, 6(4), 39–48.
- Hardi, E. H., Sukenda, E. Harris & Lusiasuti, A. M. (2011). Characteristics and Pathogenicity of *Streptococcus Agalactiae* Type B-Hemolytic and Non-Hemolytic in Tilapia (*Oreochromis niloticus*). Veterinary Journal, 12 (2) : 152 - 164.
- Ikhwanuddin, M., Achmad, N. P., & Mustahal et al., 2018. Utilization of Fermented Rice Bran Using *Aspergillus niger* as Raw Material for Tilapia Fish (*Oreochromis niloticus*) feed. Journal of Fisheries and Marine Affairs, 8 (1) : 79 -87.
- Indariyanti, N. & Rakhmawati. 2013. Improving the Nutritional Quality of Cocoa Fruit and Lamtoro Leaf Waste Through Fermentation as a Protein Base for Tilapia (*Oreochromis niloticus*). Journal of Applied Agricultural Research, (2) : 108-115.
- Kanani, H.G., Nobahar, Z., Kakoolaki, S., & Jafarian. 2014. Effect of Ginger-and Garlic - Supplemented Diet On Growth Performance, Some Hematological Parameters and Immune Responses In Juvenile Huso huso. Fish Physiol Biochem. 40(1) : 481 - 490.
- Ministry of Marine and Fisheries (KKP). 2021. The 2020 Biofloc Program Begins to be Harvested for Food Security and Community Welfare. Public.
- Kurniawan, A. H., Rita, D. R., Suwardiyono. & Imam, S. 2020. Effect of Time and Temperature of Making Activated Carbon from Water Hyacinth (*Eichhornia crassipes*) as an Effort to Utilize High Temperature Waste by Pyrolysis. Journal of Chemical Engineering Innovation, 5(2), 73–80.
- Kurniawan, R., Indra, S. & Adelina. 2022. Utilization of Fermented Water Hyacinth Leaf Flour (*Eichhornia crassipes*) Using Cow Rumen Fluid on the Growth of Baung Fish (*Hemibagrus nemurus*). Journal of Aquatic Science, 10 (1), 31 - 41.
- Lata, N., & Venapani, D. 2010. Preliminary Phytochemical Screening of *Eichhornia crassipes* : The World's Worst Aquatic Weed, Journal of Pharmacy Research, 3 (6) : 1240 - 1242.
- Lestari, D., Sri, S., & Engkus, A. Y., 2021. Comparison of Fermentation Time Using *Aspergillus niger* on Nutrient Levels of Potato Skin. Journal of Animal Husbandry Industry Science and Technology, 1(2), 6-10.
- Lestari, S., Firma, F. R., & Rahmad, J. 2018. Effect of Addition of Cinnamon (*Cinnamomum burmanii*) Leaf Powder to Feed on Blood Profile (Hematocrit Levels, Hemoglobin Levels, Total Leukocytes and Total Erythrocytes) Tilapia (*Oreochromis niloticus*) Infected by *Streptococcus agalactiae*, Journal of Pantura Fisheries (JPP), 1 (1) : 24 - 31.
- Maryanty, Y. P., Hesti, P. & Paulina, R. (2010). Produksi Crude Lipase Dari *Aspergillus niger* Pada Substrat Ongok Menggunakan Metode Fermentasi Fasa Padat. Seminar Rekayasa Kimia dan Proses Semarang Indonesia.
- Marzuqi, M., & Anjusary, D. N. 2013. Digestibility of Feed Nutrients with Different Protein and Fat Levels in Juvenile Sand Grouper (*Epinephelus corallicola*). Journal of Tropical Marine Science and Technology, 5 (2) : 311 - 323.
- Mulyani, Y. S., Yulisman & Fitriani, M. (2014). Growth and Feed Efficiency of Tilapia (*Oreochromis niloticus*) Periodically Fasted. Indonesian Journal of Swamp Aquaculture, 2 (1) : 1 - 12.
- National Standardization Agency (BSN). 2014. Catfish (*Clarias sp.*) Part 3: Parent Production.



- Pane, A., S., Yuniarti., & Juliana. 2018. The Effect of Addition of Water Hyacinth Leaf Flour (*Eichhornia crassipes*) on the Making of Feed on the Growth and Survival of Sangkuriang Catfish (*Clarias gariepinus*) Seeds. Scientific Journal of Fisheries and Marine Affairs, 6(3), 246–252.
- Prasekti, Y., H. & Irfan, Y., H. 2021. Analysis of the Sangkuriang Catfish (*Clarias gariepinus*) hatchery Case Study at BBI Bolorejo, Kauman District, Tulungagung Regency. Journal of Agribis, 7(1), 1–6.
- Pratama, F. A., Afiati, N., & Djunaedi, A. 2016. Conditions of Aquaculture Pond Water Quality with the Use of Probiotics and Without Probiotics on the Growth of Sangkuriang Catfish (*Clarias gariepinus*) in Cirebon, West Java. Diponegoro Journal of Maquares. 5(1), 38 - 45.
- Puspitowati, D., Iesje, L., & Henni, S. 2022. Leukocyte description of Jambal Siam (*Pangasianodon hypophthalmus*) fish fed with fermented papaya leaf solution (*Carica papaya L.*). Journal of Aquaculture Sebatin, 3 (1) : 78-92.
- Putra, A. N., Ristiani, S., Musfiroh, M., & Syamsunarno, M. B. 2020. Utilization of Water Hyacinth (*Eichhornia crassipes*) as Tilapia Fish Feed: Effects on Feed Growth and Digestibility. Leuit (Journal of Local Food Security), 1(2), 77. <https://doi.org/10.37818/leuit.v1i2.10016>
- Putri, R. R., Basuki, F., & Hastuti, S. 2013. Blood profile and survival rate of F5 Pandu Tilapia (*Oreochromis niloticus*) Infected with *Streptococcus agalactiae* Bacteria with Different Densities. Journal Of Aquaculture Management and Technology, 2 (2) : 47 - 56.
- Rahayu, W. T. 2020. Effect of Various Carriers in Solid Starters on Proximate Content in Fermented Water Hyacinth (*Eichhornia crassipes*). Journal of Agromedia, 38(2), 10–17.
- Raudah, P., Indra, S., & Hamdan, A. 2018. Utilization of Fermented Lamtoro Gung (*Leucaena leucocephala*) Leaf Flour *Aspergillus niger* as a Protein Substitute for Soy Flour in Feed Against the Growth of Siamese Catfish (*Pangasius hypophthalmus*). Journal of Fisheries and Marine Affairs, 23(2), 1-8.
- Salim, M. A., Indriyani, N., & Muhammad, I. 2016. Effect of Gradually Increasing Salinity on Differential Leukocytes in Tilapia (*Oreochromis niloticus*). Journal of Aquatic Media, 1 (1) : 52 - 58.
- Santoso, B. B., Fajar, B., & Sri, H. 2013. Analysis of Body Resistance of Tilapia larasati (*Oreochromis niloticus*) Hybrid Seeds Generation 5 (F5) Infected with *Streptococcus agalactiae* Bacteria with Different Concentrations. Journal of Aquaculture Management and Technology, 2 (3) : 64 - 75.
- Setiawan, A. S. 2013. Efficiency of Protein Use in Male Pengging Ducks (*Anas platyrhynchos*) Given Water Hyacinth (*Eichhornia crassipes*) Fermented in Ration. Journal of Agromedia, 31(2), 9–19.
- Setiawati, J. E., Adiputra, Y. T., & Hudaidah, S. 2013. Effect of Addition of Probiotics to Feed with Different Doses on Growth, Life Graduation, Feed Efficiency and Protein Retention of Catfish (*Pangasius hypophthalmus*). Journal of Aquaculture Engineering and Technology, 1(2), 7 – 13.
- Simanjuntak, N., Putra, I., & Pamukas, N. A. (2020). Effect of EM4 Probiotics in Feed on Growth and Survival of Sangkuriang Catfish (*Clarias sp*) Seeds with Biofloc Technology. SEBATIN Journal of Aquaculture, 1(1), 63 - 69.
- Sorongan, J.V., Andaki, J. A., & Kotambunan, O. V. 2019. Farmers' Perceptions of Catfish Cultivation Development in Warukapas Village, Dimembe District, North Minahasa Regency, North Sulawesi Province. ACULTURATION (Scientific Journal of Fisheries Agribusiness), 6(12).
- Suci, R., Rudy, A., N., Retno, A., & Hetty, M., Rudianto. 2020. Evaluation of Dietary Choline and Methionine on Growth and Feed Efficiency of Carp (*Cyprinus carpio*). Indonesian Journal Of Fisheries Science And Technology. 16 (2) : 97-103.
- Syaraswati, M., Indra, S. & Adelina. 2022. Utilization of *Aspergillus niger* (*Eichhornia crassipes*) Leaf Flour Fermented in Feed for Growth of

- Nilem Fish (*Osteochilus hasselti*) Seeds. The Best Fisheries Periodic Journal, 50(1), 1394 - 1405.
- Tamba, J. T. 2020. Supplementation of Cinnamon Leaf (*Cinnamomum burmannii*) Different Concentrations in Feed on the Growth of Snakehead Fish (*Channa striata*). *Journal of Tropical Fisheries*, 7(20), 35–44.
- Widyastuti., Sukanto. & Rukaya. 2010. Use of Fermented Feed in Floating Cage Fish Cage System to Reduce Eutrophication Potential in Wadas Lintang Reservoir. *Limnotek Journal*, 17 (2) : 191 - 200.
- Wijaya, D., Putri, P. Y., Raffty, S. A., & Muhammad, R. 2015. Screening of Phytochemicals and Antioxidant Activity of Water Hyacinth Leaves (*Eichhornia crassipes*). *Journal of VALENCY Chemistry*, 1(1), 65–69. <https://doi.org/10.15408/jkv.v0i0.4965>
- Zhai, S. W., & Liu, S. L. 2013. Effects of Dietary Quercetin on Growth performance Serum Lipids Level and Body Composition of Tilapia (*Oreochromis niloticus*). *Italian Journal of Animal Science*, 12 (4) : 523 - 527.
- Zissalwa, F., Henni, S., & Iesje, L. 2020. Erythrocyte profile of Jambal Siam (*Pangasius hypophthalmus*) which is fed with extracts of mangrove leaves (*Rhizophora apiculata*) and kept in cages. *Journal of Fisheries and Marine Affairs*, 25 (1) : 70 - 78.

ORIGINALITY REPORT

13%

SIMILARITY INDEX

13%

INTERNET SOURCES

5%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Syiah Kuala University Student Paper	3%
2	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1%
3	<a href="http://fison.org.ng">fison.org.ng</a> Internet Source	1%
4	L Anna, Muhammadar, I Sahidhir. " The effect of probiotics (RABAL) on application in diet with different doses on growth, survival and feed conversion of Barramundi ( ) ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1%
5	<a href="http://hdl.handle.net">hdl.handle.net</a> Internet Source	1%
6	<a href="http://ajoas.ejournal.unri.ac.id">ajoas.ejournal.unri.ac.id</a> Internet Source	1%
7	<a href="http://www.jurnal.unsyiah.ac.id">www.jurnal.unsyiah.ac.id</a> Internet Source	1%

8	<a href="http://www.iosrjournals.org">www.iosrjournals.org</a> Internet Source	1 %
9	<a href="http://zsp.com.pk">zsp.com.pk</a> Internet Source	1 %
10	<a href="http://ojs.unimal.ac.id">ojs.unimal.ac.id</a> Internet Source	1 %
11	<a href="http://ijariie.com">ijariie.com</a> Internet Source	1 %
12	I Suharman, I Lukistyowati, S Ramayani, C M A Caipang, A Adelina, N Aryani. "Quality Improvement of Water Hyacinth (Eichornia Crassipes) Leaf Meal Fermented with Aspergillus Niger as Fish Feed Ingredient", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1 %
13	Shahida Naseem, Sami Ullah Bhat, Adil Gani, Farooz Ahmad Bhat. "Perspectives on utilization of macrophytes as feed ingredient for fish in future aquaculture", Reviews in Aquaculture, 2020 Publication	1 %
14	<a href="http://brec-currentops.wm.com">brec-currentops.wm.com</a> Internet Source	1 %
15	<a href="http://f1000researchdata.s3.amazonaws.com">f1000researchdata.s3.amazonaws.com</a> Internet Source	1 %

---

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography On