(26) Financial Analysis of Arabica Coffee Cultivation of Agroforestry system

by Yosep Ruslim

Submission date: 03-May-2021 09:20AM (UTC+0700)

Submission ID: 1576290004

File name: nalysis_of_Arabica_Coffee_Cultivation_of_Agroforestry_system.pdf (541.19K)

Word count: 4579

Character count: 24216



Proceedings of the Joint Symposium on Tropical Studies (JSTS-19)

Financial Analysis of Arabica Coffee Cultivation of Agroforestry Systems in Lembang Bokin, North Toraja Indonesia

Andi Lisnawati^{1,*} Abubakar M. Lahjie² Syahrir Yusuf² Yosep Ruslim²

¹Department of Agriculture Technology, Agricultural Polytechnic of Samarinda, Kampus Gunung Panjang, 13 marinda

² Faculty of Forestry, Mulawarman University, Campus of Gunung Kelua, Jl. Penajam, Samarinda *Corresponding author. Email: andilisnawatismrd75@gmail.com; ** Email:yruslim@gmail.com

ABSTRACT

Toraja is a centre for a coffee product in South Sulawesi. One of the most famous species is Arabica, one of the national superior-commodities with a high economic value. Given the demand for coffee, Arabica coffee needs to develop the plantation of Arabica coffee in particular. Therefore, it is crucial to analyze the financial feasibility of Arabica coffee for each type of shade. This research was conducted in some cultivated areas of Arabica coffee in Lembang Bokin, North Toraja. It is located at ± 1,237 meters above sea level, and it has coordinates of 03°01'04,7" – 03°01'45,2"N and 119°59'44,9" – 120°00'03,8"S. The study was aimed to analyze the financial feasibility of Arabica coffee cultivation with a combination of *Leucaena glauca* and *Calliandra calothyrsus* and *Elmerrellia ovalis* (agroforestry models) by calculating: (i) the Net Present Value (NPV); (ii) the Net Benefit-Cost Ratio (Net B/C) and (iii)) Internal Return Rate (IRR). The financial analysis of Arabica coffee under combination *Leucaena glauca* and *Calliandra Calothyrsus* was at an interest rate of 10%. The financial analysis result suggests that the production of Arabica coffee shaded by Leucaena glauca and *Calliandra calothyrsus* was feasible because financial analysis showed a positive NPV value B/C>1, and IRR value (greater than MAR value =10%). On the other hand, the Arabica coffee shaded by *Elmerrellia ovalis* was not economically feasible because it gave a negative NPV value and Net B/C <1, IRR less than the MAR value of 5.4%.

Keywords: Agroforestry, Arabica Coffee cultivation, financial feasibility analysis

1. INTRODUCTION

The plantation commodity sectors have significant roles in the economic growth of a country worldwide. Coffee is one of the plantation commodity exports, which gives considerable benefits to farmers, employers, and governments in terms of income and foreign exchange; and affects global aspects, including economy, ecology, and social sectors. Besides, the coffee plantation can also support the development of regional and agro-industry in Indonesia. Lembang Bokin in North Toraja is one of the districts located in South Sulawesi Province with the abundant natural resources of coffee types. It is also best known as the center of coffee plantation, for Arabica and Robusta type, which is intensively developed by local farmers, foreign and domestic industries.

Quantitative data on Arabica coffee cultivation using an agroforestry system is needed to stimulate people's interest in conserving trees. Agroforestry is one of the suggested methods for conserving biodiversity, producing food crops, and providing other ecosystem services such as climate change mitigation and carbon deposition. It's land utilization system consists of a mixture of hard plants with or without annual plants [1,2]. The agroforestry system applied to the coffee plant with shade trees is a promising alternative to improve the agricultural system [3]. In a simple agroforestry system, understanding the characteristics and varieties of the shade trees and its function can advance the sustainability of coffee agroecosystem [4] and improve biodiversity conservation [5].

Shade trees used to shade Arabica coffee trees are a combination of (Leucaena glauca and Calliandra



calothyrsus) and Ermerellia ovalis. This plant is used as shade due easy to grow and can adapt to the environment. Growing shade trees is a way that can be undertaken to reach an environmental condition that suitable for coffee growth. Subsequently, giving shade trees on Arabica coffee will reduce light intensity that exposes plants to change the plants' temperature and humidity. The use of shade trees on coffee will influence biotic and abiotic factors, such as other research findings that would affect the temperature, humidity, and wind [6,7]. Agroforestry can be considered a land-sharing strategy to conserve biodiversity and increase agricultural production [8]. An agroforestry coffee system can also maintain the amount of mycorrhiza in the soil effectively compared to the monoculture system [9].

Different shade trees will also give a different light intensity that affects plant physiological and morphological factors and affect plant production [10]. The study was aimed to analyze (1) the financial feasibility of Arabica coffee cultivation with a combination of (*Leucaena glauca* and *calliandra calothyrsus*) and *Elmerrellia ovalis* (agroforestry models) by calculating: the Net Present Value (NPV), the Net Benefit-Cost Ratio (Net B/C), and Internal Return Rate (IRR); (2) How is the development of coffee based on the type of shade and market opportunity.

2. MATERIAL AND METHODS

2.1. Study Area

This study was conducted in Lembang Bokin North Toraja, South Sulawesi province. Bokin village is administratively located in Rantebua district, north Toraja, South Sulawesi province. The geographical location of Bokin village was 03 01 '04,7" - 03 01 '45,2" n and 119°58'44,9" - 120° 00'03,8" (Fig 1) with the temperature of 22°-24°c and relative humidity (Rh) 80 %. From Rantepao City, Bokin village can be reached in 45 minutes by motor vehicles. The total area of Bokin village is about 2.069 ha, which consists of the settlement area, forest estate, and people's farm and orchard. The majority of the Bokin village population was the Toraja tribe people, and their main livelihood came from coffee cultivation. North Toraja is one of South Sulawesi's districts, Indonesia, famous for its Arabica coffee. One of the most famous species is Arabica, one of the national superior-commodities with a high economic value. The research location was selected by purposive sampling on 17-year-old Arabica coffee plants with spacing 2.5 m x 2.5 m, and direct variables measured in this study including light intensity and several models of shade trees used in the coffee plantation such as the combination of (Leucaena glauca and Calliandra calothyrsus) with spacing 6 m x 4 m and Ermerellia ovalis 8 m x 7 m.




Figure 1 Research location in Lembang Bokin village, Rantebua district village, North Toraja South Sulawesi Province, Indonesia.



2.2. Data Analysis

Production and financial analyses were done for two models of Arabica coffee cultivation, namely: (1) Arabica coffee plantation under shade tree combination/mixed *Leucaena glauca* and *Calliandra Calothyrsus* (Model 1), (2) Arabica coffee under shade tree *Elmerrellia ovalis* (Model 2).

Specifically, data collection was done in the following manner [11]: (1) observation by conducting direct observation on Arabica coffee cultivation activities, including activities undertaken, production costs, and earned income, (2) direct measurements in the field, including diameter Arabica coffee trees, production of cheery, the weight of cherry and dry bean coffee, (3) library research, data collecting through literature review [15]d reports from institutions related to Arabica coffee, Arabica coffee cultivation activities in north Toraja district, south Sulawesi Province; (4) structured interviews with questionnaires, discussions and direct interviews with Arabica coffee farmers and companies.

The method used in the financial feasibility study of this Arabica coffee cultivation is the present value (present value) because it comes from the data at the time of the study. The assumptions used are as follows: (1) costs and benefit are the present value, a value that applied on the market in the year of the research, (2) the age of Arabica coffee is 3-25 years. It is seen from the data obtained when researching primary data and secondary data obtained from the field the selling price applied in 2017, (3) the prices that used in 2017, (4) labour cost is IDR 80,000 day-1, (5) discount rate 10%

In the financial analysis, data analysis of each activity step of Arabica coffee cultivation is cost component analysis, income from Arabica coffee cultivation business, and feasibility analysis using NVP, Net B/C Ratio, and IRR parameters necessary. Analysis of maximum coffee yield was done based on the measurement of time interval (cycle) by calculating average annual production average product (AP) and marginal product (MP) [12] with the formula as follows:

AP = Pt/t

Where:

AP = average product of arabica coffee
Pt = total product at age t of arabica coffee

T = tree age of coffee

 $MP = \frac{Pt - P_{t-1}}{Tt - T_{t-1}}$

Where:

MP = marginal product of arabica coffee

Pt = total product at age t of arabica coffee

P_{t-1} = total product at age t-1 of arabica coffee

T = time interval between each measurement age of arabica coffee

To calculate how much investment, the financial feasibility in coffee cultivation with agroforestry systems by the company was analyzed using investment criteria [13,14].

 $NPV = \sum_{t=0}^{t=n} \frac{Bt - Ct}{(1+i)^t}$

$$Net \, B/C = \frac{\sum_{t=1}^{t=n} \frac{Bt-Ct}{(1+t)^t}}{\sum_{t=n}^{t=n} \frac{Bt-Ct}{(1+t)^t}} \longrightarrow Bt - Ct > 0$$

The criter 19 used in evaluating the business feasibility was the internal rate of return (IRR). IRR is a mean annual return derived from an investment and expressed in percentage [15]. IRR value indicates an interest rate that can be paid by a business, or in other words, the ability to gain income from the cost invested.

IRR =
$$i_1 + \frac{NPV1}{NPV1 - NPV2} (i_2 - i_1)$$

Where:

NPV1 = positive NPV of arabica coffee NPV2 = negative NPV of arabica coffee i_1 = interest rate when NPV is positive. i_2 = interest rate when NPV is negative.

3. RESULTS AND DISCUSSION

Coffee has become one of the plantation crops that have been cultivated and have high economic value [16]. The coffee growth needs shade trees in the cultivation. Without preparing the fine quality of land and shade trees, it is hard to reach successful coffee growth. Ten species (28%) of thirty-six kinds of a shade tree in Sumberjaya Lampung are originated from legumes, Gliricida sepium, Dalbergia latifolia, namely Paraserianthes falcataria, Parkia speciosa, Acacia sp., Archidendron pauciflorum and Archidendron microcarpum [17], Calliandra calothyrsus, and Leucaena glauca [18].

The use of shade trees in coffee plantations depends on many factors. The most crucial factor is the one related to production, increasing crop yields, and as a 11 dition modifier of environment that appropriate [19]. The use of different shade trees on coffee undoubtedly results in a different physiological response from the other canopy of shade trees plants. The shade tree is used to manage the intensity of sunlight that enters and is used by coffee for the flowering and fertilization process. The growth and development of coffee can also be affected by the shade trees' condition [19]. The difference occurred because of the different shade trees being used. The mixed shade trees of Calliandra calothyrsus and Leucaena glauca are different from



Ermerellia ovalis trees in terms of the trunks, size of the trunk, and leaves, so that it causes the light intensity received by the Arabica coffee differently. Some studies reported that the dominant shade tree species affect coffee's growth and productivity [20,21].




Figure 2 Correlation between average product (AP) and marginal product (MP) of Arabica coffee under mixed *C. Calothyrsus* and *L. Glauca* shading.




Figure 3 Correlation between average product (AP) and marginal product (MP) of Arabica coffee under *Emerallis ovalis* shading.

The total production of dried coffee seeds from the research result shows the increase of production in each year along with the increase of the age of the tree, that age of coffee trees affects its productivity [22] for the shade tree mixed *Calliandra calothyrsus* and *Leucaena glauca* annual average of production 1135 kg ha⁻¹year⁻¹ or coffee production 1.42 kg tree⁻¹ha⁻¹ with increased by 12.92% year⁻¹ (Fig 2). Meanwhile, under shade trees, *Elmerrellia ovalis* production reached 489 kg ha⁻¹year⁻¹ or coffee production, 0.60 kg tree⁻¹ha⁻¹ increased by 11.59% year⁻¹ by the age of 17. It shows that coffee production maximizes both types of shade trees (Fig 3).

It can be seen (Fig 2) from the intersection of the average product (AP) reached 100 kg ha⁻¹year⁻¹ and marginal product (MP) 100 kg ha⁻¹ year⁻¹ or coffee production 1.42 kg trees⁻¹ha⁻¹for the mixed shade tree of *C. calothyrsus* and *L. glauca*. Meanwhile, tree *E. Ovalis* production reached an average 76.47 kg tree⁻¹ha⁻¹and marginal product (MP) 76.50 kg ha⁻¹year⁻¹ (Fig 3). The

dominant shade of legume tree had a positive influence on coffee productivity [23]. There was a positive correlation between shade trees with coffee productivity [17]. Coffee plant growth and productivity with and without shading showed different results. Coffee plants under shading conditions y 6d much more significant and heavier produce [24]. The use of shade trees in coffee plantations depends on many factors. The most crucial factor is the one related to production, increasing crop yields, and as a condition modifier of environment that appropriate [19]. The traditional agroforestry system is a century-old agricultural practice. It has become an essential agricultural model globally, especially in the tropical and sub-tropical areas, from the economic, ecological, and socio-cultural viewpoints, profitable to the community that runs it [25]. Companies and Toraja people use Uru-Uru's shade tree (Elmerrellia ovalis) as an agroforestry coffee cultivation system. Moreover, because the wood is substantial and round, widely used as building materials for traditional Toraja

Financial analysis on coffee cultivation is the effort made to get an idea of how much activity was undertaken to provide benefits that outweigh the costs (investment) issued. The cultivation cost was including the starting cost for preparing the field up to the cost of harvesting [26]. Activities done in the coffee cultivation consisted of planning, field preparation, supplying seeds, planting, stitching, maintenance, and harvesting. Activities such as planning, field preparation, seed supply, and planting are carried out in the first year. Plant planting is only done in the second year. Maintenance of growing plants is carried out once a year, while harvesting starts in the third year for Arabica coffee. Maintenance is an activity in the form of weeding plants against weeds and pests, and diseases. This activity is carried out every year. The day needed for maintenance in 1 ha is five days, and the cost of maintenance is IDR 2,400,000 ha⁻¹year⁻¹ for a 30 person day. Meanwhile, harvesting coffee cherries in 1 hectare in a year is needed 30 days, and the cost for harvesting is IDR 12,112,500 ha⁻¹ year⁻¹ for 150 person day.

According to the market prices, when this study was conducted, Arabica coffee's local market prices were Arabica coffee under mixed *Calliandra calothyrsus* and *Leucaena glauca* IDR 57,000 kg⁻¹, and Arabica coffee under *Elmerrellia ovalis* IDR 45,000 kg⁻¹.

The financial analysis of Aral a coffee cultivation in the community forest used an internal rate of return (IRR) criterion. This discount rate gives the result to a net present value (NPV) of zero. A feasible cultivation business is indicated by the more significant IRR value than the interest rate applied when the investment takes place [15].

Based on the research results at Lembang Bokin, North Toraja, each investment criterion's value can be



4					
Advances in	Biological	Sciences	Research,	volume	11

Table 1. Details of analyzing Arabica coffee cultivation (25-year cycle) with discount rate 10%

Shade Tree	Analysis	Total	Criteria	Value
C. calothyrsus and L. glauca	Gross Benefit	1,630,300,000	Net B/C Ratio	3.07
	Cost	944,500,000	Gross B/C Ratio	1.73
	Investasi	31,640,000	NPV	99,980,000
	Net Benefit	559,693,000	IRR	19.3
Elmerrellia ovalis	Gross Benefit	554,661,000	Net B/C Ratio	0.50
	Cost	459,051,000	Gross B/C Ratio	1.21
	Investasi	30,255,000	NPV	28,653,000
	Net Benefit	96,610,000	IRR	5.3

Source: mimary data after processed, 1\$ = IDR 13,000,-

Note: NPV = Net Present Value, IRR = Internal Rate of Return, Net B/C Ratio = Net Benefit Cost Ratio, Gross B/C Ratio = Gross Benefit Cost Ratio

seen in Table 1. According to calculation, the cash flow of coffee under mixed C. calothyrsus and L. glauca at 25 years plant cycles shows that the total cost for the entire activity amounted to IDR 944,500,000 and gross benefit IDR 1,630,300,000. Financial analysis of Arabica coffee cultivation with agroforestry systems using mixed shade (C. calothyrsi 12 nd L. glauca) at an interest rate of 10%, resulting in Net Present Value (NPV) and Net B/C resulting in Net Present Value were IDR 99,980,000 and 3,07 respectively. This statement is strengthened by an analysis of the Internal Rate of Return (IRR) model with a value of 19.3%. The results above show that Arabica coffee cultivation 24 er mixed shade (C. calothyrsus and L. glauca) at 25 years and an interest rate of 10% is feasible, which means that it is profitable. The figureial analysis of Arabica coffee cultivation used an internal rate of return (IRF23 riterion, a discount rate that gives the result to a net present value (NPV) of zero. Meanwhile, the Net of B/C of the business amounted to 3.07, which means that the value of the rupiah investment return was 3.07 times the value of the rupiah invested. The Net B/C >1 indicates that the business is profitable.

Meanwhile, according to the calculation of the cash flow of coffee under E. ovalis at 25 years plant cycles, the total cost for the entire activity amounted to IDR 459,051,000 and gross benefit of IDR 554,661,000. Financial analysis of Arabica coffee cultivation with agroforestry systems, \mathfrak{g} de tree E. ovalis at an interest rate of 10%, resulting Net Present Value (NPV) and Net B/C resulting in Net Present of values were IDR 28,653,000 and 0.50 resp3tively. This statement is strengthened by analyzing the Internal Rate of Return (IRR) model with a value of 5.4%. The results above show that the Arabica coffee cultivation under shade tree E. ovalis 18 25 years and an interest rate of 10% are not feasible, which means it is not profitable. Meanwhile, the Net of B/C of the business amounted to 0.50, which means that the value of the rupiah investment return was

0.50 times the value of the rupiah invested. The Net B/C <1 indicates that the business is not profitable and indicated the cultivation model was not feasible to be executed because the profit was still far smaller than the actual bank interest rate (10%). The Arabica coffee cultivation using a combination of *C. calothyrsus* and *L. glauca* can be further developed because this shade provides more outstanding coffee production than the shade of *E. ovalis*. Shade trees like the combination *C. calothyrsus* and *L. glauca* in the agroforestry system offer many benefits, i.e., increasing organic matter and soil nutrients through litter and nitrogen fixation legumes as shade trees, to encourage biodiversity function and to suppress the weed growth [19].

The use of shade trees in coffee plantations related to production, increase crop yields, and as a continuous of different shade trees on coffee undoubtedly results in a different physiological response that coffee received due to the different canopy of shade trees plants. The shade tree is used to manage the intensity of sunlight that enters and is used by coffee for the flowering and fertilization process. The growth and development of coffee can also be affected by the shade trees' condition [20].

4. CONCLUSION

The conclusions that can be derived from this study were the combination of (*C. calothyrsus and L glauca*) provides the highest productivity compared to *E. ovalis* with an annual average of production 1135kgha⁻¹year⁻¹ or coffee production of 1,42 kg tree⁻¹ha⁻¹ with increased by 12,92% year⁻¹, the maximum output of Arabica coffee in model 1 under mixed (*C. calothyrsus* and *L. glauca*) and model 2.

E.ovalis was reached at 17-year Arabica Coffee in model 1 under shade tree (C. calothyrsus and L.



glauca) resulted in IRR 17.2%, Net B/C Ratio 2,43 (>1) and meanwhile model 2 IRR 5.4%; Net B/C Ratio 0.5 (<1). Arabica coffee cultivation models under the shade tree mixed (*C. calothyrsus* and *L. glauca*) are feasible to be developed, while under the shade tree, *E. ovalis* is not feasible.

ACKNOWLEDGMENT

This research was funded by the Directorate General of Research and Development, The Ministry of Research, Technology and The Higher Education, the Republic of Indonesia through Doctoral Dissertation Grant 2017. The authors wish to thank the Director of Toarco Company and the staff to kindly allow us to conduct research in their place and give necessary and valuable data. We also would like to thank the people around the research site and to all people who have supported us from the planning stage, data analysis, up to the research report.

REFERENCES

- [1] M.D. Beenhouwer, L. Geeraert, J. Martens, M.V. Geel, R. Aerts, K. Vanderhaegen, O. Honnay, Godiversity and carbon storage co-benefits of coffee agroforestry across a gradient of increasing management intensity in the SW Ethiopian highlands, Agriculture, Ecosystems & Environment, 222, 2016, pp. 193–199. DOI: https://doi.org/10.1016/j.agee.2016.02.017
- [2] Wiryono, V.N.U. Puteri, G. Senoaji, The diversity 21 plant species, the types of plant uses and the estimate of carbon stock in agroforestry system in Harapan Makmur Village, Bengkulu, Indonesia, Biodiversitas, 17(1), 2016, pp. 249–255. DOI: 10.13057/biodiv/d170136
- [3] C.R. Cerdin, M.C. Rebolledo, G. Soto, B. Rapidel, F.L. Sinclair, Local knowledge of impacts of tree cover on ecosystem services in smallholder coffee production systems, Agricultural Systems, 110, 2012, pp. 119–130. DOI: https://doi.org/10.1016/j.agsy.2012.03.014
- [4] F.M. Mamani-Pati, D.E. Clay, S.A. Clay, H. Smeltekop, M.A. Yujra-Callata, The influence of strata on the nutrient recycling within a tropical certified organic coffee production system, International Scholarly Research Notices, 2012, Article ID 389290, 8 pages. DOI: https://doi.org/10.5402/2012/389290
- [5] G. Lopez-Rodriguez, D. Sotomayor-Ramirez, J.A. Amador, E.C. Schroder, Contribution of nitrogen from litter and soil mineralization to shade and sun coffee (*Coffee arabica* L.) agroecosystems, Tropical Ecology, 56, 2015, pp. 21–33.

- [6] J. Jaramillo, M. Setamou, E. Muchugu, A. Chabi-Olaye, A. Jaramillo, Climate change or 17 anization? Impacts on a traditional coffee production system in East Africa over the last 80 years, PLOS ONE, 8(1): e51815. DOI: https://doi.org/10.1371/journal.pone.0051815
- [7] Y.A. Marino, M.E. Perez, F. Gallardo, M. Trifilio, M. Cruz, P. Bayman, Sun vs. shade affects infestation, total population and sex ratio of the coffee berry borer (*Hypothenemus hampei*) in Puerto Rico, Agricultural, Ecosystems & Environment, 222, 2016, pp. 258–266. DOI: https://doi.org/10.1016/j.agee.2015.12.031
- [8] J. Fischer, DJ Abson, V Butsic, MJ Chappell, J Ekroos, J Hanspach, T Kuemmerle, HG Smith, H Wehrden, Land sparing versus land sharing: moving forward, Conservation Letters, 2014, pp. 149–157. DOI:https://dx.doi.org/10.1111/conl.12084
- [9] D. Muleta, F. Assefa, S. Nemomissa, U. Granhall, 22 tribution of arbuscular mycorrhizal fungi spores in soils of smallholder agroforestry and monocultural coffee system in southwestern Ethiopia, Biology and Fertility Soils 44, 2008, pp. 653–659. DOI: https://doi.org/10.1007/s00374-007-026-3
- [10] U. Sholikha, D.A. Arimunandar, S.A. Pradana, Robusta coffee physiological characters BP 358 on different types of shade, Agrovigor 8(1), 2015, pp. 58–67 DOI:https://doi.org/10.21107/agrovigor.v8i1.749
- [11] E. Linger, Agro-ecosystem and socio-economic role of home garden agroforestry in Jabithenan District, North-Western Ethiopia: Implication for climate change adaptation, SpringerPlus 3: 154, 2014, pp. 1–9. http://www.springerplus.com/content/3/1/154
- [12] P.R. van Gardingen, M.J. McLeish, P.D. Philips, D. Fadilah, G. Tyrie, I. Yasman, Financial and ecological analysis of management options for logged-over dipterocarp forest in Indonesia Borneo, Forest Ecology and Management 183, 2003, pp. 1–29. DOI:https://doi.org/10.1016/S03781127(03)00097-5
- [13] J.P. Gittinger. Analisa Ekonomi Proyek-proyek Pertanian, Second Edition, University of Indonesia Publisher, Jakarta, 1986.
- [14] C. Gray, P Simanjutak, L.K. Sabur, P.F.L. Maspaitella, R.C.G. Varley, Pengantar Evaluasi Proyek, Second Edition, Gramedia Jakarta, 1993.
- [15] A.R. Graves, P.J. Burgess, J.H.N. Palma, F. Herzog, G. Moreno, M. Bertomeu, C. Dupraz, F. Liagre, K. Keesman, W. van der Werf, A.K. de



- Nooy, J.P. van den Briel, Development and plication of bioeconomic modelling to compare silvoarable, arable, and forestry systems in three European countries, Ecologi Bengineering 29 (4), 2007, pp. 434–449. DOI: https://doi.org/10.1016/j.ecoleng.2006.09.018
- [16] P.K.P. Nair, The agronomy and economy of important tree crops of the developing world, 2010. DOI: 10.1016/C2010-0-64818-8.
- [17] R. Evizal, Sugiatno, F.E. I. Prasmatiwi, Nurmayasari, Shade tree species diversity and coffee productivity in Sumberjaya, West Lampung, Indonesia. Biodiversitas 17(1), 2016, pp. 234–240. DOI: https://doi.org/10.13057/biodiv/d170134
- [18] A. Lisnawati, A.M. Lahjie, B.D.A.S. Simarangkir, S. Yusuf, Y. Ruslim, Agroforestry system biodiversity of Arabica coffee cultivation in North Toraja District, South Sulawesi, Indonesia, Biodiversitas 18(2), 2017, pp. 741–751. DOI: https://doi.org/10.13507/biodiv/d180243
- [19] T. Tscharntke, Y. Clough, S.A. Bhagwat, D. Buchori, H. Faust, D. Hertel, D.H. Ischer, J. Juhrbandt, M. Kessler, I. Perfecto, C. Scherber, G. Schroth, E. Veldkamp, Multifunctional shade-tree management in tropical agroforestry landscapes-a review, Journal Applied of Ecology 48, 2011, pp. 619–629. DOI: https://doi.org/10.1111/j.1365-2664.2010.01939x
- [20] R.G. Muschler, Shade or sun for ecologically 10 ainable coffee production: a summary of environmental key factors, in: III. Semana Científica del Centro Agronómico Tropical de Investigación y Enseñanza (CATIE), Turrialba, Costa Rica 1997, pp 109–112.
- [21] A.A. Muliasari, The growth of seedlings arabica coffee (Coffea arabica L.) on the application anorganic-organic fertilizer and the level of intensity of the shade, Thesis, Institut Pertanian Bogor, Bogor, 2016.
- [22] T. Kufa, M.J. Burkhardt, Plant composition and 25 wth of wild *Coffea arabica*: Implications for management and conservation of natural forest resources, International Journal of Biodiversity and Conservation 3, 2011, pp. 131–141.
- [23] L. Ebisa, Effect of dominant shade trees on coffee production in Manasibu District, West Oromia, Ethiopia, Science, Technology and Art Research Journal 3, 2014, pp. 18–22, https://dx.doi.org/ 10.4314/star.v313.3

- [24] C. Potvin, C.T. Owen, S. Melzi, P. Beaucage, Biodiversity and modernization in four coffeeproducing villages of Mexico, Ecology and Society 10(1), 18, 2005, http://www.ecologyand society.org/vol10/iss1/art18
- [25] R.A. Rice, Agricultural intensification within agroforestry: The case of coffee and wood products, Agriculture, Ecosystem and Environment 128, 2008, pp. 212–218. DOI: https://doi.org/10.1016/j.agee.2008.06/007
- [26] A.D. Bote, P.C. Struik, Effects of shade on growth, production and quality of coffee (*Coffea arabica*) in Ethiopia, Journal of Horticulture Forestry, 3, 2011, pp. 336–341.

(26) Financial Analysis of Arabica Coffee Cultivation of Agroforestry system

ORIGINALITY REPORT

8% SIMILARITY INDEX

%
INTERNET SOURCES

8%

%

PUBLICATIONS

STUDENT PAPERS

PRIMARY SOURCES

Jenisa Devy, Ali Ibrahim Hasyim, Suriaty Situmorang. "ANALISIS KELAYAKAN FINANSIAL DAN RISIKO USAHA BUDIDAYA JAMUR TIRAM DI PROVINSI LAMPUNG", Jurnal Ilmu-Ilmu Agribisnis, 2019

1 %

Publication

Nengah I. K. Besung, Gusti Ketut Suarjana, Karang Kadek Agustina, Oka Ida Bagus Winaya et al. "Isolation and Identification of Streptococcus suis from sick pigs in Bali, Indonesia", Research Square, 2019

1 %

van Gardingen, P.R.. "Financial and ecological analysis of management options for logged-over Dipterocarp forests in Indonesian Borneo", Forest Ecology and Management, 20030915

<1%

Publication

Publication

4

"Quantitative systems pharmacology for shifting the drug discovery and development

<1%

paradigm", Biopharmaceutics & Drug Disposition, 2013

Publication

Bioenergy, 2020

Publication

Chris N. Namah, Dina V. Sinlae. "The <1% 5 Feasibility Study of Keprok SoE Citrus Farming and the Development Strategis of Local Citrus Farming", Procedia Economics and Finance, 2012 Publication "Directions in Tropical Agroforestry Research", <1% 6 Springer Science and Business Media LLC, 1998 Publication Castellazzi, M.S.. "Simulation scenarios of <1% spatio-temporal arrangement of crops at the landscape scale", Environmental Modelling and Software, 201012 **Publication** Mirza Faisal Qaseem, Ai - Min Wu. "Marginal <1% 8 lands for bioenergy in China; an outlook in status, potential and management", GCB

Sandro L.S. Moreira, Cleverson V. Pires,
Gustavo E. Marcatti, Ricardo H.S. Santos et al.
"Intercropping of coffee with the palm tree,
macauba, can mitigate climate change

Publication

Fábio M DaMatta. "Ecophysiological constraints on the production of shaded and unshaded coffee: a review", Field Crops Research, 2004

<1%

Publication

J. Beer, R. Muschler, D. Kass, E. Somarriba.
"Chapter 6 Shade management in coffee and cacao plantations", Springer Science and Business Media LLC, 1998

<1%

Publication

Newnan, Donald G.. "Engineering Economic Analysis", Oxford University Press

<1%

Tabileación

Rachmat B. SUBA, Nils G. P. BEVERIDGE, Wawan KUSTIAWAN, Geert R. DE SNOO, Hans H. DE IONGH. "Foraging ecology and diet of Bornean elephants in the Sebuku forest area, North Kalimantan Province of Indonesia: Do the choices matter? ", Integrative Zoology, 2018

<1%

Publication

Eman Hamza, Robert Sims, Günter Stolz.
"Dynamical Localization in Disordered
Quantum Spin Systems", Communications in
Mathematical Physics, 2012

<1%

15

Endang Hernawan, Mia Rosmiati, Tien Lastini, N.A. Sofiatin, Angga Dwiartama, Heri Rahman. "Agroforestry as a model of sustainable land use of small-scale private forest: a case study in Sumedang, West Java, Indonesia", International Journal of Business Continuity and Risk Management, 2020

<1%

Publication

16

Kulandaivelu Velmourougane, Rajeev Bhat. "Sustainability Challenges in the Coffee Plantation Sector", Wiley, 2017

<1%

Publication

17

Yobana A. Mariño, Maria-Eglée Pérez, Fernando Gallardo, Marella Trifilio, Michelle Cruz, Paul Bayman. "Sun vs. shade affects infestation, total population and sex ratio of the coffee berry borer (Hypothenemus hampei) in Puerto Rico", Agriculture, Ecosystems & Environment, 2016 Publication


<1%

18

B O Nababan, Y Christian, A Afandy, A Damar. "Integrated Marine and Fisheries Center and priority for product intensification in East Sumba, Indonesia", IOP Conference Series: Earth and Environmental Science, 2020

<1%

Publication



Tesfay Gidey, Tânia Sofia Oliveira, Josep Crous-Duran, João H. N. Palma. "Using the yield-SAFE model to assess the impacts of climate change on yield of coffee (Coffea arabica L.) under agroforestry and monoculture systems", Agroforestry Systems, 2019

<1%

Publication

Exclude quotes

On

Exclude matches

< 3 words

Exclude bibliography On