

Novel Amides Derivative with Antimicrobial Activity of Piper betle var. nigra Leaves from Indonesia

by Hadi.kuncoro82@gmail.com 1

Submission date: 08-Sep-2021 03:11AM (UTC-0500)

Submission ID: 1643631117




File name: molecules-26-00335_3.pdf (922.45K)

Word count: 5140

Character count: 30030

Article

Novel Amides Derivative with Antimicrobial Activity of *Piper betle* var. *nigra* Leaves from Indonesia

Fajar Prasetya ¹ , Supriatno Salam ², Agung Rahmadani ^{1,2,3}, Kansy Haikal ², Lizma Febrina ¹, Hady Anshory ⁴, Muhammad Arifuddin ¹, Vita Olivia Siregar ¹, Angga Cipta Narsa ¹, Herman Herman ¹, Islamudin Ahmad ¹ , Niken Indriyanti ¹, Arsyik Ibrahim ¹, Rolan Rusli ¹, Laode Rijai ¹ and Hadi Kuncoro ^{1,*} 

¹ Faculty of Pharmacy, Universitas Mulawarman, Samarinda 75123, Kalimantan Timur, Indonesia;

fajarprasetya@farmasi.unmul.ac.id (F.P.); agungchempharm8794@gmail.com (A.R.);

lizmafebrina@farmasi.unmul.ac.id (L.F.); marifuddin@farmasi.unmul.ac.id (M.A.);

vitaoliviasiregar@farmasi.unmul.ac.id (V.O.S.); angga@farmasi.unmul.ac.id (A.C.N.);

herman@farmasi.unmul.ac.id (H.H.); islamudinahmad@farmasi.unmul.ac.id (I.A.);

nikenindriyanti@farmasi.unmul.ac.id (N.I.); arsyik@farmasi.unmul.ac.id (A.I.);

rolan@farmasi.unmul.ac.id (R.R.); laoderijai@farmasi.unmul.ac.id (L.R.)

² Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Jatinangor, Sumedang 45363, Indonesia; supriatno.salam@gmail.com (S.S.); kansyhaikal@unpad.ac.id (K.H.)

³ Department of Chemistry Education, Faculty of Teaching and Education, Mulawarman University, Samarinda 75123, Kalimantan Timur, Indonesia

⁴ Departement of Pharmacy, Faculty of Mathematics and Natural Sciences, Islamic University of Indonesia, Jogjakarta 55584, Indonesia; hadyanshory@uii.ac.id

* Correspondence: hadikuncoro@farmasi.unmul.ac.id; Tel.: +62-541-739-491 or +62-82-158-220-088



Citation: Prasetya, F.; Salam, S.; Rahmadani, A.; Haikal, K.; Febrina, L.; Anshory, H.; Arifuddin, M.; Siregar, V.O.; Narsa, A.C.; Herman, H.; et al. Novel Amides Derivative with Antimicrobial Activity of *Piper betle* var. *nigra* Leaves from Indonesia. *Molecules* **2021**, *26*, 335. <https://doi.org/10.3390/molecules26020335>

Academic Editor: Kevin K. Schrader

Received: 11 December 2020

Accepted: 7 January 2021

Published: 11 January 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: *Piper betle* var. *nigra* is a tropical plant closely related to the common piper. *P. betle* has also been dubbed a promising source of natural antioxidants in herbal health products, antibacterial, antifungal, antimalarial, cytotoxic activity against the cancer cell lines K562 and HL-60, and antileishmanial. The aim of this study to observation Antimicrobial activity and isolation of chemical compound. The antimicrobial activity of *P. betle* extract was performed by well diffusion method against two oral pathogenic bacteria (*Streptococcus mutans* and *Streptococcus sanguinis*) and opportunistic pathogenic yeast (*Candida albicans*). The inoculum (bacterial and yeast suspension) was prepared from a 24-h culture on NB for bacterial suspension and on TSB for yeast suspension. Extraction and isolation using various method of chromatography. Isolated compounds were characterized by spectroscopic means. Our study showed antimicrobial activity from crude ethanol extract of leaves *P. betle* L. var. *nigra* against two oral pathogenic bacteria and opportunistic pathogenic yeast with concentration 0.5% and 1%. The first report of two new amides derivatives, piperenamamide A (1) and piperenamamide B (2) in *P. betle* L. var. *nigra*.

Keywords: *Piper betle* var. *nigra*; *Streptococcus mutans*; *Streptococcus sanguinis*; *Candida albicans*; piperenamamide A; piperenamamide B

1. Introduction

Piper betle a piper species have a simple profile contain very diverse suites of secondary metabolites and responsible for their use in traditional medicines to treat several disease [1]. *Piper betle* L. var. *nigra* or black betle (in Indonesia known as Sirih Hitam) is a tropical plant closely related to the common piper and belongs to the Piperaceae family and the genus of piper. This genus consists of five subgenera and approximately 1400 species spread throughout tropical and subtropical regions and widely cultivated in Indonesia, India, Sri Lanka, Malaysia, Thailand, Taiwan, and other Southeast Asian countries and has a long history of over 2000 years. This plant use for decoration and medicinal plants. *P. betle* is one of the most potent medicinal herbs that has been used over the years. In addition to the large number of beneficial properties, *P. betle* has also

been dubbed a promising source of natural antioxidants in herbal health products, antibacterial, antifungal, antimalarial, cytotoxic activity against the cancer cell lines K562 and HL-60, and antileishmanial [2–8]. Chemical composition from *Piper betle* L. var. *nigra* included caryophyllene, cadinene, γ -lactone, allyl catechol, p-cymene, eugenol methyl ether, 4-allyl resorcinol, stigmast-4-en-3,6-dione, and aristololactam A-II, and essential oils such as chavicol, chavibetol, chavabetyl acetate, eugenol, eugenyl acetate, safrole, (E) Isoeugenol, and B-caryophyllen, piperocerebrosides A and B, and amides alkaloid [8–10]. Amides as a class of typical constituents. More than 300 members of amide alkaloids are already found in species of the genus *Piper*. It seems most of them have possible bioactivity, such as antifungal, antiepileptic, antidepressant, hepatoprotective, and antiplatelet aggregation activities [11]. Isobutyl amides are one of the most frequently known classes of the amides in the plants. These amides are primarily found as long chain conjugates in *piper* genus. Arboreumine, pellitorine, fagaramide, brachystamides-C, D, E, retrofractamide-D, N-isobutyl-4-hexanoyl-hydroxypyrrolidin-1-one, (\pm)-threo-N-isobutyl-4,5-dihydroxy-2E-octaenamamide, scutifoliamide A, B, hoffmannseggiamide A, B, and cyclopiiperettine [12,13].

In this study we reported the isolation and elucidation of the chemical structures two novel amide derivatives compound and evaluated the antimicrobial activity effect of the crude ethanol extract of *Piper betle* var. *nigra* leaves from Indonesia. Antimicrobial assay to be carried out for the activity of ethanol extract from *P. betel* leaves against the activities of antimicrobial against two oral pathogenic bacteria (*Streptococcus mutans* and *Streptococcus sanguinis*) and opportunistic pathogenic yeast (*Candida albicans*).

2. Results and Discussion

2.1. Antimicrobial Activity

Table 1 shows the antibacterial activity of *P. betle* extract against two oral pathogenic bacteria and an oral opportunistic fungal. *P. betle* extract in all concentration from 0.5% to 1% inhibited growth of *S. mutans*, *S. sanguinis*, and *C. albicans* with increasing diameters of inhibition observed with increasing concentration of *P. betle* extract (Figure 1). *S. mutans* was the most sensitive strain compared to *S. sanguinis* and *C. albicans* against *P. betle* extract (0.5%) with inhibition diameters of 18.2 mm, 9.9mm, and 16.7 mm, respectively. The difference in sensitivity of these organisms is probably due to different cell types. *Candida* (yeast) cell types are different from streptococci (bacteria). This causes differences in sensitivity to *P. betle* extract. Sensitivity of *S. mutans* are more susceptible than *S. sanguinis*. The possible causes are many factors, including differences in virulence. In 2015, Azizi et al. assed the sensitivity of *S. mutans* and *S. sanguinis* to *Zingiber officinale*. They showed that *S. mutans* more susceptible than *S. sanguinis* with MIC value of 0.02 and 0.3 mg/mL, respectively [14]. In addition, their sensitivity to chlorhexidine were also reported by Medina-Flores et al. (2016), where *S. mutans* more susceptible than *S. sanguinis* with growth inhibition 23.97 mm and 19.80 mm, respectively. Alkaloid, terpenoid, flavonoid, polyphenols, tannin, and saponin compounds have been identified from *P. betle* leaf extract [15]. Flavonoids, polyphenols, and tannins are known to have antibacterial activity with at least five possible mechanisms: damage the cell membrane permeability, inhibit protein synthesis, damage the bacterial cell wall, inhibit ATP synthesis, and interfere with cell [16]. The antibacterial activity that we obtained is thought to be derived from the activity of the isolate compounds of Piperamide A and B was isolated from *Piper betle* var. *nigra*. This is as has been reported from other plants of the *Piper* genus because of the similarity in structure to the compounds we obtained although further testing is needed to ensure the strength of the antibacterial activity of these compounds [17–19]. Table 2 shows none of the tested samples of Piperamide A and B showed antimicrobial activity against *S. mutans*, *S. sanguinis*, and *C. albicans* up to 0.02%. These result from amides derivative showed the same results in tests for cyclopiiperettine on the *Piper nigrum* [13]. In recent years, *Candida* species has demonstrated resistance to many synthetic medications, indicating the need for new antifungal drugs with less side effects to treat candidiasis effectively. Several experiments used natural substances against multiresistant strains and

in particular against azole resistant candidiasis have indicated that certain species of plants have promising antimicrobial compounds from natural substances [20–22].

Table 1. Growth inhibition diameters of *Candida albicans*, *Streptococcus mutans*, and *Streptococcus sanguinis* by *Piper betle* var. *nigra* extract.

Microbial	Inhibition Diameters \pm SD (mm)	
	Extract Concentration of 0.5%	Extract Concentration of 1%
<i>Candida albicans</i>	16.7 \pm 0.8	18.1 \pm 0.7
<i>Streptococcus mutans</i>	18.2 \pm 1.8	19.6 \pm 1.9
<i>Streptococcus sanguinis</i>	9.9 \pm 1.8	12.3 \pm 1.1




Figure 1. Antimicrobial activity of *Piper betle* var. *nigra* extract. (A) Antifungal activity against *Candida albicans*; (B) Antibacterial activity against *Streptococcus mutans*; (C) Antibacterial activity against *Streptococcus sanguinis*. 1 (concentration of 1%); 2 (concentration of 0.5%).

Table 2. Growth inhibition diameters of *Candida albicans*, *Streptococcus mutans*, and *Streptococcus sanguinis* by *Piper betle* var. *nigra* isolate piperamide A and B.

Microbial	Inhibition Diameters \pm SD (mm)	
	Piperamide A	Piperamide B
	Concentration of 0.02%	Concentration of 0.02%
<i>Candida albicans</i>	-	-
<i>Streptococcus mutans</i>	-	-
<i>Streptococcus sanguinis</i>	-	-

2.2. Isolation of Crude Ethanol Extract of *Piper betle* var. *nigra*

The EtOH extract from the leaf of *P. betle* L. var. *nigra* was chromatographed over a vacuum-liquid chromatographed (VLC) column, packed with silica gel 60 by gradient elution. The VLC fractions were repeatedly subjected to normal and reverse phase column chromatography, as well as preparative TLC on silica gel GF₂₅₄ to accommodate compounds 1–2.

Piperenamide A (1) was observed as a colorless amorphous solid, with its molecular composition established as C₁₈H₂₃NO₃, based on HR-TOFMS. This showed a [M + H]⁺ ion peak at m/z 302.1749 (calcd. C₁₈H₂₄NO₃ m/z 302.1756), requiring to eight degrees of unsaturation (Figure 2). The UV spectrum showed maximum absorption at 270 and 241 nm, indicating the presence of a benzene and conjugated alkene. The IR spectrum showed bands which were ascribed to an amine (ν_{\max} 3295 cm⁻¹), amide carbonyl (ν_{\max} 1656 cm⁻¹), isolated double bond conjugated (ν_{\max} 1611 and 1504 cm⁻¹), and ether groups

(ν_{\max} 1241 cm^{-1}). Furthermore, the $^1\text{H-NMR}$ spectrum (Table 3) showed two primary methyls at δ_{H} 0.90 (6H, d, 6.7 Hz, Me-4 and 5), four sp² methine protons at δ_{H} 5.92 (1H, dd, $J = 4.6; 15.1$ Hz, H-1'), 7.02 (1H, dd, $J = 10.6; 15.1$ Hz, H-2'), 6.09 (1H, dd, 10.6; 15.2 Hz, H-3'), and 6.18 (1H, dd, 7.2; 15.2 Hz, H-4'), suggested the presence of an $\alpha, \beta, \gamma, \delta$ -unsaturated secondary amide, two trans double bonds ($J_{1',2'} = 15.1$ Hz, $J_{3',4'} = 15.2$ Hz), one cis double bond ($J_{2',3'} = 10.6$ Hz), tri-substituted benzene at δ_{H} 6.63 (1H, dd, $J = 1.5; 7.9$ Hz, *m*; o, H-6''), 6.69 (1H, d, 1.5 Hz, *m*, H-2''), and 6.69 (1H, d, 7.9 Hz, *m*, H-5''), a methylene dioxyphenyl δ_{H} 5.88 (2H, s) and some aliphatic signals in the upfield region (Supplementary Materials). Comparison of the NMR data of **1** with those of guineensine [23] shows the same amide chain but with different N-substituent positions. The N-substituent of **1** position is attached to the olefin proton resonances, this is indicated δ_{H} 3.06 (2H, d, 6.7 Hz, H-2) no correlation to δ_{H} 4.62 (NH s), δ_{H} 5.92 (1H, dd, $J = 4.6; 15.1$ Hz, H-1') and guineensine position is attached to the iso-butyl, this is indicated δ_{H} 3.16 (2H, t, 6.4; 12.9 Hz, H-1') correlation to δ_{H} 5.60 (NH brs), δ_{H} 5.76 (1H, d, 15.0 Hz, H-2). The ^{13}C NMR together with the DEPT spectra revealed eighteen carbons consisting of an amide carbonyl at δ_{C} 169.1, $\alpha, \beta, \gamma, \delta$ -unsaturated secondary amide at δ_{C} 123.3, 141.9, 142.8 and 130.2, tri-substituted benzene at δ_{C} 136.5, 109.7, 149.0, 147.9, 147.9, 108.8, and 122.3, carbon methylene dioxyphenyl at δ_{C} 102.0, two methyls at δ_{C} 20.1 (Me-4) and 20.1 (Me-5). The $^1\text{H-}^1\text{H}$ COSY spectrum of compound **1** showed correlations in H2-H3-H4 and H5, H1'-H2'-H3'-H4'-H5'-H6' and H5''-H6'', supporting the presence of a secondary amide [10]. The HMBC correlations from H-2 to C-1, C-3, C-4 and C-5, H-1' to C-1 and C-3', H-2' to C-4, H-3' to C-1 and C-5, H-4' to C-2' and C-6', H-5' to C-3' and C-1'', H-6' to C-4', C-2'' and C-6'', H-2'' to C-4', C-4'' and C-6'', H-5'' to C-1'' and C-3'', H-6'' to C-6', C-2'' and C-4'', and OCH₂O to C-3'' and 4'', which was verified by correlations observed in the $^1\text{H-}^1\text{H}$ COSY and HMBC spectra (Figure 3) Therefore, the structure of compound **1** was elucidated as the new amide and namely as piperenamide A (Figure 2).




Figure 2. Structures of piperenamide A-B (1-2).

Piperenamide B (**2**) was observed as a colorless amorphous solid, with its molecular composition established as $\text{C}_{17}\text{H}_{21}\text{NO}_3$, based on HR-TOFMS. This showed a $[\text{M} + \text{H}]^+$ ion peak at m/z 288.1567 (calcd. $\text{C}_{17}\text{H}_{22}\text{NO}_3$ m/z 288.1521), requiring to eight degrees of unsaturation (Figure 2). The UV spectrum showed maximum absorption at 270 and 241 nm, indicating the presence of a benzene and conjugated alkene. The IR spectrum showed bands which were ascribed to an amine (ν_{\max} 3235 cm^{-1}), amide carbonyl (ν_{\max} 1690 cm^{-1}), isolated double bond conjugated (ν_{\max} 1610 and 1504 cm^{-1}), and ether groups (ν_{\max} 1241 cm^{-1}). The NMR spectra of two was very similar to those of one (Table 3), except with lost of methylene at δ_{H} 2.66 (2H, t, 7.5 Hz, H-6'); δ_{C} 35.9, C-6') and methylene at δ_{H} 2.44 (2H, dd, 7.2; 7.5 Hz, H-5'), while two, methylene at δ_{H} 2.39 (2H, d, 7.2 Hz, H-5'), indicating that **2** was a demethylene derivative of one. In the HMBC spectrum, methylene correlations from H-5' to C-1'', C-2'' and C-6'', suggest methylene δ_{H} 2.39; δ_{C} 35.4, directly apply to benzene (Supplementary Materials). Therefore, leading to the structure of **2** had been elucidated as the new amide, namely as piperenamide B (Figure 2).




Figure 3. Selected HMBC and COSY correlations for piperenamide A-B (1–2).

Table 3. NMR data compound 1–2 (500 MHz for ^1H dan 125 MHz for ^{13}C).

Position Carbon	1		2	
	^{13}C -NMR δC (mult.)	^1H -NMR δC [(ΣH , mult, J(Hz))]	^{13}C -NMR δC (ppm)	^1H -NMR δC [(ΣH , mult, J(Hz))]
1	169.1 (s)	-	168.4 (s)	-
2	48.0 (t)	3.06 (2H, d, 6.7)	47.3 (t)	3.10 (2H, d, 6.7)
3	29.7 (d)	1.78 (1H, qt, 6.7)	28.9 (d)	1.82 (1H, qt, 6.7)
4	20.1 (q)	0.90 (3H, d, 6.7)	20.1 (q)	0.94 (3H, d, 6.7)
5	20.1 (q)	0.90 (3H, d, 6.7)	20.1 (q)	0.94 (3H, d, 6.7)
1'	123.3 (d)	5.92 (1H, dd, 4.6; 15.1)	122.6(d)	5.96 (1H, dd, 4.6; 15.1)
2'	141.9 (d)	7.02 (1H, dd, 10.6; 15.1)	141.2 (d)	7.12 (1H, dd, 10.6; 15.1)
3'	142.8 (d)	6.09 (1H, dd, 10.6; 15.2)	142.0 (d)	6.12 (1H, dd, 10.6; 15.2)
4'	130.2 (d)	6.18 (1H, dd, 7.2; 15.2)	129.6 (d)	6.19 (1H, dd, 7.2; 15.2)
5'	36.1 (t)	2.44 (2H, dd, 7.2; 7.5)	35.4 (t)	2.39 (2H, d, 7.2)
6'	35.9 (t)	2.66 (2H, t, 7.5)	-	-
1''	136.5 (s)	-	135.8 (s)	-
2''	109.7 (d)	6.69 (1H, d, 1.5)	109.8 (d)	6.73 (1H, d, 1.5)
3''	149.0 (s)	-	148.3 (s)	-
4''	147.9 (s)	-	146.4 (s)	-
5''	108.8 (t)	6.69 (1H, d, 7.9)	108.2 (t)	6.73 (1H, d, 7.9)
6''	122.3 (d)	6.63 (1H, dd, 1.5, 7.9)	121.6 (d)	6.68 (1H, dd, 1.5, 7.9)
OCH ₂ O	102.0 (t)	5.88 (2H, s)	101.2 (t)	5.91 (2H, s)
NH	-	4.62 (1H, br.s)	-	4.65 (1H, br.s)

3. Materials and Methods

3.1. General Experimental Procedures

UV spectra was measured using a TECAN Infinite M200 pro, with MeOH. The IR spectra and mass spectra were recorded on a SHIMADZU IR Prestige-21 in KBr and Waters Xevo QTOF MS, respectively. Using an NMR JEOL ECZ-500 and Variant Unity INOVA-500 Spectrometer (Agilent Technologies, Santa Carla, CA, USA), the NMR data was recorded at 500 MHz for ^1H and 125 MHz for ^{13}C , using TMS as internal standard. Column chromatography was conducted on the silica gel 60 (70–230 and 230–400 mesh, Merck, Kenilworth, NJ, USA), after which TLC analysis was carried out on 60 GF₂₅₄ (Merck, 0.25 mm) using various solvent systems, in order to detect spots by irradiating under ultraviolet-visible light (257 and 364 nm) and heating of silica gel plates, sprayed with H_2SO_4 in EtOH (10%).

3.2. Plant Material

The leaf of *P. betle* L. var. *nigra* were collected from Samarinda, East Kalimantan, Indonesia in June 2020. Furthermore, the plant was identified By Dr. Atik Retnowati a

staff of the Bogoriense Herbarium, Bogor, Indonesia. Finally, a voucher specimen (No. 749/IPH.1.01/If.07/VII/2020) was deposited at the Herbarium.

3.3. Extraction and Isolation

Method for extraction and isolation was referred to supriatno et al. (2018) [24]. The dried ground leaf (473.21 g) of *P. betle* L. var *nigra* was extracted with ethanol 70% (14 L), at room temperature for 7 days. After removal of the solvent under vacuum, the viscous concentrated EtOH extract (14.61 g) was obtained. The EtOH extract (14.61 g) was fractionated by column chromatography on silica gel, using a gradient of n-hexane, EtOAc and MeOH (10% stepwise) resulting into eight fractions (A–H). Fraction D (2.32 g) was subjected to column chromatography on silica gel using n-hexane-CHCl₃-EtOAc (5% stepwise), as eluting solvents to afford seven subfractions (D1–D7). Subfraction D3 (882.2 mg) was chromatographed on a column of silica gel, eluted with n-hexane: EtOAc (7:3), to give seven subfractions (D3A–D3G). Similarly, subfraction D3D (120.3 mg) was chromatographed on silica gel eluted with n-hexane: CHCl₃: EtOAc (7:2:1), to give 1 (8.2 mg). Subfraction D3E (90.1 mg) was chromatographed on silica gel eluted with petroleum ether: CHCl₃ (4.5:5.5), to give 2 (2.8 mg) [24].

3.3.1. Piperenamide A (1)

Colorless amorphous solid; mp 185–190 °C; UV (MeOH) λ_{\max} (log ϵ) 270 (4.01) and 241 (3.91) nm; IR (KBr) ν_{\max} 3295, 1656, 1611, 1504, 1490, 1443, 1141 and 1030 cm⁻¹; HR-TOFMS m/z 302.1751 [M + H]⁺, (calcd. C₁₈H₂₃NO₃ m/z 301,1749); ¹H-NMR (CDCl₃, 500 MHz), see Table 1; ¹³C-NMR (CDCl₃, 125 MHz), see Table 1.

3.3.2. Piperenamide B (2)

Colorless amorphous solid; mp 199–211 °C; UV (MeOH) λ_{\max} (log ϵ) 268 (3.91), and 219 (3.67) nm; IR (KBr) ν_{\max} 3235, 1690, 1610, 1504, 1480, 1443, 1141 and 1030 cm⁻¹; HR-TOFMS m/z 288.1567 (calcd. C₁₇H₂₂NO₃ m/z 288.1521) ¹H-NMR (CDCl₃, 500 MHz), see Table 1; ¹³C-NMR (CDCl₃, 125 MHz), see Table 1.

3.4. Antimicrobial Activity

3.4.1. Material

Liquid media for this study were nutrient broth (NB; Oxoid, Hampshire, UK) and tryptic soy broth (TSB; Merck). Solid media were Mueller Hinton Agar (MHA; Oxoid) and Tryptic soy agar (TSA; Merck)

3.4.2. Method

The antimicrobial activity of *P. betle* extract was performed by well diffusion method against two oral pathogenic bacteria (*Streptococcus mutans*, *Streptococcus sanguinis*) and opportunistic pathogenic yeast (*Candida albicans*). The inoculum (bacterial and yeast suspension) was prepared from a 24-h culture on NB for bacterial suspension and on TSB for yeast suspension. Each of inoculum was diluted with sterile physiological solution (0.9%) to 10⁸ CFU/mL (McFarland standard 0.5). 20 mL of each agar media were melted (NA media for bacterial and TSA media for yeast), cooled to 50 °C and then inoculated with 0.2 mL of the microbial suspension. The inoculated agar was poured into sterile petri dish, and then allowed to cool down on a leveled surface. Once the media had compacted, two wells were cut out of the agar, each 6 mm in diameter. Then, 30 μ L of the extract sample (concentration of 0.5 and 1%) were added into each well and incubated for 24 h at 36 °C \pm 1 °C under aerobic condition. Inhibition of microbial growth was measured in mm using SCAN 500[®] tools. Tests were performed in duplicate.

4. Conclusions

The presents study showed antimicrobial activity from crude ethanol extract of leaves *P. betle* L. var. *nigra*. The first report of two new amides derivatives, piperenamide A (1) and piperenamide B (2) in *P. betle* L. var. *nigra*.

Supplementary Materials: The following are available online. Figure S1: 1H-NMR Spectra of (1), Figure S2: 13C-NMR Spectra of (1), Figure S3: DEPT-135° Spectrum of (1), Figure S4: HMQC Spectrum of (1), Figure S5: HMBC Spectrum of (1), Figure S6: 1H-1H-COSY Spectra of (1), Figure S7: TOF MS Spectra of (1), Figure S8: 1H-NMR Spectra of (2), Figure S9: 13C-NMR Spectrum of (2), Figure S10: DEPT-135° Spectrum of (2), Figure S11: HMQC Spectrum of (2), Figure S12: HMBC Spectrum of (2), Figure S13: 1H-1H-COSY Spectra of (2), Figure S14: TOF MS Spectra of (2)

Author Contributions: Conceptualization, H.K. F.P. and L.R.; data curation and formal analysis, S.S., A.R., L.F., K.H., H.A., M.A., V.O.S., N.I., A.I., R.R.; software, A.C.N., H.H., and I.A.; validation, H.K., F.P., and L.R.; formal analysis, S.S.; investigation, H.K.; resources, F.P.; data curation, F.P., R.R.; writing—original draft preparation, H.K.; writing—review and editing, H.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the RISPRO-LPDP Research Grant from Ministry of Finance, Republic of Indonesia.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: Our thanks go to the Managing Director of the Educational Fund Management Agency of the Ministry of Finance of the Republic of Indonesia who has provided funding for Productive Innovative Research (RISPRO) according to the funding agreement number: PRJ-52/LPDP/2019 dated 22 August 2019.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Salehi, B.; Zakaria, Z.A.; Gyawali, R.; Ibrahim, S.A.; Rajkovic, J.; Shinwari, Z.K.; Khan, T.; Sharifi-Rad, J.; Ozleyen, A.; Turkdonmez, E.; et al. Piper Species: A Comprehensive Review on Their Phytochemistry, Biological Activities and Applications. *Molecules* **2019**, *24*, 1364. [[CrossRef](#)] [[PubMed](#)]
- Budiman, A.; Aulifa, D.L. A Study Comparing Antibacterial Activity of Ageratum Conyzoides, L. Extract and Piper Betle, L. Extract in Gel Dosage Forms Against *Staphylococcus Aureus*. *Pharmacogn. J.* **2020**, *12*, 473–477. [[CrossRef](#)]
- Choudhary, D.; Kale, R.K. Antioxidant and Non-Toxic Properties Of Piper Betle Leaf Extract: In Vitro And In Vivo Studies. *Phytother. Res.* **2002**, *16*, 461–466. [[CrossRef](#)] [[PubMed](#)]
- Sarkar, A.; Sen, R.; Saha, P.; Ganguly, S.; Mandal, G.; Chatterjee, M. An Ethanolic Extract of Leaves of Piper betle (Paan) Linn Mediates Its Antileishmanial Activity via Apoptosis. *Parasitol. Res.* **2008**, *102*, 1249–1255. [[CrossRef](#)] [[PubMed](#)]
- Al-Adhroey, A.H.; Nor, Z.M.; Al-Mekhlafi, H.M.; Amran, A.A.; Mahmud, R. Antimalarial Activity of Methanolic Leaf Extract of Piper Betle L. *Molecules* **2010**, *16*, 107–118. [[CrossRef](#)] [[PubMed](#)]
- Ali, I.; Khan, F.G.; Suri, K.A.; Gupta, B.D.; Satti, N.K.; Dutt, P.; Afrin, F.; Qazi, G.N.; Khan, I.A. In Vitro Antifungal Activity of Hydroxychavicol Isolated from Piper Betle L. *Ann. Clin. Microbiol. Antimicrob.* **2010**, *9*, 7. [[CrossRef](#)] [[PubMed](#)]
- Rijai, H.R.; Fakhruddin, N.; Wahyuono, S. Isolation and Identification of DPPH Radical (2,2-Diphenyl-1-Pikrylhidrazyl) Scavenging Active Compound in Ethyl Acetat Fraction of Piper Acre Blume. *Maj. Obat Tradis.* **2019**, *24*, 204. [[CrossRef](#)]
- Chen, D.-Z.; Xiong, H.-B.; Tian, K.; Guo, J.-M.; Huang, X.-Z.; Jiang, Z.-Y. Two New Sphingolipids from the Leaves of Piper Betle L. *Molecules* **2013**, *18*, 11241–11249. [[CrossRef](#)]
- Ghosh, K.; Bhattacharya, T. Chemical Constituents of Piper Betle Linn. (Piperaceae) Roots. *Molecules* **2005**, *10*, 798–802. [[CrossRef](#)]
- Tang, G.-H.; Chen, D.-M.; Qiu, B.-Y.; Sheng, L.; Wang, Y.-H.; Hu, G.-W.; Zhao, F.-W.; Ma, L.-J.; Wang, H.; Huang, Q.-Q.; et al. Cytotoxic Amide Alkaloids from Piper Boehmeriaefolium. *J. Nat. Prod.* **2011**, *74*, 45–49. [[CrossRef](#)]
- Shi, Y.-N.; Liu, F.-F.; Jacob, M.; Li, X.-C.; Zhu, H.-T.; Wang, D.; Cheng, R.-R.; Yang, C.-R.; Xu, M.; Zhang, Y.-J. Antifungal Amide Alkaloids from the Aerial Parts of Piper Flaviflorum and Piper Sarmentosum. *Planta Med.* **2016**, *83*, 143–150. [[CrossRef](#)] [[PubMed](#)]
- Kumar, V. Amides from Plants: Structures and Biological Importance. In *Studies in Natural Products Chemistry*; Elsevier: Amsterdam, The Netherlands, 2018; pp. 287–333.

13. Ren, J.; Zeng, T.; Ali, Z.; Wang, M.; Bae, J.; Chittiboyina, A.G.; Wang, W.; Li, S.; Khan, I.A. Cyclopiperettine, A New Amide from *Piper Nigrum*. *Nat. Prod. Commun.* **2017**, *12*, 1934578X1701201210. [[CrossRef](#)]
14. Azizi, A.; Aghayan, S.; Zaker, S.; Shakeri, M.; Entezari, N.; Lawaf, S. In Vitro Effect of *Zingiber Officinale* Extract on Growth of *Streptococcus Mutans* and *Streptococcus Sanguinis*. *Int. J. Dent.* **2015**, *2015*, 1–5. [[CrossRef](#)] [[PubMed](#)]
15. Junairiah, J.; Ni'matuzahroh, N.; Zuraidassanaaz, N.I.; Sulistyorini, L. Isolation and identification of secondary metabolites of black betel (*Piper betle* L. var *Nigra*). *J. Kim. Ris.* **2019**, *3*, 131. [[CrossRef](#)]
16. Tamhid, H.A. Chemical Compounds and Antibacterial Activity of *Garcinia Dulcis* (Roxb) Kurz. *J. Kedokt. Dan Kesehat. Indones.* **2019**, *10*, 71–85. [[CrossRef](#)]
17. Chanprapai, P.; Chavasiri, W. Antimicrobial Activity from *Piper sarmentosum* Roxb. against Rice Pathogenic Bacteria and Fungi. *J. Integr. Agric.* **2017**, *16*, 2513–2524. [[CrossRef](#)]
18. Mgbeahuruike, E.E.; Yrjönen, T.; Vuorela, H.; Holm, Y. Bioactive Compounds from Medicinal Plants: Focus on Piper Species. *South Afr. J. Bot.* **2017**, *112*, 54–69. [[CrossRef](#)]
19. Nascimento, S.A.; Araújo, E.A.; Da Silva, J.M.; Ramos, C.S. Chemical study and antimicrobial activities of *piper arboreum* (piperaceae). *J. Chil. Chem. Soc.* **2015**, *60*, 2837–2839. [[CrossRef](#)]
20. Sivareddy, B.; Reginald, B.; Sireesha, D.; Samatha, M.; Reddy, K.; Subrahamanyam, G. Antifungal Activity of Solvent Extracts of *Piper Betle* and *Ocimum Sanctum* Linn on *Candida Albicans*: An in Vitro Comparative Study. *J. Oral Maxillofac. Pathol.* **2019**, *23*, 333. [[CrossRef](#)]
21. Taweechaisupapong, S.; Singhara, S.; Khunkitti, W. Antimicrobial effects of *boesenbergia pandurata* and *piper sarmentosum* leaf extracts on planktonic cells and biofilm of oral pathogens. *Pak. J. Pharm. Sci.* **2010**, *9*, 224–231.
22. Donadu, M.G.; Usai, D.; Marchetti, M.; Usai, M.; Mazzarello, V.; Molicotti, P.; Montesu, M.A.; Delogu, G.; Zanetti, S. Antifungal Activity of Oils Macerates of North Sardinia Plants against *Candida* Species Isolated from Clinical Patients with *Candidiasis*. *Nat. Prod. Res.* **2020**, *34*, 3280–3284. [[CrossRef](#)] [[PubMed](#)]
23. Park, S.-D.; Oh, J.-H.; Lim, D. A New Type of Amide Formation from Thiocarboxylic Acid and Alkyl Azide. *Tetrahedron Lett.* **2002**, *43*, 6309–6311. [[CrossRef](#)]
24. Supriatno; Nurlelasari; Herlina, T.; Harneti, D.; Maharani, R.; Hidayat, A.T.; Mayanti, T.; Supratman, U.; Azmi, M.N.; Shiono, Y. A new limonoid from stem bark of *Chisocheton pentandrus* (Meliaceae). *Nat. Prod. Res.* **2018**, *32*, 2610–2616. [[CrossRef](#)] [[PubMed](#)]

Novel Amides Derivative with Antimicrobial Activity of Piper betle var. nigra Leaves from Indonesia

ORIGINALITY REPORT

0%

SIMILARITY INDEX

0%

INTERNET SOURCES

0%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

Exclude quotes On

Exclude bibliography On

Exclude matches < 64%

Novel Amides Derivative with Antimicrobial Activity of Piper betle var. nigra Leaves from Indonesia

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
