

Mineralogical Characterization of Soils In Plantation Forest of East Kalimantan, Indonesia

by Triyono Sudarmadji

Submission date: 05-Oct-2021 10:41AM (UTC+0700)

Submission ID: 1665621966

File name: article_13.pdf (158.38K)

Word count: 1412

Character count: 7995



DOI <https://doi.org/10.18551/rjoas.2021-01.13>

MINERALOGICAL CHARACTERIZATION OF SOILS IN PLANTATION FOREST OF EAST KALIMANTAN, INDONESIA

Paranoan Ria Rachel*

Faculty of Agriculture, University of Mulawarman, East Kalimantan, Indonesia

Kustiawan Wawan, Marjenah, Hartati Wahjuni, Syahrudin, Sukartiningsih

Laboratory of Budidaya Hutan, Faculty of Forestry, University of Mulawarman, East Kalimantan, Indonesia

Sudarmadji Triyono

Laboratory of KTAL, Faculty of Forestry, University of Mulawarman, East Kalimantan, Indonesia

*E-mail: rachelria_paranoan@yahoo.co.id

ABSTRACT

Soil minerals including primary mineral and (secondary) clay mineral play an important role which has greatly contributed in determining the properties of soils and availability of soil nutrient reserve in soil. The objective of the study was to identify of clay minerals and primary minerals (sand fraction) at forest plantation area in East Kalimantan, Indonesia. The mineralogical composition of fine sand fraction of soils was identified using a polarizing microscope (PM). The mineralogical composition of clay fractions was determined using X-ray diffraction analysis (XRD). The research plot is circular with an area of 0.05 ha and a radius of 12.6 m. Four soil sample from four plots collected for soil mineral analysis. The results showed that the clay fraction in plot A 45 has moderate amounts of kaolinite and vermiculite, trace amounts of goethite and illite. Plot A 34 was dominated by kaolinite, small amounts of vermiculite, trace amounts of illite. Plot A 36 was dominated by kaolinite, small amounts of vermiculite and illite, trace amounts of goethite. Plot A 33 was dominated by kaolinite small amounts of vermiculite and quartz. Compositions of Sand fraction at study area were dominated by resistant minerals (quartz).

KEY WORDS

Quartz, vermiculite, primary minerals, illite, clay minerals, kaolinite, industrial plantation forest.

Minerals are the main constituent elements of soil and play an important role in determining the chemical and physical properties of soil (Prastyo et al. 2004; Anda, et al. 2012). Minerals in the soil can be differentiated into primary minerals (fine sand fraction) minerals and secondary minerals or minerals from clay fraction (Prasetyo et al. 2004).

In general, the management of plants that continuously produce wood with fast growing plants that require a lot of nutrients will increase the depletion of large amounts of nutrients, this is closely related to land productivity, including efforts to increase land fertility by recognizing its potential presence nutrient reserve through the identifications of primary minerals and clay (secondary) minerals which greatly affect soil properties. So that it can help in formulating strategies for increasing soil productivity in industrial plantation forest location.

To determine the level of nutrient reserves of a soil type it is necessary to analyze primary mineral composition from the soil. Considering the importance of clay minerals, it is necessary to know the type of dominant clay minerals in order to further interpret the potential of soil resources and land capabilities (Munir, 1995).



Based on research studied, the clay mineral has been identified in East Kalimantan (Ohta, et al., 1992; Prasetyo et al., 2001). Nevertheless, there is still little information of mineral found in plantation forest industry, Sebulu Subdistrict, East Kalimantan.

The purpose of this study was to identify primary minerals and clay minerals at plantation forest industry in East Kalimantan, Indonesia.

METHODS OF RESEARCH

The study was conducted at Industrial Plantation Forest, Sebulu-sub distric, East Kalimantan, Indonesia from March to July 2020. The research plot is circular with an area of 0.05 ha and a radius of 12.6 m (Syahrudin, 1997; Syahrudin, 2005; Mackenzen, 1998). Soil samples were taken from 4 plots for primary mineralogical analysis consists of 500 g from each sample form selected horizon and for clay mineral analysis consists of 500 g from each sample from selected horizon. The mineralogical compositions of sand fractions of soils were analyzed at using a polarizing microscope (PM). Mineralogical analysis to determine clay mineral composition was carried out using X-ray diffractometer (XRD), clay treatments, i.e., Mg then heated 550°C.

RESULTS AND DISCUSSION

Based on sand fraction (primary minerals) analysis using polarizing microscope, the composition of primary minerals in sand fraction of soils is given in Table 1. The resistant minerals at plot A45 was dominated by quartz (96%). Composition of sand fraction minerals at plot A34 was dominated by quartz mineral (85%), plot A36 was dominated by quartz mineral (72%) and plot A33 was dominated by quartz mineral (92%). This indicates that the dominance of quartz minerals means that there is no potential for nutrient reserves. If the soil is dominated by quartz (resistant mineral), the soil is considered poor in nutrients (Tafakresnanto and Prasetyo, 2001).

Table 1 – Mineralogical Composition of fine sand fractions (50 -250 µm)

Plot	Op	Zi	Qt	Wm	Rf	Konkresi Besi	Lm	An	Lb	Bw	An	Or	Hb	Vg	Total
A45	sp	3	96	Sp	1	-	-	-	-	-	-	-	-	-	100
A34	9	4	85	Sp	2	sp	-	sp	-	-	-	-	-	-	100
A36	3	3	72	17	-	5	-	-	-	-	-	-	-	-	100
A33	2	-	92	1	5	-	-	sp	-	-	-	-	-	-	100

Note:Op=opaque, Zi=zircon, Qt=quartz, Wm=weathered mineral, Rf=rock fragment, Lm=limonite, An=anatas, Lb=labradorite, Bw=bytownite, An=anorthite, Or=orthoclase, Hb=hornblende, Vg=volcaic glass, sp=sporadic, - =not detected.




Figure 1 – X-ray diffraction patterns of the clay fraction of the soil treated with Mg



The XRD patterns of clay fraction are illustrated in Fig.1. The XRD patterns of clay fractions from selected horizon of four plots indicated the presence of kaolinite, vermiculite, illite minerals in soils. The kaolinite mineral is characterized by the peak at 7.17 Å on Mg saturated treatment. The peak of kaolinite and goethite minerals disappeared after heating treatment at 500°C.




Figure 2- X-ray diffraction patterns of the clay fraction of the soil heated 550°C

Table 2 – Composition of clay fraction

No	Plot	Smectite	Kaolinite	Goethite	Vermiculite	Illite	Quartz
1	A45	-	++	(+)	++	(+)	-
2	A34	-	+++	-	+	(+)	-
3	A36	-	+++	(+)	+	+	-
4	A33	-	+++	-	+	-	+

Note: +++ = predominant, ++ = dominant, ++ = moderate, + = small, (+) = trace - = not detected.

Clay minerals (secondary minerals) of soils all plots presented in Table 2. There are moderate amounts of kaolinite and vermiculite, trace amounts of goethite and illite in plot A45. Composition of clay mineral in plot A34 dominated by kaolinite, small amounts of vermiculite and trace amounts of illite. The clay fraction of plot A36, dominated by kaolinite mineral, small amounts of vermiculite and illite, and trace amounts of goethite. Composition of clay mineral at plot A33 dominated by kaolinite, and small amounts of vermiculite and quartz.

CONCLUSION

Clay mineral compositions presented in all plots. Clay minerals composition of soils was dominated by kaolinite minerals. The dominance of resistant minerals at study area in forest plantation, sub-district Sebulu, East Kalimantan dominated by quartz mineral in all plots. indicated that the soils have no potential reserved nutrients or soil has low fertility.

REFERENCES

1. Anda, M., Mulyani, A., Surapto. 2012. Mineralogical and Characterization and Chemical Properties of Soils as a Consideration for Establishing Sustainable Soil Management Strategies. Indonesian Journal of Agricultural Science 13(2):54-67.



2. Mackensen, J. 1998. Untersuchung zur nachhaltigen Nährstoffversorgung in schnellwachsenden Plantagensystemen in Ost-Kalimantan, Indonesien-ökologische und ökonomische Implikationen. Göttinger Beiträge zur Land-und Forstwirtschaft in den Tropen und Subtropen. 127.
3. Munir, M. 1995. Geologi & Mineralogi Tanah. Pustaka Jaya. ISBN: 979-419-146-9.
4. Prasetyo, B.H., J.S. Adiningsih, K. Subagyo, & R.D.M. Simanungkalit. 2004. Mineralogi, Kimia, Fisika and Biologi Lahan Sawah. Hlm. 36-50. Dalam F. Agus (Wd.) Lahan Sawah and Teknologi Pengelolaanya. Pusat Penelitian Tanah and Agroklimat, Bogor.
5. Prasetyo, B.H., Suharta, N., Subagyo, H., Hikmatullah. Chemical and Mineralogical Properties of Ultisols of Sasamba Area, East Kalimantan. Indonesian Journal of Agricultural Science 2(2): 37-47.
6. Pramuji & Bastaman, M. (2009). Teknik Analisis Mineral Tanah Untuk Menduga Cadangan Sumber Hara. Buletin Teknik Pertanian. Vol. 14. No. 2. 80-82.
7. Ohta, S., Effendi, S., Tanaka, N., Miura S. 1992. Characteristics of Major Soils Under Lowland Dipterocarp Forest in East Kalimantan, Indonesia. PUSREHUT Special Publication No.2.
8. Syahrudin. 1997. The Role of Undergrowth on Timber Estate of Eucalyptus deglupta in East Kalimantan. [Thesis]. The Master of Science Degree in Tropical Forestry and Ecology Georgia Augusta University, Göttingen. [Germany].
9. Syahrudin. 2005. The potential of oil palm and forest plantations for carbon sequestration on degraded land in Indonesia. [Dissertation]. Zentrum für Entwicklungsforschung Center Development Research, University of Bonn. [Germany].
10. Tafakresnanto, C. and B.H. Prasetyo. 2001. Peranan data mineral tanah dalam menunjang interpretasi sumber daya tanah. Jurnal Tanah and Air 2(1): 47-56.

Mineralogical Characterization of Soils In Plantation Forest of East Kalimantan, Indonesia

ORIGINALITY REPORT

6%

SIMILARITY INDEX

0%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** I. Unamba-Oparah, M.J. Wilson, B.F.L. Smith. "Exchangeable cations and mineralogy of some selected Nigerian soils", *Applied Clay Science*, 1987 2%
Publication
 - 2** I Suryani, C Lopulisa, A Ahmad. "The Potential estimation of soil fertility based mineral types in Papalang areas, Mamuju Regency, West Sulawesi", *IOP Conference Series: Earth and Environmental Science*, 2021 2%
Publication
 - 3** Anda, Markus, and Djadja Subardja. "Assessing soil properties and tidal behaviors as a strategy to avoid environmental degradation in developing new paddy fields in tidal areas", *Agriculture Ecosystems & Environment*, 2013. 2%
Publication
-
-

Exclude quotes On

Exclude bibliography On

Exclude matches < 10 words