

## **Phytochemical, TLC Profile, and Antioxidant Activity of Malinau Endemic Plant of Tabar Kedayan (*Aristolochia papilifolia* Ding Hou) Root Fractions**

Islamudin Ahmad\*<sup>1</sup>, Mirhansyah Ardana<sup>1</sup>, Riski Sulistyarini<sup>1</sup>, Wisnu Cahyo Prabowo<sup>2</sup>, Muhammad Arifuddin<sup>2</sup>

Departement of Pharmaceutical Sciences, Faculty of Pharmacy, Mulawarman University, Samarinda, East Kalimantan, Indonesia, 75116

Laboratory of Research and Depelovment of Farmaka Tropis, Faculty of Pharmacy, Mulawarman University, Samarinda, East Kalimantan, Indonesia, 75116

**Abstract :** Tabar Kedayan (*Aristolochia papilifolia* Ding Hou) is Malinau endemic plant as a traditional medicine by Dayak community. However, the scientific data from this plant is limited. This study aimed to know the secondary metabolite groups, TLC profile and antioxidant activity in the fraction of this plant. The dried sample (one kg) was macerated with ethanol for 2 x 24 hour. The obtained extract (80 gram) was partitioned using liquid-solid extraction with ethyl acetate, then ethyl acetate soluble fraction (22.4 gram) was conducted fractionation using vacuum liquid chromatography. Each fraction was analyzed of preliminary phytochemical, TLC profile, and antioxidant activity (qualitative and quantitative). Based on the results, had obtained data, such as preliminary phytochemical was showed the groups of alkaloid and steroid were present in fractions A, B, and C. Phenol groups was present only in the fraction F, while the fraction C, D, E, and F was present flavonoid. The TLC profile had demonstrated the ability of fluorescent components of based on the different properties of compounds contained in each fraction. Antioxidant activity determination of ethyl acetate soluble fraction from Tabar Kedayan (*A. papilifolia* Ding Hou) roots using DPPH (2,2-diphenyl-1-picrylhydrazyl) method with a qualitative activity (TLC autography method) was fraction C, and a quantitative activity (IC<sub>50</sub> value) was 139.11 µg/ml.

**Keywords:** *Aristolochia papilifolia* Ding Hou, preliminary phytochemical, 2,2-diphenyl-1-picrylhydrazyl, IC<sub>50</sub>, TLC autography.

### **Introduction**

Malinau is one of the districts in North Kalimantan Province, Indonesia. Most areas of Malinau is still a primary forest with abundant biodiversity. One of the species used as medicine by the local community derived from the genus of *Aristolochia* (family *Aristolochiaceae*)<sup>1</sup>.

The genus of *Aristolochia* comprises about 500 species, and the species of this genus has efficacy to be used as a medicine. This genus is a vines shrub and most widely spread in Asia, Africa, North and South America<sup>2,3,4</sup>. Many plants of this genus have efficacy as an anti-asthmatic, anti-inflammatory<sup>5</sup>, anticancer, analgesic, infections diseases, cardiovascular<sup>6</sup>, nephrotoxicity<sup>7</sup>, antioxidant, antimicrobial, and insect repellent or antiphenom<sup>8,9</sup>. Some research was reported from this genus to contain secondary metabolites such as

flavonoid<sup>5</sup>, alkaloid<sup>10,11</sup>, tannic acid, fixed oil, nitrophenanthrene carboxylic acid, resin<sup>12</sup>, and aristolochic acids<sup>7</sup>.

Tabar Kedayan (*A. Papilifolia*) is a Malinau endemic plant widely traditionally used by Dayak community as a traditional medicine, such as detoxifying, neutralizing venom (particularly the insect and snake), and all sorts of venomous animal bites, food poisoning, diarrhea, toothache and can also be used to neutralizing alcohol in the human body (alcoholic). However, scientific data from this species (mainly secondary metabolites and activity potential) has not reported. Therefore, this study aimed to know the secondary metabolite groups, TLC profile and antioxidant activity in the fraction of Tabar Kedayan (*A. Papilifolia*).

## Experimental

### Material

#### Plant

Tabar Kedayan (*A. Papilifolia*) root was collected from Malinau, North Kalimantan, Indonesia. Moreover, this plant was identified at Laboratory of Dendrology, Faculty of Forestry, Mulawarman University, Samarinda, East Kalimantan, Indonesia and the voucher specimen was deposited at Pharmaceutical Research & Development Laboratory of FARMAKA TROPIS, Faculty of Pharmacy, Mulawarman University, Samarinda, East Kalimantan, Indonesia.

#### Chemical and equipment

The chemical was used in this study such as; aqua DM, n-hexane, ethyl acetate, ethanol were purchased from SmartLab Indonesia, Indonesia. DPPH (2,2-diphenyl-1-picrylhydrazyl) was purchased from Sigma-Aldrich, Germany. Silica gel GF 60, silica gel GF 254 and TLC plate were purchased from Merck, Germany. The equipment was used such as; Water bath (Memmert<sup>®</sup>), the rotary evaporator (Buchi<sup>®</sup>), Chamber (Camaq<sup>®</sup>), hot plate (Stuart<sup>®</sup>), vortex (Health<sup>®</sup>), and Spectrophotometry UV-Visible Double Beam (Dynamica<sup>®</sup>).

#### Extraction and Fractionation Process

The extraction process carried out by extracting The dried sample (1 kg) using maceration method with ethanol based on standard conventional extraction procedures. The extracts obtained (80 grams) then partitioned using solid-liquid extraction methods with ethyl acetate. The ethyl acetate soluble extract (22.4 grams) carried out fractionation using a vacuum liquid chromatography method with a solvent combination (n-hexane and ethyl acetate) in a ratio of 5: 5, 4: 5, 3: 5, 2: 5, 1: 5, and 0: 5, respectively.

#### Phytochemical Screening

Each fraction was analyzed using chemical reagent for preliminary analysis of various groups of phytoconstituents such as alkaloids, flavonoids, steroid, phenolic, and polyphenols by using standard procedure<sup>13,14,15</sup>.

#### TLC Profile

The TLC profile determination performed based on the literatures<sup>14</sup>, with slight modification. The stationary phase using TLC plate (10x20 cm) and the mobile phase using an eluent comparison based on the polarity levels was used in this study. The retention factor (Rf) value was a data of TLC profile.

#### Antioxidant Activity


Determination of antioxidant activity conducted using DPPH free radical scavenging assay (qualitative and quantitative assay) as the standard procedures<sup>16,17,18,19</sup> with slight modification. For qualitative activity, each solution of extract sample was spotted (5 x 2 µg/ml) d on TLC-plates (silica gel thin layer chromatography) and were spread using a mobile phase (appropriate solvent systems). The plates developed with 0.1 % DPPH in methanol and placed in the dark condition for 30 minutes, and the color changes (yellow on purple background) were noted. For quantitative activity, the sample solution was used for experimental at 30 times dilution. The reaction mixture contained two mL of 40 µg/mL DPPH and two mL of sample solution. The DPPH free radical scavenging activity of the sample was quantified using spectrophotometry at 517 nm wavelength. Results were reported as the IC<sub>50</sub> (the amount of antioxidant necessary to decrease by 50% the initial DPPH concentration).

## Results and Discussion

### Extraction and Fractionation Processes

Tabar Kedayan (*Aristolochia papilifolia* Ding Hou) is a native plant of Kalimantan, particularly Malinau region. This plant is a snub, root russet, edged, elongated and usually grown in the tropical rainforests. The local communities take advantage of this plant as a traditional medicine from generation to generation. The roots can be used in the form of thin slices and mixed with warm water.

The fresh samples sorted, washed, chopped, and dried. The dried samples were extracted by maceration using ethanol. Subsequently, the crude extract was partitioned using solid-liquid extraction method using ethyl acetate. The ethyl acetate soluble extract (the ethyl acetate extract) was fractionated with vacuum liquid chromatography method using n-hexane: ethyl acetate eluent (a ratio of 5: 5, 4: 5, 3: 5, 2: 5, 1: 5, and 0: 5, respectively). Extraction and fractionation process (shown in **Figure 1**).



Where: A was the dried root sample of Tabar Kedayan (*Aristolochia Papilifolia* Ding Hou); B was maceration processes; C was vacuum liquid chromatography procedures; D was the prepared sample ready for analysis; E was the TLC spot at 254 nm wavelength, and F was the TLC spot at 366 nm wavelength.

**Figure 1** Extraction and Fractionation processes

### Phytochemical analysis

Preliminary phytochemical was performed on all the ethyl acetate soluble fraction including fraction A, B, C, D, E, and F. The content of secondary metabolites examined, namely alkaloids, flavonoids, steroids, polyphenols, and phenolic (shown in **Table 1**).

**Table 1** Results of preliminary phytochemical of ethyl acetate soluble fraction

No	Secondary Metabolite	Ethyl Acetate Soluble Fraction					
		A	B	C	D	E	F
1	Alkaloid	(+)	(+)	(+)	(-)	(-)	(-)
2	Polyphenolic	(-)	(-)	(-)	(-)	(-)	(-)
3	Flavonoid	(-)	(-)	(+)	(+)	(+)	(+)
4	Steroid	(+)	(+)	(+)	(-)	(-)	(-)
5	Phenolic	(-)	(-)	(-)	(-)	(-)	(+)

where: (+)= Present, (-) = Absent

Based on **Table 1** shows that the groups of alkaloid and steroid were present in fractions A, B, and C. Phenol groups was present only in the fraction F, while the fraction C, D, E, and F was present flavonoid.

### TLC Profile

TLC profile of the ethyl acetate soluble fraction (including fraction A, B, C, D, E, and F) shown in **Table 2**. TLC profile at UV 254 nm wavelength was detected spots are different in each of the fractions, namely four spots (fraction A), seven spots (fraction B), one spot (fraction C), one spot (fraction D and E), and two spots (fraction F). Profile TLC at UV 366 nm wavelength was also present with the spot differently on each of the fractions, namely three spots (fraction A and F), five spots (fraction B and C), seven spots (fraction D), and six spots (fraction E). Moreover, profiles TLC at visible wavelength using 10% H<sub>2</sub>SO<sub>4</sub> spray was detected four spots (fraction B), three spots (fraction A, C, and D), and one spot (fraction E and F).

**Table 2 TLC Profile of Ethyl Acetate Soluble Fraction**

Ethyl acetate soluble fraction	Total Spot and Rf value			
	Spot	UV (254nm)	UV (366nm)	H <sub>2</sub> SO <sub>4</sub> Spray
Fraction A (5:5)	1	-	0,51	-
	2	-	0,62	0,62
	3	0,74	-	0,69
	4	0,82	-	-
	5	0,85	-	-
	6	0,92	0,92	0,92
Fraction B (4:5)	1	0,28	-	-
	2	0,32	0,31	-
	3	-	0,38	-
	4	-	0,51	-
	5	-	0,62	0,62
	6	-	0,69	0,69
	7	0,74	-	0,74
	8	0,82	-	-
	9	0,85	-	-
	10	0,92	-	-
	11	0,97	-	0,97
Fraction C (3:5)	1	-	0,54	-
	2	-	0,71	-
	3	-	0,77	0,77
	4	-	-	0,88
	5	0,92	-	-
	6	-	0,97	0,95
	7	-	0,98	-
Fraction D (2:5)	1	0,31	0,31	-
	2	-	0,46	-
	3	-	-	0,49
	4	-	0,54	0,54
	5	-	-	0,62
	6	-	0,71	-
	7	-	0,77	-
	8	-	0,97	-
	9	-	0,98	-
Fraction E (1:5)	1	0,31	0,31	-
	2	-	0,51	0,49
	3	-	0,57	-
	4	-	0,71	-
	5	-	0,77	-
	6	-	0,98	-


Fraction F (0:5)	1	0,38	0,34	-
	2	0,52	-	0,54
	3	-	0,66	-
	4	-	0,85	-

In UV 254 nm, the TLC plates will fluoresce while the sample will appear dark because the power of interaction between UV light with a fluorescence indicator contained on the TLC plates. Fluorescence is the emission of light emitted by the components contained in each spot when the excitation of the electrons undergo basic energy level to a higher energy level and then return to its original state while releasing energy. While at UV 366 nm has the opposite fluoresce because the power of interaction between UV light with chromophore groups are bound by auxochromes that existed at the spots. Meanwhile, the detection principle of a spot at a visible wavelength using 10% H<sub>2</sub>SO<sub>4</sub> spray reagent was based on the ability of reagents damage the chromophore groups of spot components, so that the wavelength will be shifted toward the longer (UV into VIS) and the spot components become visible. The ability of fluorescent components based on the different properties of compounds contained in each fraction.

## Antioxidant Activity

### 1. Qualitative antioxidant activity using TLC autography methods

In the present study, a qualitative test for the antioxidant activity of ethyl acetate soluble fractions using DPPH method. Determination of antioxidant activity using DPPH method is a simple, rapid method and does not require many reagents. The measurement results with DPPH samples demonstrated the ability of antioxidants in general and not based on the radical inhibitory. The spots of each fraction were sprayed using 0.1% DPPH solution, with a presence of yellow spots whitish after 30 minutes showed a positive antioxidant. The results showed the fraction that has antioxidant activity was a fraction C, E, and F (shown in **Figure 2**). At the fraction C, the presence of round yellow spots at the center of the TLC plate, the results indicate that the separation of compounds in this fraction showed the best separation of compound groups.




Where a is a positive reaction of DPPH free radical scavenging with the color changes of spot (yellow on purple background) meant that sample has an antioxidant activity

**Figure 2** Results of the antioxidant activity of the ethyl acetate soluble fraction after spraying 0.1 % DPPH (only shown the fraction has a positive reaction)

### 2. Quantitative antioxidant activity using determination of IC<sub>50</sub> of DPPH free radical scavenging

The antioxidant activity determination with DPPH free radical scavenging was used fraction C. The IC<sub>50</sub> value was obtained using linear regression equation that states the relationship between the sample concentration with the symbol (X) and the DPPH free radical scavenging activity with the symbol (Y) of a

measurements replication series. The equation result was obtained  $Y = 0.292X + 9.38$  with coefficient correlation ( $R^2$ ) = 0.998 (shown in **Figure 3**). A substance (active compound) has the antioxidant properties when the  $IC_{50}$  value of less than 200  $\mu\text{g/ml}$ <sup>18,20</sup>. Antioxidant activity determination of ethyl acetate soluble fraction (fraction C) from Tabar Kedayan (*A. papilifolia* Ding Hou) roots using DPPH method with  $IC_{50}$  value was 139.11  $\mu\text{g/ml}$  (showed in **Table 3**). At a concentration of  $\mu\text{g/ml}$  demonstrated the concentration of ethyl acetate soluble fraction could inhibit oxidation process by 50 percent.



**Figure 3** Effect of antioxidant activities against concentration of ethyl acetate soluble fraction

**Table 3** Results of measurement and determination of a DPPH free radical scavenging assay ( $IC_{50}$ ) from ethyl acetate soluble fraction (fraction C)

No.	Concentration	Approximately absorbance	DPPH free radical scavenging activity (%)	Linear regression equation	$IC_{50}$ ( $\mu\text{g/ml}$ )
1	10 ppm	0,339	7,95	Y= 0,292x + 9,38 $R^2= 0,98$	<b>139,11</b>
2	50 ppm	0,276	25,12		
3	100 ppm	0,204	44,55		
4	150 ppm	0,168	54,29		
5	200 ppm	0,133	63,84		

## Conclusion

Based on the above results, the fraction C was an active fraction from Tabar Kedayan (*A. papilifolia* Ding Hou) roots and was analyzed (including preliminary phytochemical and TLC profile). The results are preliminary data related to the potential as a source of natural medicine and are first information to explore these plants in further study.

## Acknowledgement

This study was supported by grants via "HIBAH KELOMPOK PENELITI SAINS FARMASI 2015." Faculty of Pharmacy, Mulawarman University, Samarinda, East Kalimantan, Indonesia (1909.4/UN17.12/LT/2015).

### Conflict of Interest

We declare that we have no conflict of interest

### References:

1. Supriningrum R, Jubaidah S, Sapri. Acute toxicity test of extract *Aristolochia foveolata* Merr. roots. *Media Sains*. 2014;7(2):247-251.
2. Santander R, Urzúa A, Olguín Á, Sánchez M. Temporal Variation of *Aristolochia chilensis*. Aristolochic Acids during Spring. *Molecules*. 2015;20(12):20391-20396.
3. Polevova S V. Ultrastructure and development of sporoderm in *Aristolochia clematitis* (Aristolochiaceae). *Rev Palaeobot Palynol*. 2015;222(3):104-115.
4. Dhouioui M, Boulila A, Chaabane H, Zina MS, Casabianca H. Seasonal changes in essential oil composition of *Aristolochia longa* L. ssp. *paucinervis* Batt. (Aristolochiaceae) Roots and its antimicrobial activity. *Ind Crops Prod*. 2016;83:301-306.
5. Machado MB, Lopes LMX. Tetraflavonoid and biflavonoids from *Aristolochia ridicula*. *Phytochemistry*. 2008;69(18):3095-3102.
6. Heinrich M, Chan J, Wanke S, Neinhuis C, Simmonds MSJ. Local uses of *Aristolochia* species and content of nephrotoxic aristolochic acid 1 and 2-A global assessment based on bibliographic sources. *J Ethnopharmacol*. 2009;125(1):108-144.
7. Huang T-C, Chen S-M, Li Y-C, Lee J-A. Increased renal semicarbazide-sensitive amine oxidase activity and methylglyoxal levels in aristolochic acid-induced nephrotoxicity. *Life Sci*. 2014;114(1):4-11.
8. Li M, Au KY, Lam H, Cheng L, But PPH, Shaw PC. Molecular identification and cytotoxicity study of herbal medicinal materials that are confused by *Aristolochia* herbs. *Food Chem*. 2014;147:332-339.
9. Li ZJ, Njateng GSS, He WJ, et al. Chemical composition and antimicrobial activity of the essential oil from the edible aromatic plant *Aristolochia delavayi*. *Chem Biodivers*. 2013;10(11):2032-2041.
10. Ferreira MLR, de Pascoli IC, Nascimento IR, Zukerman-Schpector J, Lopes LMX. Aporphine and bisaporphine alkaloids from *Aristolochia lagesiana* var. *intermedia*. *Phytochemistry*. 2010;71(4):469-478.
11. Guillén AP, Michel FJS, Amezcua JMR, et al. Four new species of *Aristolochia pentandrae* from Western Mexico. *Syst Bot*. 2016;41(1):128-141.
12. Bhattacharjee P, Bhattacharyya D. Characterization of the aqueous extract of the root of *Aristolochia indica*: Evaluation of its traditional use as an antidote for snake bites. *J Ethnopharmacol*. 2013;145(1):220-226.
13. Sumithra M, Arunachalam G. Pharmacognostical study and phytochemical evaluation of Agardh, *Sargassum ilicifolium* Turner C. *Int J PharmaTech Res*. 2014;6(7):2022-2027.
14. Ahmad I, Arifuddin M, Rijai L. The effect of extraction methods of Bawang Dayak (*Eleutherine palmifolia* L. MERR) against TLC profiles and sunscreen activities. *Int J PharmTech Res*. 2016;9(9):428-436.
15. Anupama N, Madhumitha G. Pharmacognostic standardization and anti-microbial studies of dried *Carissa carandas* fruits. *Int J PharmaTech Res*. 2015;8(8):206-210.
16. Fuad Hafid A, Ismail, Wardiyanto S, Tumewu L, Rahman A, Widyawaruy Anti A. Free-radical scavenging activity screening of some Indonesian plants. *Int J Pharm Pharm Sci*. 2014;6(6):115-117.
17. Lee Wei HAR, Ismail IS, others. Antioxidant activity, total phenolics and total flavonoids of *Syzygium polyanthum* (Wight) Walp leaves. *Int J Med Aromat Plants*. 2012;2(2):219-228.
18. Ahmad I, Rijai L, Sulistiarini R. Antioxidant Activity of Some Selected East Borneo Plants. *Int J Public Health Sci*. 2015;4(1):58-62.
19. Shalaby E, Shanab S. Antioxidant compounds, assays of determination and mode of action. *African J Pharm Pharmacology*. 2013;7(10):528-539.
20. Molyneux P. The use of the stable free radical diphenyl picrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarinn J Sci Technol*. 2004;26(2):211-219.

\*\*\*\*\*