

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/351358469>

# Antioxidant Activity of Secondary Metabolite Compounds from Lichen *Teloschistes flavicans*

Research · March 2021

DOI: 10.33263/BRIAC116.1387813884

CITATIONS

0

READS

32

6 authors, including:



**Ahmad Darmawan**  
Research Center for Chemistry

48 PUBLICATIONS 173 CITATIONS

SEE PROFILE



**Usman Usman**

6 PUBLICATIONS 7 CITATIONS

SEE PROFILE



**Muhammad Nurdin**  
Universitas Haluoleo

86 PUBLICATIONS 557 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Natural Product Chemistry [View project](#)



natural products chemistry and drug discovery from natural products [View project](#)

# Antioxidant Activity of Secondary Metabolite Compounds from Lichen *Teloschistes flavicans*

Maulidiyah Maulidiyah<sup>1,\*</sup>, Akhmad Darmawan<sup>2</sup>, Usman Usman<sup>3</sup>, Andi Musdalifah<sup>1</sup>, La Ode Agus Salim<sup>1</sup>, Muhammad Nurdin<sup>1</sup>

<sup>1</sup> Department of Chemistry, Faculty of Mathematics and Natural sciences, Universitas Halu Oleo, Kendari 93231, Southeast Sulawesi – Indonesia

<sup>2</sup> Research Center for Chemistry, Indonesian Institute of Sciences, Kawasan Puspitek, Serpong 15314, Banten – Indonesia

<sup>3</sup> Post Graduate of Chemistry Education Program, Faculty of Teacher Training and Education, Universitas Mulawarman, Samarinda 75119, East Kalimantan – Indonesia

\* Correspondence: maulid06@yahoo.com (M. M.);

Scopus Author ID 56648049800

Received: 16.01.2021; Revised: 15.02.2021; Accepted: 18.02.2021; Published: 1.03.2021

**Abstract:** In the last decade, lichen has become a research interest related to exploring secondary metabolites compounds. Lichen is a unique organism and contains specific chemical compounds rarely found in other natural sources; one species is the lichen *Teloschistes flavicans*. The objective study explores the toxicity effects and antioxidants activity properties from lichen *Teloschistes flavicans* isolate. The isolation of secondary metabolites was carried out by extraction, separation, and purification by means of gravity column chromatography, thin-layer chromatography, and purity tests. Toxicity activity test used the Mayer method by looking at the LC<sub>50</sub> value, while the antioxidant activity test used the DPPH method by looking at the IC<sub>50</sub> value. The results of the isolation obtained a needle-shaped crystal with the name of the compound is 3-[1'-(2",3"-dihydroxy-phenyl)-propyl]-7-hydroxy-chroman-4-one and the molecular formula C<sub>18</sub>H<sub>18</sub>O<sub>5</sub>. The toxicity test showed that extract and isolate compounds have LC<sub>50</sub> values of 9.38 and 162.18 µg/mL, indicating that the isolated compound was toxic. The antioxidant test showed that the extract and isolated compounds have antioxidant activity with IC<sub>50</sub> values of 54.05 and 127.38. The results of these studies provide scientific development of natural compounds that have the potential to act as antioxidants for diseases in the human body.

**Keywords:** lichen; isolation; DPPH; toxicology, antioxidant.

© 2021 by the authors. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The unique lichen organism is very interesting to study because it has many benefits, such as medicinal ingredients, textile materials, perfume ingredients, and an indicator of a polluted environment. Lichen is a unique organism formed from two different organisms: fungi and algae [1–3]. Lichen produces secondary metabolites of the depsides, depsidone, dibenzofuran, xanthone, and terpenes, which are generally only found in lichen organisms [4–6]. Secondary metabolite compounds are currently a concern for researchers to study more seriously. The use is widely used in the health sector to seek remedies from natural ingredients. The secondary metabolite compounds of lichen have active substances with biological activity that can be developed as medicinal substances [7–10]. Several biological activities have been reported, namely antioxidant [11,12], antimalarial [13], antimicrobial [14,15], anticancer [16,17], antidiabetic [18], and anti-inflammatory [19].

One of the lichens that have the potential to be developed is the lichen *Teloschistes*. Lichen *Teloschistes* have been used by the general public as traditional medicine [20]. The pharmacological activity provided by lichen depends on the secondary metabolites contained therein. Lichen *Teloschistes flavicans* is a lichen species known as 'Golden Hair Lichen' [21–23]. Its bright color is one of the identifying identities of this lichen. The metabolite compounds produced by lichen *T. flavicans* need further investigation.

This research studies the active compounds of lichen *T. flavicans* by looking at their biological activity against antioxidants. Oxidative stress in the body is the beginning of degenerative diseases, such as hypertension, heart disease, diabetes, stroke, and cancer [24,25]. Antioxidant testing becomes necessary to report, given the lifestyle of people who are less concerned with health. The human body can provide defense by producing its own antioxidant compounds [26–28]. Defenses that are not optimal cause these healthy cells to be attacked or sick if the number of free radicals is greater than the body's supply of antioxidants. Therefore, antioxidants are needed from outside the human body. Some of the secondary metabolites of lichen that have antioxidant bioactivity are protolichesterinic acid [29], cryptostictinolide [30], resorcinol [31], and gallic acid [32].

This study is a continuation of the exploration of secondary metabolites from lichen of the *T. flavicans* type. Previous studies have reported the results of isolation from *T. flavicans* moss, namely compounds vicanicin [33] and 3-[1'-(2",3"-dihydroxy-phenyl) -propyl] -7hydroxy-chroman-4-one have antidiabetic and antifungal activity [34]. This isolate was obtained from the extraction using acetone and purification using gravity column chromatography, eluent n-hexane and ethyl acetate. It is necessary to explore other biological activities of 3-[1'-(2",3"-dihydroxy-phenyl)-propyl]-7hydroxy-chroman-4-one, namely antioxidants, and to test their toxicity effects.

## 2. Materials and Methods

### 2.1. Materials.

Lichen *T. flavicans*, acetone, hydrochloric acid, Bouchardic Reagent, Mayer Reagent, Dragendorf Reagent, magnesium powder, and FeCl<sub>3</sub> were obtained from Merck-Germany, distilled water (IPHA, Indonesia), Whatman No. 42 filter paper (Merck-Germany), 2,2-difenil-1-pikrilhidrazil (DPPH, Merck-Germany), *Artemia Salina* Leach larvae, and 3-[1'-(2",3"-dihydroxy-phenyl) -propyl] -7hydroxy-chroman-4-one compound [34].

### 2.2. Toxicity activity test.

The toxicity test used the Meyer method and *A. salina* Leach eggs as the test material [35]. *A. salina* Leach eggs were hatched in a container filled with seawater with the help of a 14-watt lamp. The extracts and isolates tested were made into various solutions with concentrations of 1000, 500, 100, and 10 µg/mL. The test was carried out for 24 hours and counted the number of dead and living shrimp larvae. The mortality rate is calculated by comparing the number of dead larvae divided by the number of dead and living shrimp larvae.

### 2.3. Antioxidant activity test.


The antioxidant activity test was carried out based on the Blois method using DPPH as the free radical [36]. The IC<sub>50</sub> value was calculated based on the percentage of inhibition of

each concentration, namely 2.5, 10, 50, and 100 µg/mL. The percentage of inhibition (y) from each concentration (x), the points (x and y) are plotted on the coordinate plane then the line equation  $y = ax + b$  is determined by calculation using linear regression where a and b are constants, x is the concentration sample (ppm), and y is the percentage of inhibition (%). Antioxidant activity is expressed by IC<sub>50</sub>, namely the sample's concentration that can reduce 50% of DPPH radicals [37].

### 3. Results and Discussion

#### 3.1. Isolation of lichen *T. Flavicans*.

In previous studies, we have reported that the compound 3-[1'-(2'',3''-dihydroxy-phenyl)-propyl]-7hydroxy-chroman-4-one has been isolated from lichen *T. flavicans*. The structural formula can be seen in Figure 1.



**Figure 1.** Structure of the compound 3-[1'-(2'',3''-dihydroxy-phenyl)-propyl]-7hydroxy-chroman-4-one.

**Table 1.** Elucidation data of 3-[1'-(2'',3''-dihydroxy-phenyl)-propyl]-7hydroxy-chroman-4-one [34].

Spectra Data	
Molecular weight ([M <sup>+</sup> ] (m/z))	315
UV (λ <sub>max</sub> , (nm))	270
IR (ν̄, (cm <sup>-1</sup> ); functional groups)	978, C-O; 1037, C-O; 1625, C=C; 1740, C=O; 2932, Csp <sup>3</sup> -H; 3559, OH and 3619, OH
<sup>1</sup> H-NMR (δ <sub>H</sub> (ppm) (ΣH, mult, J (Hz))	7.612 (1H, d, 1.7); 7.296 (1H, d, 1.6); 7.196 (1H, d, 1.69); 6.892 (1H, d, 1.6); 4.353 (2H, s); 3.784 (3H, s); 2.966 (2H, t, 7.18); 1.59 (1H, q); 1.337 (2H, m); 0.876 (3H, t, 7.12)

#### 3.2. Toxicity test.

Based on the toxicity testing results, the extract and isolated compounds have LC<sub>50</sub> values of 9.38 and 162.18 µg/mL, which indicated that the isolated compound was toxic, where a compound was said to be toxic if the LC<sub>50</sub> value was ≤ 1000 µg/mL (Table 1). The mechanism of larval death is related to the content of flavonoid class compounds in an extract. The compound acts as stomach poisoning, so it irritates the larvae' digestive organs [38]. In addition, the taste receptors in the mouth area of the larvae are also inhibited, causing the failure of the taste stimuli in the larvae so that they are unable to recognize their food. Based on its toxic properties, it can be developed as an anticancer drug.

**Table 1.** Toxicity data of acetone extract and isolate compounds of lichen *T. Flavicans*.

Sample	K (µg/mL)	Log K	Death	Life	Accumulation		Mortality	IC <sub>50</sub> (µg/mL)
					D	L		
Control	0	0	0	10	0	40	0	-
	0	0	0	10	0	30	0	
	0	0	0	10	0	20	0	
	0	0	0	10	0	10	0	
Aseton extract of <i>T. flavicans</i>	10	1	9	21	9	30	23.08	9.38
	100	2	21	9	30	9	76.92	
	500	2.7	30	0	60	0	100	

Sample	K (µg/mL)	Log K	Death	Life	Accumulation		Mortality	IC <sub>50</sub> (µg/mL)
					D	L		
	1000	3	30	0	90	0	100	
Isolate	10	1	5	25	5	63	7.35	162.18
	100	2	3	27	8	38	17.39	
	500	2.7	22	8	30	11	73.17	
	1000	3	27	3	57	3	95.00	

### 3.3. Antioxidant test.

Antioxidant activity testing provides information about the ability of the test compound to react directly with DPPH radicals. The measurement results were obtained by observing the change in purple color to colorless due to its reduction to non-radical forms by monitoring its absorbance at the optimum wavelength of DPPH using a spectrophotometer [39,40]. The parameter as an interpretation of the test results is determined by the IC<sub>50</sub> value, which is referred to as the concentration of the test sample, causing a reduction of DPPH activity by 50%. Table 2 shows the IC<sub>50</sub> value of antioxidant activity by *T. flavicans* isolates and acetone extracts. The extract has an antioxidant activity to ward off DPPH radicals. The resulting IC<sub>50</sub> value is 54.05 indicating strong antioxidant activity. The isolated compound has an IC<sub>50</sub> value of 127.38, which is in the medium category.

**Table 2.** Antioxidant data of acetone extract and isolate compounds of lichen *T. Flavicans*.

Sample	Concentration (ppm)	IC <sub>50</sub> (ppm)
Ascorbic acid positive control	10	5.05
	25	
	50	
	100	
Aseton extract of <i>T. flavicans</i>	10	54.05
	25	
	50	
	100	
Isolate	10	127.38
	25	
	50	
	100	

## 4. Conclusion

Isolation of lichen *T. flavicans* obtained pure compounds 3-[1'-(2",3"-dihydroxy-phenyl)-propyl]-7hydroxy-chroman-4-one. The toxicity test showed that extract and isolate compounds have LC<sub>50</sub> values of 9.38 and 162.18 µg/mL, indicating that the isolated compound was toxic. The antioxidant activity showed an IC<sub>50</sub> value of 161.273 µg/mL. The antioxidant test showed that the extract and isolated compounds have antioxidant activity with IC<sub>50</sub> values of 54.05 and 127.38. The results of these studies provide scientific development of natural compounds that have the potential to act as antioxidants for diseases in the human body.

## Funding

This research was funded by DRPM Menristek/BRIN of the Republic of Indonesia, grant number 171/SP2H/AMD/LT/DRPM/2020.

## Acknowledgment

We are grateful for the laboratory facilities from the Research Center for Chemistry, Indonesian Institute of Sciences in conducting this research.

## Conflicts of Interest

We declare that this article has no conflict of interest.

## References

1. Goga, M.; Elečko, J.; Marcinčinová, M.; Ručová, D.; Bačkorová, M.; Bačkor, M. Lichen Metabolites: An Overview of Some Secondary Metabolites and Their Biological Potential. In *Co-Evolution of Secondary Metabolites*, Mérillon, J.-M., Ramawat, K.G., Eds. Springer International Publishing: Cham, 2020; [https://doi.org/10.1007/978-3-319-96397-6\\_57](https://doi.org/10.1007/978-3-319-96397-6_57).
2. Solárová, Z.; Liskova, A.; Samec, M.; Kubatka, P.; Büsselberg, D.; Solár, P. Anticancer potential of lichens' secondary metabolites. *Biomolecules* **2020**, *10*, 87, <https://doi.org/10.3390/biom10010087>.
3. Ranković, B.; Kosanić, M. Lichens as a Potential Source of Bioactive Secondary Metabolites. In *Lichen Secondary Metabolites: Bioactive Properties and Pharmaceutical Potential*, Ranković, B., Ed. Springer International Publishing: Cham, 2019; [https://doi.org/10.1007/978-3-030-16814-8\\_1](https://doi.org/10.1007/978-3-030-16814-8_1).
4. Sabarwati, S.H.; Harjuliarto, R.; Watoni, A.H.; Nurdin, M. Antibacterial activity of usnic acid from *Usnea longissima* Ach. *Pak. J. Pharm. Sci.* **2020**, *33*.
5. Stanojković, T. Investigations of Lichen Secondary Metabolites with Potential Anticancer Activity. In *Lichen Secondary Metabolites: Bioactive Properties and Pharmaceutical Potential*, Ranković, B., Ed. Springer International Publishing: Cham, 2019, [https://doi.org/10.1007/978-3-030-16814-8\\_5](https://doi.org/10.1007/978-3-030-16814-8_5).
6. Olivier-Jimenez, D.; Chollet-Krugler, M.; Rondeau, D.; Beniddir, M.A.; Ferron, S.; Delhaye, T.; Allard, P.-M.; Wolfender, J.-L.; Sipman, H.J.M.; Lücking, R. A database of high-resolution MS/MS spectra for lichen metabolites. *Scientific data* **2019**, *6*, 1-11, <https://doi.org/10.6084/m9.figshare.10265234>.
7. Maulidiyah, I.; Watu, M.; Nurdin, M. Secondary metabolites identification from Lichen *Usnea Longissima* Ach: Bioactivity test of antibacterial. *International journal of applied chemistry* **2016**, *12*, 347-357.
8. Cahyana, A.H.; Suwarso, W.P.; Nurdin, M. Isolation and structure elucidation of eumitrin A1 from lichen *Usnea blepharea* Motyka and its cytotoxic activity. *International journal of pharmtech research* **2015**, *8*, 782-789.
9. Maulidiyah, M.; Cahyana, A.H.; Suwarso, W.P. A new phenolic compound from acetone extract of lichen *usnea flexuosa* tayl. *Indonesian Journal of Chemistry* **2011**, *11*, 290-294.
10. Nayaka, S.; Haridas, B. Bioactive Secondary Metabolites from Lichens. In *Plant Metabolites: Methods, Applications and Prospects*, Sukumaran, S.T., Sugathan, S., Abdulhameed, S., Eds. Springer Singapore: Singapore, 2020; [https://doi.org/10.1007/978-981-15-5136-9\\_12](https://doi.org/10.1007/978-981-15-5136-9_12).
11. El-Garawani, I.; Emam, M.; Elkhateeb, W.; El-Seedi, H.; Khalifa, S.; Oshiba, S.; Abou-Ghanima, S.; Daba, G. In Vitro Antigenotoxic, Antihelminthic and Antioxidant Potentials Based on the Extracted Metabolites from Lichen, *Candelariella vitellina*. *Pharmaceutics* **2020**, *12*, <https://doi.org/10.3390/pharmaceutics12050477>.
12. Shameera Ahamed, T.K.; Rajan, V.K.; Sabira, K.; Muraleedharan, K. DFT and QTAIM based investigation on the structure and antioxidant behavior of lichen substances Atranorin, Evernic acid and Diffractaic acid. *Comput. Biol. Chem.* **2019**, *80*, 66-78, <https://doi.org/10.1016/j.compbiolchem.2019.03.009>.
13. Pamenta, A.F.A.; Azis, T.; Watoni, A.H.; Natsir, M.; Ansharullah; Adiba, I.F.; Dermawan, A.; Nurdin, M.; Maulidiyah. Antimalarial activity of lichen *Usnea longissima*. *AIP Conf. Proc.* **2020**, *2243*, 020017, <https://doi.org/10.1063/5.0001501>.
14. Sudarwanti, C.S.; Imran, I.; Yanti, N.A.; Musdalifah, A.; Nurdin, M.; Maulidiyah, M. Antimicrobial activity of acetone extract and usnic acid constituent of lichen *usnea longissima* (Ach.). *Int Res J Pharm* **2018**, *9*, 89–98, <https://doi.org/10.7897/2230-8407.097131>.
15. Nugraha, A.S.; Untari, L.F.; Laub, A.; Porzel, A.; Franke, K.; Wessjohann, L.A. Anthelmintic and antimicrobial activities of three new depsides and ten known depsides and phenols from Indonesian lichen: *Parmelia cetrata* Ach. *Nat. Prod. Res.* **2020**, 1-10, <https://doi.org/10.1080/14786419.2020.1761361>.
16. Mohammadi, M.; Zambare, V.; Malek, L.; Gottardo, C.; Suntres, Z.; Christopher, L. Lichenochemicals: <https://biointerfaceresearch.com/>



- extraction, purification, characterization, and application as potential anticancer agents. *Expert Opinion on Drug Discovery* **2020**, *15*, 575-601, <https://doi.org/10.1080/17460441.2020.1730325>.
17. Mallavadhani, U.V.; Tirupathamma, R.S.; Sagarika, G.; Ramakrishna, S. Isolation, Chemical Modification, and Anticancer Activity of Major Metabolites of the Lichen Parmotrema mesotropum. *Chem. Nat. Compd.* **2019**, *55*, 825-831, <https://doi.org/10.1007/s10600-019-02824-2>.
  18. Kosanić, M.; Ranković, B. Antineurodegenerative and Antidiabetic Activity of Lichens. In *Lichen Secondary Metabolites: Bioactive Properties and Pharmaceutical Potential*, Ranković, B., Ed. Springer International Publishing: Cham, 2019; [https://doi.org/10.1007/978-3-030-16814-8\\_8](https://doi.org/10.1007/978-3-030-16814-8_8).
  19. Studzińska-Sroka, E.; Dubino, A. Lichens as a source of chemical compounds with anti-inflammatory activity. *Herba Polonica* **2018**, *64*, 56-64, <https://doi.org/10.2478/hepo-2018-0005>.
  20. Sanjaya, A.; Avidlyandi, A.; Adfa, M.; Ninomiya, M.; Koketsu, M. A new depsidone from *teloschistes flavicans* and the antileukemic activity. *Journal of Oleo Science* **2020**, *69*, 1591-1595, <https://doi.org/10.5650/jos.ess20209>.
  21. Nugraha, A.S.; Pratoko, D.K.; Damayanti, Y.D.; Lestari, N.D.; Laksono, T.A.; Addy, H.S.; Untari, L.F.; Kusumawardani, B.; Wangchuk, P. Antibacterial and Anticancer Activities of Nine Lichens of Indonesian Java Island. *Journal of Biologically Active Products from Nature* **2019**, *9*, 39-46, <https://doi.org/10.1080/22311866.2019.1567383>.
  22. Suharno, S.; Chrystomo, L.Y.; Sujarta, P.; Tanjung, R.H. Rapid assessment of lichen diversity in Baliem Valley, Jayawijaya, Papua, Indonesia. *Biodiversitas Journal of Biological Diversity* **2020**, *21*, <https://doi.org/10.13057/biodiv/d210610>.
  23. Nayaka, S.; Mishra, G.K.; Upreti, D.K.; Kondratyuk, S.Y. Species and Chemical Diversity in Lichen Family Teloschistaceae, and their Bioprospecting Potential: A Review in Indian Context. *Cryptogam Biodiversity and Assessment* **2018**, *3*, 08-22, <https://doi.org/10.21756/cab.v3i2.16254>.
  24. Nandi, A.; Yan, L.-J.; Jana, C.K.; Das, N. Role of Catalase in Oxidative Stress- and Age-Associated Degenerative Diseases. *Oxid. Med. Cell. Longev.* **2019**, *2019*, 9613090, <https://doi.org/10.1155/2019/9613090>.
  25. García-Sánchez, A.; Miranda-Díaz, A.G.; Cardona-Muñoz, E.G. The Role of Oxidative Stress in Physiopathology and Pharmacological Treatment with Pro- and Antioxidant Properties in Chronic Diseases. *Oxid. Med. Cell. Longev.* **2020**, *2020*, 2082145, <https://doi.org/10.1155/2020/2082145>.
  26. Abbasi, A.M.; Shah, M.H. Assessment of phenolic contents, essential/toxic metals and antioxidant capacity of fruits of viburnum foetens decne. *Biointerface Research in Applied Chemistry* **2018**, *8*, 3178-3186.
  27. Asghari, S.; Movafeghi, A.; Lisar, S.Y.S.; Barar, J.; Omidi, Y. Effects of phenanthrene on growth parameters and antioxidant systems in the green microalga *Chlorella vulgaris*. *Biointerface Research in Applied Chemistry* **2018**, *8*, 3405-3411.
  28. Bahramian, G.; Golestan, L.; Khosravi-Darani, K. Antimicrobial and antioxidant effect of nanoliposomes containing zataria multiflora boiss essential oil on the rainbow trout fillets during refrigeration. *Biointerface Research in Applied Chemistry* **2018**, *8*, 3505-3513.
  29. Brisdelli, F.; Perilli, M.; Sellitri, D.; Piovano, M.; Garbarino, J.A.; Nicoletti, M.; Bozzi, A.; Amicosante, G.; Celenza, G. Cytotoxic Activity and Antioxidant Capacity of Purified Lichen Metabolites: An In Vitro Study. *Phytother. Res.* **2013**, *27*, 431-437, <https://doi.org/10.1002/ptr.4739>.
  30. Papadopoulou, P.; Tzakou, O.; Vagias, C.; Kefalas, P.; Roussis, V.  $\beta$ -Orcinol Metabolites from the Lichen Hypotrachyna revoluta. *Molecules* **2007**, *12*, <https://doi.org/10.3390/12050997>.
  31. Bhattarai, H.D.; Paudel, B.; Hong, S.G.; Lee, H.K.; Yim, J.H. Thin layer chromatography analysis of antioxidant constituents of lichens from Antarctica. *J. Nat. Med.* **2008**, *62*, 481, <https://doi.org/10.1007/s11418-008-0257-9>.
  32. Sisodia, R.; Geol, M.; Verma, S.; Rani, A.; Dureja, P. Antibacterial and antioxidant activity of lichen species Ramalina roesleri. *Nat. Prod. Res.* **2013**, *27*, 2235-2239, <https://doi.org/10.1080/14786419.2013.811410>.
  33. Maulidiyah, M.; Darmawan, A.; Hasan, A.; Wibowo, D.; Salim, L.O.A.; Ansharullah, A.; Mustapa, F.; Nurdin, I.F.A.; Nurdin, M. Isolation, structure elucidation, and antidiabetic test of vicanicin compound from lichen *Teloschistes flavicans*. *Journal of Applied Pharmaceutical Science* **2020**, *10*, 001-009, <https://doi.org/10.7324/JAPS.2020.10111>.
  34. Maulidiyah, M.; Hasan, A.; Irna, W.O.; Adiba Nurdin, I.F.; Kusmalawati, T.; Imran, I.; Watoni, A.H.; Azis, T.; Darmawan, A. Antifungal potential against *Aspergillus flavus*: secondary metabolite compound from unique organism of lichen *Teloschistes flavicans*. *Int Res J Pharm* **2018**, *9*, 30-5, <https://doi.org/10.7897/2230-8407.098160>.
  35. Meyer, B.N.; Ferrigni, N.R.; Putnam, J.E.; Jacobsen, L.B.; Nichols, D.E.j.; McLaughlin, J.L. Brine shrimp:

- a convenient general bioassay for active plant constituents. *Planta Med.* **1982**, *45*, 31-34, <https://doi.org/10.1055/s-2007-971236>.
36. Akar, Z.; Küçük, M.; Doğan, H. A new colorimetric DPPH• scavenging activity method with no need for a spectrophotometer applied on synthetic and natural antioxidants and medicinal herbs. *Journal of enzyme inhibition and medicinal chemistry* **2017**, *32*, 640-647, <https://doi.org/10.1080/14756366.2017.1284068>.
37. Villaño, D.; Fernández-Pachón, M.S.; Moyá, M.L.; Troncoso, A.M.; García-Parrilla, M.C. Radical scavenging ability of polyphenolic compounds towards DPPH free radical. *Talanta* **2007**, *71*, 230-235, <https://doi.org/10.1016/j.talanta.2006.03.050>.
38. Usman, U.; Soekamto, N.H.; Natsir, M.; Maulidiyah, M. Isolation of natural derivated steroidal from bark stem of *Melochia umbellata* (Houtt) Stapf var. *Degrabrata* K. *Asian J Chem* **2020**, *32*, 765–70, <https://doi.org/10.14233/ajchem.2020.22414>.
39. Yulizar, Y.; Kusriani, E.; Apriandanu, D.O.B.; Nurdini, N. Datura metel L. Leaves extract mediated CeO<sub>2</sub> nanoparticles: Synthesis, characterizations, and degradation activity of DPPH radical. *Surfaces and Interfaces* **2020**, *19*, 100437, <https://doi.org/10.1016/j.surfin.2020.100437>.
40. Zayadi, R.A.; Abu Bakar, F. Comparative study on stability, antioxidant and catalytic activities of bio-stabilized colloidal gold nanoparticles using microalgae and cyanobacteria. *Journal of Environmental Chemical Engineering* **2020**, *8*, 103843, <https://doi.org/10.1016/j.jece.2020.103843>.