

Wound healing potency of *Terminalia catappa* in mice (*Mus musculus*)

by Rudy Nugroho

Submission date: 28-Jan-2020 08:59PM (UTC+0700)

Submission ID: 1247657713

File name: J1.pdf (387.3K)

Word count: 4110

Character count: 21751



Wound healing potency of *Terminalia catappa* in mice (*Mus musculus*)

Rudy Agung Nugroho^{1*}, Retno Aryani¹, Hetty Manurung², Rudianto¹,
Dinda Prameswari¹

¹ Laboratorium Fisiologi, Perkembangan dan Molekuler Hewan, Jurusan Biologi Fakultas Matematika dan Ilmu

Pengetahuan Alam Universitas Mulawarman, Samarinda, Kalimantan Timur, INDONESIA

² Laboratorium Fisiologi Tumbuhan, Jurusan Biologi Fakultas Matematika dan Ilmu Pengetahuan Alam Universitas Mulawarman, Samarinda, Kalimantan Timur, INDONESIA

*Corresponding author: rudyagung.nugroho@fmipa.unmul.ac.id

Abstract

Leaves of *Terminalia catappa* and its phytoconstituents are well known as a natural promising medicine especially for wound healing agents. The purpose of the current research was to evaluate the wound healing potency of ethanolic and water extract of *T. catappa* either green or brown leaves in mice (*Mus musculus*) model. A group of mice were used and randomly divided into control, vaseline, povidone iodine groups and treatment groups consisting ethanolic extract either green (GLE) or brown leaves (BLE), and water extract green (GLW) and brown (BLW) leaves of *T. catappa* as ointment. All mice were prepared for excision and treated with simple ointment once a day until the wound was enclosure. The percentage of wound enclosure was recorded every three days, while total DNA and hydroxyproline content were determined at day 12. Qualitative phytochemical assay was performed to determine the phytoconstituent in the leaves extract. The results found that mice treated with GLE and GLW had better wound healing at the day 12, confirming that the wound healing activity was found to be better than control and Povidone-iodine group. Furthermore, hydroxyproline content were also found to be higher in all treated groups than control group. Present finding suggested that *T. catappa* leaves extract which contains important phytochemicals shows a good wound healing activity and can be used as alternative medicine for wound healing.

Keywords: mice, phytochemicals, *Terminalia catappa*, wound healing

Nugroho RA, Aryani R, Manurung H, Rudianto, Prameswari D (2019) Wound healing potency of *Terminalia catappa* in mice (*Mus musculus*). Eurasia J Biosci 13: 2337-2342.

© 2019 Nugroho et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

The Indian almond tree or known as *Terminalia catappa* is a native to Southeast Asia and widely distributed in both tropical and sub-tropical regions. The *T. catappa* is an attractive, long-lived tree well suited to ornamental and amenity plantings. The leaves of *T. catappa* are useful as a folk medicine such as antidiabetic, analgesic, antiulcer, hyper glycaemia, and liver-related diseases (Anand et al. 2015, Behl and Kotwani 2017, Pinheiro Silva et al. 2015). Previously reported also obtained that the leaf of *T. catappa* has an antibacterial properties (Allyn et al. 2018, Nugroho et al. 2017) and it contains benefit phytochemicals such as saponin, triterpenoid, quinone, phenolic, tannin, and flavonoid (Nugroho et al. 2016), that may be useful for wound healing.

Wound healing is a natural repairing process of the damage tissue that consists of four biochemical events: blood coagulation, inflammation which is immediately begins after injury, cell proliferation, lesion contraction and remodeling (Ghuman et al. 2019, Hermes et al. 2013). In the proliferation event, a forming of granulation

tissue is built by fibroblast and the angiogenesis process (Dwivedi et al. 2017). According to Lodhi and Singhai (2013), a successful of wound healing process depends on the several factors such as blood cells and parenchymal cells component in a time frame, extracellular matrix molecules, soluble and biochemical mediators.

Further, determining the DNA and hydroxyproline content at wound healing tissue is a well-established procedure for monitoring normal cell proliferation (Nordin et al. 2018). The enhance in the DNA and hydroxyproline content is a result of increased cell division (Ahmad et al. 2017). Moreover, the increase of DNA content can be stimulated by Platelet-derived growth factor (PDGF) which also triggered chemotaxis of fibroblasts and production of collagen, glycosaminoglycan, and collagenase (Lynch et al. 1987). Meanwhile, previous study also revealed that there was a high content of hydroxyproline due to the

Received: June 2019

Accepted: November 2019

Printed: December 2019

effects of *Rhus coriaria* extract on Sprague Dawley rats excision wound model (Amin 2018).

Though some wound model studies have been done by using several medicinal plant extracts, study of *T. catappa* which related to wound healing activity in mice model is limited. Thus, present project was aimed to determine the wound healing properties of *T. catappa* leaves by measuring the percentage of wound enclosure, DNA and hydroxyproline content. Some preliminary qualitatively phytochemical test in the crude extract of *T. catappa* leaves was also evaluated to detect presence possible phytochemical such as: flavonoid, quinon, triterpenoid, steroid, saponin, tannin, phenolic, and alkaloid.

34

MATERIALS AND METHODS

Plant Materials and Ethanol Extract

Preparation

The *T. catappa* green and brown leaves were collected from around Mulawarman University, Samarinda, East Kalimantan, Indonesia. To eliminate extraneous matter, the leaves were cleaned with distilled water and immediately dried in an oven at 40°C for 12h. The leaves were later cut and ground to make them powdery using a mill and the powder extracted using ethanol 95% for 2 days (100 g L⁻¹). The obtained extracts were filtered and put in the dark bottle and keep in 4°C until used.

Preliminary Test for Phytochemicals

To detect the presence for flavonoid in either green or brown leaves, a 2 mL leaves extract was added with concentrated hydrochloric acid and magnesium ribbon. The pink-red color indicated the presence of flavonoids. Meanwhile, quinon test was performed by adding 1 mL of leaves extract with 1 mL of concentrated sulfuric acid. A red color indicated the presence of quinon. The presence of triterpenoid and steroid in the *T. catappa* leaves extract can be detected by Liebermann-Burchard methods. Respectively, a 2 mL of leaves extract was added with 1 mL chloroform, a few drops of acetic anhydride and concentrated sulphuric acid. The presence of triterpenoid was shown by a red or brown color, while steroid by a blue or green color. The occurrence of saponin in the leaves extract can be qualitatively monitored by mixing 1 mL of extract and 5 mL distilled water. The mixture was shaken vigorously to develop the froth. The occurrence of stable froth for 15 minutes indicated the presence of saponin. The presence of tannin and phenolic was determined by adding leaves extract with 2 mL of 2% solution of FeCl₃. A blue-green or black color results revealed the presence of phenols and tannins. To screen possible alkaloid content in the leaves extract, Dragendorff Test was done by mixing 2 mL of the leaves extract with 1 mL of Dragendorff reagent. The orange or orange-reddish-

brown precipitate stated a positive result for alkaloid presence.

Acclimated Animals

30

In total of thirty-five mice (20–25g, male, 3 months old) were purchased from the local animal husbandry unit in Samarinda, Indonesia. Mice were reared and acclimated in standard housing conditions at room temperature (±25°C) with a light/dark cycle of 12/12h for 3 days. Feed and water were given ad libitum methods. After acclimatization, mice were randomly divided into seven groups, namely Control group without treatment as negative control, vaseline group as placebo group which treated with vaseline (base ointment), Povidone iodine group as positive control, treatment groups consisting base ointment with 60% of either ethanol or water extract of green leaves of *T. catappa*, and another treatment groups with 60% of either ethanol or water extract of brown leaves of *T. catappa*. Each group consisted of five mice as replication.

Ointment Preparation

The ointment was made by adding vaseline with 60% of either green or brown leaves extract of *T. catappa*. The mixture was stirred using spatula until homogeneous. The ointment was stored in the freezer and ready to be used in wound healing treatment.

Treatment Procedure

Before simple ointment treatment in each group, the hair around the back was shaved and the skin was smeared with alcohol. All mice were then anesthetized using 2% ketamine, followed by 1 cm incision with ± 1 mm depth in the back using a sterile scalpel to reach the hypodermic layer. The ointment was done by applying it to the wound in mice by using a spatula every day in the morning, from day 1 until the wound was completely closed. Each mouse was observed for wound enclosure every three days until day 15 using a callipers and documentation was carried out on the wound using a digital camera. During the trial, mouse was reared in standard housing conditions at room temperature (±25°C) with a light/dark cycle of 12/12h. Feed and water were given ad libitum methods. All procedures followed NC3Rs Animal research reporting of in vivo experiments (ARRIVE) animal ethical guidelines.

Total DNA content

Total DNA content was measured by using a method that previously used by Fikru et al. (2012). Briefly, the ointment was topically applied on the excision wound area of the mice once daily and tissue samples were taken on the 15th day. DNA was isolated with a commercial extraction kit (Genomic DNA Extraction Kit D1700; Solarbio, Beijing). Each total amount of DNA from each tissue sample was calculated in triplicates performed with Qubit 2.0 fluorometer (Invitrogen, USA).

Table 1. Qualitative test for possible phytochemical in the *T. catappa* leaves extract

Phytochemicals	Green leaves	Brown leaves
Flavonoid	+	+
Quinon	+	+
Triterpenoid	+	+
Steroid	-	-
Saponin	+	+
Tannin	+	+
Phenolic	+	+
Alkaloid	+	-

Noted: (+) Present; (-) Absent

Determination of Hydroxyproline Content

To calculate the hydroxyproline content, on the day 12 of the post excision wound procedure, a piece of skin from the healed wound area was collected. Respectively, tissues from healed wound were dried in oven at 60-70°C until weight was constant. The constant weight of healed wound was then hydrolysed using 6 N HCl at 130°C for 4 h in tight sealed tube. The resulting hydrolyte was neutralized to pH 7.0 followed by adding Chloramine T oxidation for 20 min. The reaction was terminated by addition of 0.4M per chloric acid. To develop the colour Ehrlich reagent was added at 60°C and calculated at 557 nm using UV/Vis spectrophotometer (Jenway 6305, UK).

Statistical Analysis

All values are presented as mean $\pm$ standard error mean (S.E.M.). Analysis of variance (ANOVA) and Duncan multiple range test (DMRT) as post hoc were performed to find a significant different among the groups and A value of $P < 0.05$ was considered significant. All statistical analysis was done by using SPSS for windows 20 (SPSS, Inc., USA).

RESULTS AND DISCUSSION

Present study found that both green and brown ethanolic extract leaves of *T. catappa* contained phytochemicals such as flavonoid, quinon, triterpenoid, steroid, saponin, tannin, phenolic, and alkaloid (Table 1). Either green or brown ethanolic extract *T. catappa* leaves did not contain steroid. This finding is consistent with previous study stating that no steroid has been found in both leaves (Nugroho et al. 2019). However, ethanolic extract of brown leaves of *T. catappa* showed no alkaloid. This difference might be due to the fact that water/moisture content, terrain, geographical location, seasonal variation, soil type influence phytochemicals in plants.

38

Table 2. The percentage of wound healing enclosure of mice after ointment with *Terminalia catappa* leaves

Day	Control	Vaseline	Povidone Iodine	GLE	GLW	BLE	BLW
0	0.00 $\pm$ 0.00 ^a	0.00 $\pm$ 0.00 ^a	0.00 $\pm$ 0.00 ^a	0.00 $\pm$ 0.00 ^a	0.00 $\pm$ 0.00 ^a	0.00 $\pm$ 0.00 ^a	0.00 $\pm$ 0.00 ^a
3	18.66 $\pm$ 0.66 ^a	17.00 $\pm$ 1.00 ^a	29.33 $\pm$ 2.60 ^b	12.00 $\pm$ 1.30 ^a	5.66 $\pm$ 1.45 ^c	9.33 $\pm$ 2.90 ^a	2.66 $\pm$ 0.33 ^c
6	24.00 $\pm$ 2.51 ^a	34.33 $\pm$ 2.90 ^b	40.66 $\pm$ 5.23 ^c	40.00 $\pm$ 4.16 ^c	48.00 $\pm$ 1.15 ^c	53.33 $\pm$ 5.45 ^c	13.33 $\pm$ 0.33 ^a
9	47.00 $\pm$ 1.70 ^a	74.33 $\pm$ 2.90 ^b	50.00 $\pm$ 8.66 ^a	52.66 $\pm$ 1.70 ^a	70.33 $\pm$ 1.20 ^b	71.33 $\pm$ 1.85 ^b	52.00 $\pm$ 1.15 ^a
12	94.66 $\pm$ 5.33 ^a	84.66 $\pm$ 1.76 ^{ab}	95.22 $\pm$ 4.66 ^a	100.00 $\pm$ 0.00 ^c	100.00 $\pm$ 0.00 ^c	93.33 $\pm$ 6.66 ^a	93.00 $\pm$ 1.7 ^a
15	97.00 $\pm$ 3.00 ^a	96.33 $\pm$ 3.66 ^a	100.00 $\pm$ 0.00 ^c	100.00 $\pm$ 0.00 ^b	100.00 $\pm$ 0.00 ^b	100.00 $\pm$ 0.00 ^b	100.00 $\pm$ 0.00 ^b

Note: Different letter indexes superscripts (a, b, c, d) indicate significantly different means for different treatments at $P < 0.05$. Different numerical indexes subscripts (1-6) indicate significantly different means at different times at $P < 0.05$. GLE = Green leaves ethanolic extract, GLW = Green leaves water extract, BLE = Brown leaves ethanolic extract, BLW = Brown leaves water extract

27

The wound healing is natural process to repair injury or damage tissue. Current result revealed that the percentage of wound enclosure in mice, treated topically with GLE was from 12% on day 3 to 100% on day 12, while GLW was 5.66% to 100% from day 3 to day 12, respectively. Meanwhile, mice in povidone iodine, BLE and BLW group showed completely wound enclosure at the day 15. On the other hand, the complete wound enclosure of mice in the control and vaseline group was found on the day 15 (Table 2 and Fig. 1). In agreement with these results, Sharath et al. (2010) revealed that alcoholic extract of *Bacopa monniera* was found to increase wound healing enclosure by enhancing tensile strength, wound epithelization, and the formation of connective tissue. Another study also demonstrated that, *Ceylon cinnamon* is an effective plant to stimulate the enclosure of wounds in mice (Farahpour and Habibi 2012).

Several and various phytochemicals from plants extract have been claimed as a natural phytoconstituent that responsible in wound healing. The wound healing process may involve in enhancing antioxidant production, especially in the wound area (Habibpour et al. 2003) which related to the phytochemical content from *T. catappa* extract. The results current findings confirmed that *T. catappa* leaves extract promote the healing process due to phytochemical contents.

The phytochemicals from plant such as tannin and flavonoid are responsible in wound healing enclosure that act as astringent and increased rate of epithelization (Tsuchiya et al. 1996). Further, flavonoids are stated to be a health-promoting substances which anti-inflammatory properties, having a crucial role in wound healing (Gomathi et al. 2003, Lodhi et al. 2016). Meanwhile, (Kim et al. 2011) revealed that saponin from the ginseng is also beneficial in wound healing process. Saponin has some beneficial such as hemolytic, antibacterial, anti-viral, anti-oxidative functions, anti-inflammatory activity which decrease edema and skin inflammation (Just et al. 1998, Kim et al. 2011, Navarro et al. 2001). In fact, saponin helps in accelerating neovascularization in burn wound area in mice skin. Further, saponin also relates to vascular endothelial growth factor and interleukin (IL)-1 β , both of them are a group of inflammatory cytokines that trigger the accumulation of macrophages at skin wound area, boosting wound healing (Kimura et al. 2006). In addition, phenolic may also involve in reducing inflammatory and




Fig. 2. Comparison wound healing in mice between day 0 and day 12. Note: GLE = Green leaves ethanolic extract, GLW = Green leaves water extract, BLE = Brown leaves ethanolic extract, BLW = Brown leaves water extract

Table 3. Total DNA and hydroxyproline content of new wound tissue of mice after topical application of *Terminalia catappa* leaves

Day	Control	Vaseline	Povidone iodine	GLE	GLW	BLE	BLW
Total DNA ($\mu\text{g mL}^{-1}$)	0.253 $\pm$ 0.004 ^a	0.261 $\pm$ 0.002 ^a	0.349 $\pm$ 0.01 ^b	0.849 $\pm$ 0.17 ^c	0.852 $\pm$ 0.12 ^c	0.577 $\pm$ 0.05 ^d	0.426 $\pm$ 0.07 nd
Hydroxyproline content (mg g ⁻¹ tissue)	105.10 $\pm$ 1.02	133.60 $\pm$ 1.04	199.67 $\pm$ 1.12	176.45 $\pm$ 1.13	148.97 $\pm$ 1.14	198.22 $\pm$ 1.11	188.60 $\pm$ 1.15

Note: Different letter indexes (a, b, c, d) indicate significantly different means for different treatments at $P < 0.05$. GLE = Green leaves ethanolic extract, GLW = Green leaves water extract, BLE = Brown leaves ethanolic extract, BLW = Brown leaves water extract

increase in healing activity of cutaneous wound area in mice (Song et al. 2017).

Total DNA and Hydroxyproline content are parameters that generally used in wound healing study. Present research found that total DNA from new wound tissue of mice after topical application of ethanolic (0.849 $\mu\text{g mL}^{-1}$) and water (0.852 $\mu\text{g mL}^{-1}$) extract *Terminalia catappa* green leaves were significantly higher than other groups (Table 3). Buffoni et al. (1993) stated that the increase of DNA content is shown during the skin healing, which is followed by protein content, reaching a maximum after several days of healing periods.

Meanwhile, the hydroxyproline contents in mice treated with povidone iodine, ethanolic or water extract of *T. catappa* brown or green leaves were found to be significantly ($p < 0.05$) higher than the control group. The increase in hydroxyproline content was due to a process of collagen synthesis or active proliferation of fibroblast cells that responsible for synthesizing collagens.

In conclusion, present finding stated that both ethanolic and water extract of *T. catappa* either green or brown have important phytoconstituent that useful for wound healing in mice. The ethanolic and water extract of green leaves has higher potency as wound healing in term of wound enclosure, total DNA and hydroxyproline content. Further research need to be conducted to determine the protein content in newly wound tissue and molecular aspects such as interleukin and responsible gene for wound healing.

ACKNOWLEDGEMENT

All authors thanking to IsDB that support this study through funding contract number: 136/UN17.11/PL/2019. Our gratitude also extend to research center of medicine and cosmetic, Mulawarman University and Faculty of Mathematics and Natural Sciences Mulawarman University for all kind helps.

REFERENCES

- Ahmad SB, Rehman MU, Ahmad SP, Mudasir S, Mir M (2017) Rheum emodi L. (Rhubarb) promotes wound healing by decreasing inflammatory markers and enhancing accumulation of biomolecules. *Annals of Phytomedicine*, 6: 114-118. <https://doi.org/10.21276/ap.2017.6.2.1>

- Allyn OQ, Kusumawati E, Nugroho RA (2018) Antimicrobial activity of *Terminalia catappa* brown leaf extracts against *Staphylococcus aureus* ATCC 25923 and *Pseudomonas aeruginosa* ATCC 27853. *F1000Research*, 7. <https://doi.org/10.12688/f1000research.15998.1>
- Amin ZA (2018) Effect of *Rhus coriaria* extract on wound healing potential in Sprague Dawley rats. *Zanco Journal of Medical Sciences*, 22: 89-95. <https://doi.org/10.15218/zjms.2018.012>
- Anand AV, Divya N, Kotti PP (2015) An updated review of *Terminalia catappa*. *Pharmacognosy Reviews*, 9: 93-98. <https://doi.org/10.4103/0973-7847.162103>
- Behl T, Kotwani A (2017) Proposed mechanisms of *Terminalia catappa* in hyperglycaemia and associated diabetic complications. *Journal of pharmacy and pharmacology*, 69: 123-134. <https://doi.org/10.1111/jphp.12676>
- Buffoni F, Banchelli G, Cambi S, Ignesti G, Pirisino R, Raimondi L, Vannelli G (1993) Skin wound healing: some biochemical parameters in Guinea-pig. *Journal of pharmacy and pharmacology*, 45: 784-790. <https://doi.org/10.1111/j.2042-7158.1993.tb05685.x>
- Dwivedi D, Dwivedi M, Malviya S, Singh V (2017) Evaluation of wound healing, anti-microbial and antioxidant potential of *Pongamia pinnata* in wistar rats. *J Tradit Complement Med.*, 7: 79-85. <https://doi.org/10.1016/j.jtcme.2015.12.002>
- Farahpour M, Habibi M (2012) Evaluation of the wound healing activity of an ethanolic extract of Ceylon cinnamon in mice. *Veterinari Medicina*, 57: 53-57.
- Fikru A, Makonnen E, Egale T, Debelia A, Abie Mekonnen G (2012) Evaluation of in vivo wound healing activity of methanol extract of *Achyranthes aspera* L. *Journal of ethnopharmacology*, 143: 469-474. <https://doi.org/10.1016/j.jep.2012.06.049>
- Ghuman S, Ncube B, Finnie J, McGaw L, Njoya EM, Coopoosamy R, Van Staden J (2019) Antioxidant, anti-inflammatory and wound healing properties of medicinal plant extracts used to treat wounds and dermatological disorders. *South African Journal of Botany*. <https://doi.org/10.1016/j.sajb.2019.07.013>
- Gomathi K, Gopinath D, Ahmed MR, Jayakumar R (2003) Quercetin incorporated collagen matrices for dermal wound healing processes in rat. *Biomaterials*, 24: 2767-2772. [https://doi.org/10.1016/S0142-9612\(03\)00059-0](https://doi.org/10.1016/S0142-9612(03)00059-0)
- Habibipour S, Oswald TM, Zhang F, Joshi P, Zhou XC, Dorsett-Martin W, Lineaweaver WC (2003) Effect of sodium diphenylhydantoin on skin wound healing in rats. *Plastic and reconstructive surgery*, 112: 1620-1627. <https://doi.org/10.1097/01.PRS.0000086773.96319.DA>
- Hermes D, Dudek DN, Maria MD, Horta LP, Lima EN, de Fatima A, Sanches AC, Modolo LV (2013) In vivo wound healing and antilucer properties of white sweet potato (*Ipomoea batatas*). *Journal of Advanced Research*, 4: 411-415. <https://doi.org/10.1016/j.jare.2012.06.001>
- Just MJ, Recio MC, Giner RM, Cuéllar MJ, Máñez S, Bilia AR, Ríos J-L (1998) Anti-inflammatory activity of unusual lupane saponins from *Bupleurum fruticosum*. *Planta medica*, 64: 404-407. <https://doi.org/10.1055/s-2006-957469>
- Kim YS, Cho I-H, Jeong M-J, Jeong S-J, Nah SY, Cho Y-S, Kim SH, Go A, Kim SE, Kang SS, Moon CJ, Kim JC, Kim SH, Bae CS (2011) Therapeutic effect of total ginseng saponin on skin wound healing. *Journal of ginseng research*, 35: 360-367. <https://doi.org/10.5142/jgr.2011.35.3.360>
- Kimura Y, Sumiyoshi M, Kawahira K, Sakanaka M (2006) Effects of ginseng saponins isolated from Red Ginseng roots on burn wound healing in mice. *British journal of Pharmacology*, 148: 860-870. <https://doi.org/10.1038/sj.bjp.0706794>
- Lodhi S, Jain A, Jain AP, Pawar RS, Singhai AK (2016) Effects of flavonoids from *Martynia annua* and *Tephrosia purpurea* on cutaneous wound healing. *Avicenna journal of phytomedicine*, 6: 578.
- Lodhi S, Singhai AK (2013) Wound healing effect of flavonoid rich fraction and luteolin isolated from *Martynia annua* Linn. on streptozotocin induced diabetic rats. *Asian Pacific Journal of Tropical Medicine*, 6: 253-259. [https://doi.org/10.1016/s1995-7645\(13\)60053-x](https://doi.org/10.1016/s1995-7645(13)60053-x)
- Lynch SE, Nixon JC, Colvin RB, Antoniades HN (1987) Role of platelet-derived growth factor in wound healing: synergistic effects with other growth factors. *Proceedings of the National Academy of Sciences*, 84: 7696-7700. <https://doi.org/10.1073/pnas.84.21.7696>
- Navarro P, Giner RM, Recio MC, Máñez S, Cerdá-Nicolás M, Ríos J-L (2001) In vivo anti-inflammatory activity of saponins from *Bupleurum rotundifolium*. *Life Sciences*, 68: 1199-1206. [https://doi.org/10.1016/S0024-3205\(00\)01019-5](https://doi.org/10.1016/S0024-3205(00)01019-5)
- Nordin A, Omar N, Sainik NQAV, Chowdhury SR, Omar E, Bin Saim A, Bt Hj Idrus R (2018) Low dose stingless bee honey increases viability of human dermal fibroblasts that could potentially promote wound healing. *Wound Medicine*, 23: 22-27. <https://doi.org/10.1016/j.wndm.2018.09.005>

- Nugroho RA, Manurung H, Nur FM, Prahastika W (2017) *Terminalia catappa* L. extract improves survival, hematological profile and resistance to *Aeromonas hydrophila* in *Betta* sp. Archives of Polish Fisheries, 25: 103-115. <https://doi.org/10.1515/aopf-2017-0010>
- Nugroho RA, Manurung H, Saraswati D, Ladyescha D, Nur FM (2016) The effects of *Terminalia catappa* leaf extract on the haematological profile of ornamental fish *Betta splendens*. Biosaintifika: Journal of Biology and Biology Education, 8: 241-248. <https://doi.org/10.15294/biosaintifika.v8i2.6519>
- Nugroho RA, Utami D, Aryani R, Nur FM, Sari Y, Manurung H (2019) In vivo wound healing activity of ethanolic extract of *Terminalia catappa* L. leaves in mice (*Mus musculus*). In: Journal of Physics: Conference Series. p 012031.
- Pinheiro Silva L, Damacena de Angelis C, Bonamin F, Kushima H, José Mininel F, Campaner dos Santos L, Karina Delella F, Luis Felisbino S, Vilegas W, Regina Machado da Rocha. Aparecido dos Santos Ramos, M, Maria Bauab, T, Toma, W, Akiko Hiruma-Lima, C (2015) *Terminalia catappa* L.: A medicinal plant from the Caribbean pharmacopeia with anti-*Helicobacter pylori* and antiulcer action in experimental rodent models. Journal of ethnopharmacology, 159: 285-295. <https://doi.org/10.1016/j.jep.2014.11.025>
- Sharath R, Harish B, Krishna V, Sathyanarayana B, Swamy HK (2010) Wound healing and protease inhibition activity of Bacoside-A, isolated from *Bacopa monnieri* wettest. Phytotherapy Research, 24: 1217-1222. <https://doi.org/10.1002/ptr.3115>
- Song Y, Zeng R, Hu L, Maffucci KG, Ren X, Qu Y (2017) In vivo wound healing and in vitro antioxidant activities of *Bletilla striata* phenolic extracts. Biomedicine and Pharmacotherapy, 93: 451-461. <https://doi.org/10.1016/j.biopha.2017.06.079>
- Tsuchiya H, Sato M, Miyazaki T, Fujiwara S, Tanigaki S, Ohyama M, Tanaka T, Iinuma M (1996) Comparative study on the antibacterial activity of phytochemical flavanones against methicillin-resistant *Staphylococcus aureus*. Journal of ethnopharmacology, 50: 27-34. [https://doi.org/10.1016/0378-8741\(96\)85514-0](https://doi.org/10.1016/0378-8741(96)85514-0)

Wound healing potency of Terminalia catappa in mice (Mus musculus)

ORIGINALITY REPORT

18%	11%	13%	12%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	ivsa.ir Internet Source	2%
2	Submitted to Federal University of Technology Student Paper	1%
3	Abraham Fikru, Eyasu Makonnen, Tadesse Eguale, Asfaw Debella, Getinet Abie Mekonnen. "Evaluation of in vivo wound healing activity of methanol extract of Achyranthes aspera L.", Journal of Ethnopharmacology, 2012 Publication	1%
4	Kim, Young-Soo, Ik-Hyun Cho, Moon-Jin Jeong, Soon-Jeong Jeong, Seung-Yeol Nah, Young-Sik Cho, Seung-Hyun Kim, A-Ra Go, Se-Eun Kim, Seong-Soo Kang, Chang-Jong Moon, Jong-Choon Kim, Sung-Ho Kim, and Chun-Sik Bae. "Therapeutic Effect of Total Ginseng Saponin on Skin Wound Healing", Journal of Ginseng Research, 2011. Publication	1%

5	Eduardo Sánchez, Catalina Rivas Morales, Sandra Castillo, Catalina Leos-Rivas et al. "Antibacterial and Antibiofilm Activity of Methanolic Plant Extracts against Nosocomial Microorganisms", Evidence-Based Complementary and Alternative Medicine, 2016 Publication	1 %
6	fr.scribd.com Internet Source	1 %
7	academicjournals.org Internet Source	1 %
8	Submitted to Columbia College of Missouri Student Paper	1 %
9	medcraveonline.com Internet Source	1 %
10	Submitted to Universiti Teknologi Malaysia Student Paper	1 %
11	Salnida Yuniarti Lumbessy. "Sem-sterile culture of Gracilaria salicornia seaweed (parental) on various media", AIP Publishing, 2019 Publication	1 %
12	pharmacologyonline.silae.it Internet Source	1 %
13	Nicoletta Polera, Mariateresa Badolato, Filomena Perri, Gabriele Carullo, Francesca	1 %

Aiello. "Quercetin and its Natural Sources in Wound Healing Management", Current Medicinal Chemistry, 2019

Publication

14

jneuroinflammation.biomedcentral.com

Internet Source

<1 %

15

www.ajol.info

Internet Source

<1 %

16

Sparg, S.G.. "Biological activities and distribution of plant saponins", Journal of Ethnopharmacology, 200410

Publication

<1 %

17

Gufran Ajmal, Gunjan Vasant Bonde, Pooja Mittal, Gayasuddin Khan, Vivek Kumar Pandey, Bharati V. Bakade, Brahmeshwar Mishra. "Biomimetic PCL-gelatin based nanofibers loaded with ciprofloxacin hydrochloride and quercetin: A potential antibacterial and anti-oxidant dressing material for accelerated healing of a full thickness wound", International Journal of Pharmaceutics, 2019

Publication

<1 %

18

Submitted to Our Lady of Fatima University

Student Paper

<1 %

19

Submitted to Program Pascasarjana Universitas Negeri Yogyakarta

Student Paper

<1 %

20

Submitted to Universitas Mulawarman

Student Paper

<1 %

21

Submitted to Bronx High School of Science

Student Paper

<1 %

22

Halima Saiah, Meriem Mokhtar, Wassila Saiah, Ahmed Aichouni, Fatima Zohra EL Kebir, Rachida Allem. " In vivo wound healing potential and HPLC-PDA-ESI-MS profiling of L. (Desf.) leaves methanol extract ", Journal of Food Biochemistry, 2018

Publication

<1 %

23

Abid Nordin, Norazzila Omar, Nur Qisya Afifah Veronica Sainik, Shiplu Roy Chowdhury et al. "Low dose stingless bee honey increases viability of human dermal fibroblasts that could potentially promote wound healing", Wound Medicine, 2018

Publication

<1 %

24

Deepak Kasote, Aijaz Ahmad, Alvaro Viljoen. "Proangiogenic Potential of Medicinal Plants in Wound Healing", Elsevier BV, 2015

Publication

<1 %

25

ediss.sub.uni-hamburg.de

Internet Source

<1 %

26

tessera.spandidos-publications.com

Internet Source

<1 %

27	Sunday Ositadinma Udegbunam, Rita Ijoema Udegbunam, Chijioke Clement Muogbo, Madubuike Umunna Anyanwu et al. "Wound healing and antibacterial properties of methanolic extract of <i>Pupalia lappacea</i> Juss in rats", BMC Complementary and Alternative Medicine, 2014 Publication	<1 %
28	Ekta Yadav, Deepika Singh, Pankajkumar Yadav, Amita Verma. "Attenuation of dermal wounds via downregulating oxidative stress and inflammatory markers by protocatechuic acid rich n -butanol fraction of <i>Trianthema portulacastrum</i> Linn. in wistar albino rats", Biomedicine & Pharmacotherapy, 2017 Publication	<1 %
29	beat.sums.ac.ir Internet Source	<1 %
30	Submitted to University of Malaya Student Paper	<1 %
31	sciencejournal.in Internet Source	<1 %
32	www.jmbfs.org Internet Source	<1 %
33	www.ijcpr.org Internet Source	<1 %

34

www.tandfonline.com

Internet Source

<1 %

35

zjms-hmu.org

Internet Source

<1 %

36

Submitted to Ibrahim Babangida University

Student Paper

<1 %

37

Majid Salehi, Maryam Niyakan, Arian Ehterami,
Saeed Haghi-Daredeh et al. "Porous
electrospun poly(ϵ -caprolactone)/gelatin
nanofibrous mat containing cinnamon for wound
healing application: in vitro and in vivo study",
Biomedical Engineering Letters, 2019

Publication

<1 %

38

Yusuf Özay, Sevda Güzel, Önder Yumrutaş,
Burçin Pehlivanoglu et al. "Wound Healing
Effect of Kaempferol in Diabetic and
Nondiabetic Rats", Journal of Surgical
Research, 2019

Publication

<1 %

39

Submitted to Universiti Sains Malaysia

Student Paper

<1 %

40

Submitted to Kenyatta University

Student Paper

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On