

Design of ionic liquid-based microwave-assisted extraction of flavonoids from *Cyclea barbata* Miers. and its lipoxxygenase inhibitory activity

by Islamudin Ahmad

Submission date: 06-Sep-2020 09:02AM (UTC-0500)

Submission ID: 1380637399

File name: 3102_pdf.pdf (1.88M)

Word count: 5026

Character count: 24962

Design of ionic liquid-based microwave-assisted extraction of flavonoids from *Cyclea barbata* Miers. and its lipoxygenase inhibitory activity

Islamudin Ahmad^{1*}, Berna Elya^{2*}, Yayuk Isneyni Ismail², Arikadia Noviani², Hadi Kuncoro¹, Maryono³, Neneng Siti Silfi Ambarwati⁴

¹Department of Pharmaceutical Sciences, Faculty of Pharmacy, Mulawarman University, Kampus UNMUL Samarinda, East Kalimantan, Indonesia.

²Department of Pharmacognosy-Phytochemistry, Faculty of Pharmacy, Universitas Indonesia, Kampus UI Depok, West Java, Indonesia.

³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar, Makassar, South Sulawesi, Indonesia.

⁴Department of Cosmetology, Faculty of Engineering, Universitas Negeri Jakarta, East Jakarta, Indonesia.

ARTICLE INFO

Received on: 03/08/2019
Accepted on: 22/12/2019
Available online: 04/04/2020

Key words:

Cyclea barbata Miers., ionic liquid-microwave assisted extraction, response surface methodology, total flavonoid content, lipoxygenase activity inhibition.

ABSTRACT

Green grass jelly (*Cyclea barbata* Miers.) is a plant from Indonesia that is believed to have anti-inflammatory activity. This study aims to find the optimum condition in grass jelly extraction using the ionic liquid-microwave assisted extraction toward total flavonoid content (TFC) and lipoxygenase activity (LIA). The experimental design was performed using the parameters variable including extraction time, liquid-solid ratio, and ionic liquids concentration to obtain the optimum condition. The optimization analysis used response surface methodology (RSM) with Box-Behnken design (17 trials) to obtain a predictive model with TFC and LIA as a response surface value. In the present study, the optimum condition was suggested by RSM analysis with parameter variables including extraction time of 17 minutes, 1-butyl-3-methylimidazolium bromide ([BMIM]Br) concentration of 1.76 mol/l, and the liquid-solid ratio of 38.21 ml/g. The equation of regression quadratic model was obtained to predict TFC and LIA as follows: TFC = $2.43A + 2.43B + 1.42C + 0.33AB - 3.20AC - 0.46BC - 4.90A^2 - 3.10B^2 - 3.10C^2 + 28.32$ with $R^2 = 0.8336$ and LIA = $0.066A + 8.22B + 0.97C + 2.47AB - 5.86AC + 1.96BC - 9.99A^2 - 13.75B^2 - 13.11C^2 + 63.53$ with $R^2 = 0.9207$, respectively.

INTRODUCTION

Green grass jelly (*Cyclea barbata* Miers.) is a species of Menispermaceae family which is widespread and utilized by people in many parts of Asia such as Indonesia, Thailand, India, and Burma. In Java and Sumatra island of Indonesia, the local community traditionally uses this plant as an anti-inflammatory (Siregar and Miladiyah, 2011). Some research has stated that the

green grass jelly leaves contain phenolic compounds (Arkarapanthu *et al.*, 2005; Sahasakul *et al.*, 2015) flavonoids (flavon saponins (Kusmardiyani *et al.*, 2014), rich in pectin (Hertanto *et al.*, 2017; Yuliarti *et al.*, 2017) and nutrients (Sahasakul *et al.*, 2015). In general, flavonoids have anti-inflammatory activities as described by Kumar and Pandey (2013).

Inflammation is the body's response to dangerous conditions such as infections and tissue injuries to maintain homeostasis (Medzhitov, 2008). Arachidonic acid metabolism influences inflammatory mechanisms through two pathways: lipoxygenase (LOX) and cyclooxygenase (COX). Both these pathways produce the biological mediators of inflammation including leukotriene, prostaglandin, and thromboxane (Akula and Odhav, 2008).

Conventional extraction methods take a long time and have low efficiency, and also uses excessive organic solvents

*Corresponding Author

Islamudin Ahmad, Department of Pharmaceutical Sciences, Faculty of Pharmacy, Mulawarman University, Kampus UNMUL Samarinda, East Kalimantan, Indonesia. E-mail: islamudinahmad@farmasi.unmul.ac.id; Berna Elya, Department of Pharmacognosy-Phytochemistry, Faculty of Pharmacy Universitas Indonesia, Kampus UI Depok, West Java, Indonesia. E-mail: berna.elya@gmail.com

As the process of extraction develops, many new extraction methods use fewer solvents and consider the efficiency of the extraction time (Azmir *et al.*, 2013; Tang *et al.*, 2012; Wang *et al.*, 2011). Recently, alternative solvents have been found that can replace organic solvents that are low-volatile ionic liquids, capable of dissolving specific compounds, heat-stable, and non-flammable (Chemat *et al.*, 2012). An ionic liquid as a green solvent can be combined with non-conventional extraction methods, mainly using microwave-assisted extraction methods (IL-MAE) (Chemat and Vian, 2014). Some studies have reported that the IL-MAE method successfully extracts secondary metabolite compounds, mainly including flavonoid group (Ahmad *et al.*, 2017b; Tang *et al.*, 2013; Wei *et al.*, 2012; Zhang *et al.*, 2014; 2015).

However, the studies related to optimization and application of green extraction using the IL-MAE method and research on inhibition of the lipoxygenase activity from green grass jelly leaves have not been reported. Therefore, this study contributes to the knowledge of the extraction engineering of secondary metabolite and biological properties of this plant. This study aimed to obtain an optimal IL-MAE method to the extraction of flavonoids from green grass jelly leaves and investigate its lipoxygenase inhibitory activity.

MATERIALS AND METHODS

Plant materials

Green grass jelly leaves were obtained from Bogor, West Java, Indonesia and was identified at Laboratory of Biosystematics, Indonesian Institute of Science (Lembaga Ilmu Pengetahuan Indonesia) Cibinong, West Java, Indonesia. The sample specimen (DC-001984-F) was collected at the Laboratory of Pharmacognosy-Phytochemistry, Faculty of Pharmacy, Universitas Indonesia, Depok, West Java, Indonesia.

Chemicals and equipment

Chemicals and equipment used in the study, includes Lipoxygenase kit and linoleic acid, were purchased from Sigma Aldrich. Boric acid, sodium acetate, potassium chloride, sodium hydroxide, and aluminum chloride were bought from Merck, Germany (via PT. Elo Karsa Utama, Indonesia). Aqua DM, ethanol for analysis, methanol for analysis, ethyl acetate for analysis, and ethyl acetate were purchased from PT SmartLab Indonesia, Indonesia. 1-butyl-3-methylimidazolium bromide ([BMIM]Br) was purchased from Shanghai Cheng Jie Chemical Co. LTD, China. Design Expert v10 licensed software (STATE-EASE), spectrophotometer UV-Vis (Jasco V-530, Japan), Modena Microwave 900 Watt (Buono-MV3002), Vortex mixer (Stuart, Germany), Analytical scales (Sartorius 7, Germany), sonicator (Krisbow, Indonesia), micropipette 10–100 µl and 100–1,000 µl (Corning), and pH meter (Eutech Instrument, French).

Ionic liquid-based microwave-assisted extraction procedure

The IL-MAE method was performed based on literature (Ahmad *et al.*, 2017a; 2017b), with a slight modification. Briefly, a 0.5 g of the dried sample of Green grass jelly leaves were extracted using [BMIM]Br solvent in the microwave (Modena 900 W) with some conditions as can be seen in Table 1. The separation of the

residue and the extract solution was conducted by filtration. The filtrate (extract) was cooled and left for 10–12 hours at the room temperature. Then, the filtrate was partitioned with liquid–liquid extraction using ethyl acetate. The ethyl acetate layers were separated and evaporated to obtain the dried extracts.

Determination of total flavonoid content

The total flavonoid content (TFC) of each extract was analyzed by aluminum chloride colorimetry method based on literature (Baba and Malik, 2015; Chang *et al.*, 2002; Do *et al.*, 2014) with a slight modification. In brief, 6 ml of sample solution or the standard solution were mixed in 1.5 ml methanol, 0.1 ml 10% aluminum chloride, 0.1 ml of 1 M sodium acetate, and 2.8 ml aqua DM (double-distilled water). The mixture was incubated for 30 minutes at the room temperature, and absorbance was measured at 434 nm optimum wavelength. Quercetin solution (3, 4, 5, 6, 7, and 8 µg/ml, respectively) was used as standards.

The yield of TFC in extract sample was examined using equation formula of standards ($Y = 0.0568X + 0.0576$ and $R^2 = 0.9992$, where Y is the yield of TFC and X is the absorbance), and the result was expressed as milligram quercetin equivalent per gram extract (mg QE/g extract).

Lipoxygenase inhibitory activity assay

The lipoxygenase inhibitory activity (LIA) assay was performed according to the study reported by Dzoyem *et al.* (2015). Briefly, a 10-µl extract and baicalein solution were added 1,690 µl of 20-M borate buffer solution, 1,000 µl of 300 µM linoleic acid substrate solution, and the mixture was incubated for 10 minutes at 25°C. After the incubation, the mixture was added to 300 µl of 1,000 unit/ml lipoxygenase solution. The mixture was incubated again for 15 minutes at 25°C. Then, the sample solution was added to 1,000 µl methanol, and the absorbance was measured at 234 nm using a spectrophotometer UV. Percentage of LIA was examined using equation formula as follows:

$$\% \text{ Inhibition} = \frac{(A - B)(C - D)}{(A - B)} \times 100\%$$

where A is the absorbance of the blank solution with an enzyme; B is the absorbance of blank solution without enzyme; C is the absorbance of the sample solution with an enzyme; and D is the absorbance of sample solution without enzyme.

Optimization of the IL-MAE method by response surface methodology

IL-MAE method optimization was conducted using response surface methodology (RSM) to estimate the interaction of

Table 1. The parameters optimization of the IL-MAE method using RSM with Box–Behnken design.

Factors	Unit	Symbol	Range level		
			–1	0	1
Liquid-solid ratio	ml/g	A	30	40	50
Extraction time	minutes	B	5	15	25
[BMIM]Br concentration	mol/l	C	1	2	3




Figure 1. TFC of each extract using IL-MAE method.

process and factors parameters (independent parameters) against the TFC and LIA (dependent parameters). Box–Behnken design (tree-factor-three level) was used in this study, with 17 trials (in triplicate) for the extraction parameters optimization (Table 1). A regression model was examined with a multilinear quadratic model using Design-Expert v10 software.

RESULTS AND DISCUSSION

Extraction process and pre-optimization


The extraction process was performed using the IL-MAE method to obtain the optimum extraction on total flavonoid content (TFC) and LIA from green grass jelly leaves. Pre-optimization conducted with some parameters variable condition including extraction time, liquid-solid ratio, and [BMIM]Br concentration to know the best condition. In this study, each parameter's variable was selected (Table 1) based on literature (Fan *et al.*, 2012; Xu *et al.*, 2012).

Total flavonoid content determination


Based on the result of total flavonoid content (TFC) determination (as can be seen in Fig. 1) demonstrated that the maximum of TFC (30.32 mg QE/g) was obtained from fifth trials. On the other hand, the lowest of TFC (13.20 mg QE/g) showed in second trials. Based on different combination of parameter variable at the experimental design, showed extraction time effect of TFC on the range from 5 to 25 minutes performed to obtain the optimal of extraction condition and the optimum of extraction time obtained for 15 minutes extraction time. Regarding liquid–solid ratio, the maximum of TFC was obtained at 40 ml/g with the range from 30 to 50 ml/g the increase of rate (after 40 ml/g) led to a decrease on the TFC. Meanwhile, [BMIM]Br concentration effect of TFC extraction on the range from 1 to 3 mol/l conducted to give the optimal concentration and the optimum concentration obtained at 2 mol/l.

Determination of lipoxigenase inhibitory activity


The preliminary test was performed to obtain the optimum conditions in the LIA assay using some parameters including the pH buffer of the boric acid solution, the linoleic acid substrate, and the lipoxigenase enzyme according to literature (Kumari *et al.*, 2011; Xu *et al.*, 2012). In this study, the



A



B



C

Figure 2. The optimum conditions of lipoxigenase activity inhibition assay. (A) The pH buffer of the boric acid solution, (B) the linoleic acid substrate, and (C) the lipoxigenase enzyme.

optimization of pH performed on 8.0, 8.5, 9.0, and 9.5 and the optimum condition was obtained at pH 8.5 (Fig. 2A). Optimization of the linoleic acid substrate solution was conducted to determine the optimum substrate concentration, when binding correctly with the enzyme. Based on the previous study, Xu *et al.* (2012) used a 300 μM substrate solution and Kumari *et al.* (2011) used a 250-μM substrate solution. The optimization of the substrate solution

Table 2. Results of lipoxygenase inhibitory activity by baicalein.

Concentration (µg/ml)	Absorbance				(a - b)	(c - d)	% Inhibition	Mean of % Inhibition ± STDEV	Regression equation		
	Blank (a)	Blank con. 47 (b)	Baicalein (c)	Baicalein control (d)							
0.100	0.8536	0.4435	0.8802	0.6069	0.4101	0.2733	33.36	32.05 ± 2.1642	y = 0.7864x + 1.8726 R² = 0.9975 r = 0.9987		
			0.8818	0.6080		0.2738	33.23				
			0.8894	0.6005		0.2889	29.55				
			0.7915	0.5541		0.2374	42.12				
0.125	0.8536	0.4435	0.7922	0.5567	0.4101	0.2355	42.56	42.47 ± 0.3172			
			0.7938	0.5589		0.2348	42.73				
			0.7076	0.4979		0.2098	49.00				
			0.7078	0.4970		0.4112	0.2108			48.73	49.14 ± 0.4974
0.150	0.8762	0.4650	0.7038	0.4969	0.2069		49.69	R² = 0.9975			
			0.7157	0.5568	0.1589		66.18			r = 0.9987	
			0.7123	0.5539	0.4699		0.1584				
			0.7134	0.5451		0.1683	64.19				
0.200	0.8982	0.4283	0.5871	0.4594		0.1278	72.81	72.07 ± 0.6836			
			0.5906	0.4588		0.4699	0.1318			71.94	80.24 ± 0.5537
			0.5927	0.4586	0.1341		71.46				
			0.5803	0.4870	0.0933		79.81				
0.250	0.8650	0.4028	0.5747	0.4863	0.4622		0.0884	80.87	80.24 ± 0.5537		
			0.5802	0.4880		0.0922	80.05				

was done on the concentration of 100, 200, 300, 400, and 500 µM. The optimum condition was 300 µM (Fig. 2B), whereas, the optimization of enzyme solution was performed using several concentrations of 400, 800, and 1,000 unit/ml, and the optimum solution was 1,000 unit/ml (Fig. 2C).

The LIA assay was conducted using baicalein as a standard (Putri *et al.*, 2017). Baicalein is a flavonoid from flavonols groups (Kusmardiyani *et al.*, 2014) that has the capability to inhibit the lipoxygenase activity due to hydroperoxy-octadecadienoate (HPOD) was not formed (Putri *et al.*, 2017). The higher the baicalein content, the less HPOD was formed so that, higher the inhibition percentage (Putri *et al.*, 2017). Table 2 demonstrates that the baicalein concentration from 0.1 to 0.25 µg/ml can inhibit from 33.36% to 80.05% with the linear regression equation of $y = 0.7864x + 1.8526$ ($R^2 = 0.9975$). In this study, the IC_{50} value of baicalein was 0.2 µg/ml which indicates that the enzyme works well. Figure 3 shows the highest of the LIA in 12th trials with an inhibitory percentage of 69.29%. Although the highest levels of LIA were not at fifth trials (where the most top of the TFC) and 46th trials also used the optimum condition variable include 40 ml/g liquid-solid ratio, 15 minutes extraction time, and 2 mol/l [BMIM]Br concentration.

Optimization of the IL-MAE method

Three factors with three levels (-1, 0, +1), including liquid-solid ratio (30, 40, and 50 ml/g), extraction time (5, 15, and 25 minutes), and [BMIM]Br concentration (1, 2, and 3 mol/l) were optimized using RSM with Box-Behnken design. Data analysis was performed to obtain the optimum condition for extraction of flavonoid using the IL-MAE method and LIA, the specific protocols of experimental design of parameters conditions are shown in




Figure 3. Lipoxygenase inhibitory activity of each extract.

Table 1. Based on the obtained results using RSM analysis with Design Expert v10 software, in Figure 4, demonstrated the three-dimension contour of the response surface for interaction between the factor and process parameters.

To further study the response surface plot for the TFC and LIA of the obtained extracts as a function were analyzed and developed based on the equation formula from RSM analysis results to determine the optimum condition, including liquid-solid ratio to extraction time ([BMIM]Br concentration = 2 mol/l), liquid-solid ratio to [BMIM]Br concentration (extraction time = 15 minutes), and extraction time to [BMIM]Br concentration (liquid-solid ratio = 40 ml/g). The different experimental designs




Figure 4. Three dimensional surface and contour plot. (A) Interaction of liquid-solid ratio and extraction time. (B) Interaction of liquid-solid ratio and [BMIM]Br concentration. (C) Interaction of extraction time and [BMIM]Br concentration.

Table 3. Experimental design of parameters condition for average TFC and LIA by RSM with Box-Behnken design.

Run/ Trials	Factor A: Liquid- solid Ratio (mL/g)	Factor B: Extraction time (minutes)	Factor C: [BMIM]Br concentration (mol/L)	Total Flavonoid Content (TFC) (mg QE/g)	LIA (%)
1	40 (0)	25 (1)	1 (-1)	26.41	43.82
2	30 (-1)	15 (0)	1 (-1)	13.19	30.89
3	50 (1)	25 (1)	2 (0)	25.67	53.37
4	30 (-1)	15 (0)	3 (1)	26.67	51.88
5	40 (0)	15 (0)	2 (0)	30.32	63.29
6	30 (-1)	5 (-1)	2 (0)	13.25	31.15
7	40 (0)	15 (0)	2 (0)	27.40	57.84
8	50 (1)	15 (0)	3 (1)	22.91	38.25
9	40 (0)	5 (-1)	3 (1)	20.61	25.58
10	50 (1)	5 (-1)	2 (0)	19.68	28.38
11	40 (0)	5 (-1)	1 (-1)	21.11	34.90
12	40 (0)	15 (0)	2 (0)	25.90	69.29
13	30 (-1)	25 (1)	2 (0)	17.93	46.24
14	40 (0)	15 (0)	2 (0)	26.72	68.14
15	40 (0)	25 (1)	3 (1)	24.07	42.34
16	50 (+1)	15 (1)	1 (-1)	22.21	40.69
17	40 (0)	15 (0)	2 (0)	25.31	59.05

of the extraction process with response (TFC and LIA) are demonstrated in Table 3. The obtained data were calculated by multilinear regression analysis based on the response surface. The final model of TFC and LIA from green grass jelly leaf (*C. barbata*) was given by: $TFC = 2.43A + 2.43B + 1.42C + 0.33AB - 3.20AC - 0.46BC - 4.90A^2 - 3.10B^2 - 0.98C^2 + 27.13$ and $LIA = 0.066A + 8.22B + 0.97C + 2.47B - 5.86AC + 1.96BC - 9.99A^2 - 13.75B^2 - 13.11C^2 + 63.53$, respectively.

Table 4 demonstrates the significant model with a *F*-value of 3.90. There is only a 4.33% chance that a “Model *F*-value” this large could occur due to noise. The value of “Prob > *F*” less than 0.0500 indicated model terms were significant.

Table 4. Test of ANOVA significance for regression coefficient by quadratic model with TFC as a response surface value.

Source	Sum of squares	df	Mean Square	<i>F</i> value	<i>p</i> -value Prob > <i>F</i>
Model	309.85	9	34.43	3.90	0.0433
A	47.19	1	47.19	5.34	0.0541
B	47.19	1	47.19	5.34	0.0541
C	16.07	1	16.07	1.82	0.2193
AB	0.43	1	0.43	0.049	0.8319
AC	40.83	1	40.83	4.62	0.0686
BC	0.85	1	0.85	0.096	0.7659
A ²	101.15	1	101.15	11.45	0.0117
B ²	40.37	1	40.37	4.57	0.0699
C ²	4.07	1	4.07	0.46	0.5188
Residual	61.83	7	8.83		
Lack of Fit	46.59	3	15.53	4.08	0.1041
Pure Error	15.24	4	3.81		
Cor Total	71.68	16			

*: A model where *A* is the liquid-solid ratio (mL/g), *B* is extraction time (minutes), and *C* is [BMIM]Br concentration.

Table 5. Test of ANOVA significance for regression coefficient by quadratic model with LIA as a response surface value.

Source	Sum of squares	df	Mean square	<i>F</i> value	<i>p</i> -value Prob > <i>F</i>
Model	2,888.09	9	320.90	9.03	0.0042
A	0.035	1	0.035	9.881E-004	0.9758
B	540.55	1	540.55	15.21	0.0059
C	7.51	1	7.51	0.21	0.6597
AB	24.50	1	24.50	0.69	0.4337
AC	137.24	1	137.24	3.86	0.0901
BC	15.37	1	15.37	0.43	0.5318
A ²	419.94	1	419.94	11.82	0.0109
B ²	796.54	1	796.54	22.42	0.0021
C ²	723.87	1	723.87	20.37	0.0028
Residual	248.74	7	35.53		
Lack of Fit	141.98	3	47.33	1.77	0.2911
Pure Error	106.75	4	26.69		
Cor Total	366.83	16			

*: A model where *A* is the liquid-solid ratio (mL/g), *B* is extraction time (minutes), and *C* is [BMIM]Br concentration.

Table 6. The credibility of the regression equations.

Index mark ^a	Average extraction of TFC	Average extraction of LIA
Std. Dev	2.97	5.96
Mean	22.90	46.18
C.V. (%)	12.98	12.91
PRESS	769.25	2438.52
R-Squared	0.8336	0.9207
Adjusted R-Squared	0.6198	0.8188
Predicted R-Squared	-1.0696	0.2226
Adequacy Precision	5.671	7.891

^aThe results were obtained using Design Expert v10.03 software.

The “Lack of Fit F -value” of 4.08 implies the “Lack of Fit” is not significant relative to the pure error. There is a 10.41% chance that a “Lack of Fit F -value” this large could occur due to noise and non-significant “Lack of Fit” is good.

In Table 5, the model F -value of 9.03 implies the model is significant. There is only a 0.42% chance that a “Model F -values” this large could occur due to noise. The value of “Prob > F ” less than 0.05 indicate model terms are significant. The significance of the value of 1.77 “Lack of Fit” with F value > 1.95 is not significant relative to the pure error. There is a 29.11% chance that a “Lack of Fit F -value” this large could occur due to noise.

Table 6 shows the credibility of regression equations, the correlation coefficient “ R -Squared” of 0.8416 (TFC) and 0.9207 (LIA) were obtained implies this model can express correlation of model more than 80% and 90%, respectively. A negative “Pred R -Squared” of TFC implied that the overall mean might be a better predictor of response than the current model, whereas the “Pred R -Squared” of LIA (0.2226) is not as close to the “Adj R -Squared” of 0.8188 as one might normally expect. “Adequacy Precision” measures the signal-to-noise ratio. The ratio of 5.671 (TFC) and 7.891 (LIA) indicates an adequate signal, where both ratios greater than four was desirable.

Based on the results of RSM analysis, the optimum condition was suggested with the parameter variable as follows: liquid–solid ratio of 38.21 ml/g, extraction time of 17 minutes, and [BMIM]Br concentration of 1.76 mol/l with TFC and LIA prediction of 26.38 ± 1.30 mg QE/g sample and $62.8558\% \pm 2.62\%$, respectively.

CONCLUSION

Based on the results, the design of the IL-MAE method to TFC extraction and LIA from Green grass jelly leaf (*C. barbata*) has attempted. The optimum condition of the IL-MAE method was obtained the parameter variables include 17 minutes extraction time, 1.76 mol/l [BMIM]Br concentration, and 38.21 ml/g liquid–solid ratio with the TFC prediction value of 29.07 mg QE/g sample.

This research was an early stage to develop the extraction method to obtain the target compound from natural products plant in an efficient, rapid, and environmental friendly. Future research should be performed for the optimization of extraction procedure in other parameter variables, pilot scale, isolation, and identification

of flavonoid content (mainly the active target compound) from this plant by using green chemistry approaches.

CONFLICT OF INTEREST

All authors declare that they have no conflict of interest.

FUNDING

This study was funded by Directorate of Research and Humanity Engagement (DRPM), Universitas Indonesia via grant “Hibah PITTA 2017.”

REFERENCES

- Ahmad I, Yanuar A, Mulia K, Mun'im A. Application of ionic liquid as a green solvent for polyphenolics content extraction of *Peperomia pellucida* (L) Kunth Herb. J Young Pharm, 2017a; 9:486–90.
- Ahmad I, Yanuar A, Mulia K, Mun'im A. Optimization of ionic liquid-based microwave-assisted extraction of polyphenolic content from *Peperomia pellucida* (L) Kunth using response surface methodology. Asian Pac J Trop Biomed, 2017b; 7:660–5.
- Akula US, Odhav B. *In vitro* 5-lipoxygenase inhibition of polyphenolic antioxidants from undomesticated plants of South Africa. J Med Plants Res, 2008; 2:207–12.
- Arkarapanthu A, Chavasit V, Sungpuag P, Phupathanaphong L. Gel extracted from *Khruea-Ma-Noi* (*Cyclea Barbata* Miers) leaves: chemical composition and gelation properties. J Sci Food Agr, 2005; 85:1741–9.
- Azmir J, Zaidul ISM, Rahman MM, Sharif KM, Mohamed A, Sahena F, Jahurul MHA, Ghafoor K, Norulaini NAN, Omar AKM. Techniques for extraction of bioactive compounds from plant materials: a review. J Food Eng, 2013; 117:426–36.
- Baba SA, Malik SA. Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of *Arisaema Jacquemontii* Blume. J Taibah Univ Sci, 2015; 9:449–54.
- Bubalo MC, Radošević K, Redovniković IR, Halambek J, Srček VG. A brief overview of the potential environmental hazards of ionic liquids. Ecotoxicol Environ Saf, 2014; 99:1–12.
- Carda-broch S, Berthod A, Ruiz-Angel MJ. Ionic liquids in separation techniques. J Chromatogr A, 2008; 1184:6–18.
- Chang C-C, Yang M-H, Wen H-M. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. J Food Drug Anal, 2002; 10:178–82.
- Chemat F, Vian MA. Green chemistry and sustainable technology: alternative solvents for natural products extraction. Springer US, Heidelberg, New York, Dordrecht, London, 2014.
- Chemat F, Vian MA, Cravotto G. Green extraction of natural products: concept and principles. Int J Mol Sci, 2012; 13:8615–27.
- Do QD, Angkawijaya AE, Tran-Nguyen PL, Huynh LH, Soetaredjo FE, Ismadji S, Ju YH. Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila Aromatica*. J Food Drug Anal, 2014; 22:296–302.
- Dzoyem JP, Lannang AM, Fouotsa H, Mbazona CD, Nkengfack AE, Sewald N, Eloff JN. Anti-inflammatory activity of benzophenone and xanthone derivatives isolated from *Garcinia* (Clusiaceae) species. Phytochem Lett, 2015; 14:153–8.
- Fan JP, Cao J, Zhang XH, Huang JZ, Kong T, Tong S, Tian ZY, Xie YL, Xu R, Zhu JH. Optimization of ionic liquid-based ultrasonic-assisted extraction of puerarin from *Radix Puerariae Lobatae* by response surface methodology. Food Chem, 2012; 135:2299–306.
- Hertanto BS, Kartikasari LR, Swastike W, Cahyadi M, Yuliani A, Nuhriawansa AMP. Physical properties of milk Cincau curd on different concentrations of green cincau leaf (*Cyclea Barbata* L. Miers). IOP Conf Series Mat Sci Eng, 2017; 193:2–6.

Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: an overview. *Sci World J*, 2013; 2013:1–16.

Kumari R, Meyyappan A, Selvamani P, Mukherjee J, Jaisankar P. Lipoygenase inhibitory activity of crude bark extracts and isolated compounds from *Commiphora Berryi*. *J Ethnopharmacol*, 2011; 138:256–9.

Kusmardiyani S, Insanu M, Al Asyhar M. Effect a glycosidic flavonol isolated from green grass jelly (*Cyclea barbata* Miers) leaves. *Proc Chem*, 2014; 13:194–7.

Medzhitov R. Origin and physiological roles of inflammation. *Nature*, 2008; 454:428–35.

Putri NL, Elya B, Puspitasari N. Antioxidant activity and lipoygenase inhibition test with total flavonoid content from *Garcinia kydia* Roxburgh leaves extract. *Pharmacogn J*, 2017; 9:280–4.

Sahasakul Y, Thiyajai P, Kriengsinyos W, Charoenkiatkul S. Nutrients and phenolic acids in Krung Ba Dan (*Cyclea barbata* Miers) leaves and aqueous extract from Ubon Ratchathani Province. *J Sci Technol Ubon Ratchathani Univ*, 2015; 17:1–9.

Siregar IM, Miladiyah I. Protective effects of *Cyclea barbata* Miers leaves against aspirin-induced gastric ulcer in Mice. *Universa Med*, 2011; 30:88–94.

Tang B, Bi W, Tian M, Row KH. Application of ionic liquid for extraction and separation of bioactive compounds from plants. *J Chromatogr B*, 2012; 904:1–21.

Tang B, Lee YJ, Lee YR, Row KH. Examination of 1-methylimidazole series ionic liquids in the extraction of flavonoids from *Chamaecyparis obtuse* leaves using a response surface methodology. *J Chromatogr B*, 2013; 933:8–14.

Wang SY, Yang L, Zu YG, Zhao CJ, Sun XW, Zhang L, Zhang ZH. Design and performance evaluation of ionic-liquids-based microwave-assisted environmentally friendly extraction technique for camptothecin and 10-hydroxycamptothecin from samara of *Camptotheca acuminata*. *Indust Eng Chem Res*, 2011; 50:13620–7.

Wei W, Fu YJ, Zu YG, Wang W, Luo M, Zhao CJ, Li CY, Zhang L, Wei ZF. Ionic liquid-based microwave-assisted extraction for the determination of flavonoid glycosides in Pigeon Pea leaves by high-performance liquid chromatography-diode array detector with pentafluorophenyl column. *J Separat Sci*, 2012; 35:2875–83.

Xu W, Chu K, Li H, Zhang Y, Zheng H, Chen R, Chen L. Ionic liquid-based microwave-assisted extraction of flavonoids from *Bauhinia championii* (Benth.) Benth. *Molecules*, 2012; 17:14323–35.

Yuliarti O, Chong SY, Goh KKT. Physicochemical properties of pectin from green jelly leaf (*Cyclea barbata* Miers). *Int J Biol Macromol*, 2017; 103:1146–54.

Zhang Q, Zhao S-H, Chen J, Zhang LW. Application of ionic liquid-based microwave-assisted extraction of flavonoids from *Scutellaria baicalensis* Georgi. *J Chromatogr B*, 2015; 1002:411–7.

Zhang Y, Liu Z, Li Y, Chi R. Optimization of ionic liquid-based microwave-assisted extraction of isoflavones from *Radix Puerariae* by response surface methodology. *Separat Purificat Technol*, 2014; 129:71–9.

How to cite this article:

Ahmad I, Elya B, Ismail YI, Noviani A, Kuncoro H, Maryono, Ambarwati NSS. Design of ionic liquid-based microwave-assisted extraction of flavonoids from *Cyclea barbata* Miers. and its lipoygenase inhibitory activity. *J Appl Pharm Sci*, 2020; 10(04):001–007.

Design of ionic liquid-based microwave-assisted extraction of flavonoids from *Cyclea barbata* Miers. and its lipoxygenase inhibitory activity

ORIGINALITY REPORT

19%

SIMILARITY INDEX

9%

INTERNET SOURCES

15%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to University of Hull

Student Paper

2%

2

hidrojenteknojileri.org

Internet Source

1%

3

Submitted to Institute of Graduate Studies, UiTM

Student Paper

1%

4

Liu, Zaizhi, Jia Jia, Fengli Chen, Fengjian Yang, Yuangang Zu, and Lei Yang. "Development of an Ionic Liquid-Based Microwave-Assisted Method for the Extraction and Determination of Taxifolin in Different Parts of *Larix gmelinii*", *Molecules*, 2014.

Publication

1%

5

Mohsen Ali, Siyavash Soltanian, Paria Akbary, Amin Gholamhosseini. " Growth performance and lysozyme activity of rainbow trout fingerlings fed with vitamin E and selenium, marjoram (spp.), and ajwain () extracts ", *Journal of*

1%

6

eijppr.com

Internet Source

1 %

7

eprints.whiterose.ac.uk

Internet Source

1 %

8

Islamudin Ahmad, Ayun Erwina Arifianti, Aditya Sindu Sakti, Fadlina Chany Saputri, Abdul Mun'im. "Simultaneous Natural Deep Eutectic Solvent-Based Ultrasonic-Assisted Extraction of Bioactive Compounds of Cinnamon Bark and Sappan Wood as a Dipeptidyl Peptidase IV Inhibitor", *Molecules*, 2020

Publication

1 %

9

Chunjian Zhao, Yukun Zhang, Chunying Li, Xin He, Lei Yang, Yujie Fu, Jingjing Zhang, Wenyan Zhao, Yuangang Zu. "Development of an Ionic Liquid-Based Ultrasonic/Microwave-Assisted Simultaneous Distillation and Extraction Method for Separation of Camptothecin, 10-Hydroxycamptothecin, Vincoside-Lactam, and Essential Oils from the Fruits of *Camptotheca acuminata* Decne", *Applied Sciences*, 2016

Publication

1 %

10

"Polysaccharides", Springer Nature, 2015

Publication

1 %

11	staff.ui.ac.id Internet Source	1 %
12	repository.uhamka.ac.id Internet Source	<1 %
13	doaj.org Internet Source	<1 %
14	www.academicjournals.org Internet Source	<1 %
15	amsdottorato.unibo.it Internet Source	<1 %
16	Islamudin Ahmad, Neneng Silfi Siti Ambarwati, Muhammad Arifuddin, Laode Rijai, Abdul Mun'im. "Pharmacognostic and Cytotoxicity Evaluation of Indonesia Native Plant of Piper acre Blume Leaves (Piperaceae)", Pharmacognosy Journal, 2017 Publication	<1 %
17	www.ijpsonline.com Internet Source	<1 %
18	www.science.gov Internet Source	<1 %
19	Quy Diem Do, Artik Elisa Angkawijaya, Phuong Lan Tran-Nguyen, Lien Huong Huynh et al. "Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant	<1 %

activity of *Limnophila aromatica*", Journal of Food and Drug Analysis, 2014

Publication

20

Submitted to Universitas Jenderal Soedirman

Student Paper

<1 %

21

Milica Milutinović, Neda Radovanović, Marija Ćorović, Slavica Šiler-Marinković et al.

"Optimisation of microwave-assisted extraction parameters for antioxidants from waste *Achillea millefolium* dust", Industrial Crops and Products, 2015

Publication

<1 %

22

Granica, Sebastian, Monika E. Czerwińska, Jakub P. Piwowski, Maria Ziaja, and Anna K. Kiss. "Chemical Composition, Antioxidative and Anti-Inflammatory Activity of Extracts Prepared from Aerial Parts of *Oenothera biennis* L. and *Oenothera paradoxa* Hudziok Obtained after Seeds Cultivation", Journal of Agricultural and Food Chemistry, 2013.

Publication

<1 %

23

repository.wima.ac.id

Internet Source

<1 %

24

fulltxt.org

Internet Source

<1 %

25

Chen, Quan Sheng, Tao Ying Zhou, Xing Yuan

<1 %

Li, Yu Bi, and Hui De Yang. "Applied Orthogonal Experiment Design for Optimization of Ultrasonic-Assisted Extraction of Polysaccharides from *Agaricus bisporus* Mycelia", Advanced Materials Research, 2014.

Publication

26

Oni Yuliarti. "Textural Characteristics of Indonesian Foods", Wiley, 2020

Publication

<1 %

27

Nur Laily Putri, Berna Elya, Nuraini Puspitasari. "Antioxidant Activity and Lipxygenase Inhibition Test with Total Flavonoid Content from Garcinia kydia Roxburgh Leaves Extract", Pharmacognosy Journal, 2017

Publication

<1 %

28

Ming-Lu Xu, Lan Wang, Jian-He Hu, Soek Ki Lee, Myeong-Hyeon Wang. "Antioxidant activities and related polyphenolic constituents of the methanol extract fractions from Broussonetia papyrifera stem bark and wood", Food Science and Biotechnology, 2010

Publication

<1 %

29

"Biomass and Green Chemistry", Springer Science and Business Media LLC, 2018

Publication

<1 %

30

Djamila Belhachat, Fatiha Aid, Lakhdar Mekimene, Messaouda Belhachat.

<1 %

"Phytochemical screening and in vitro antioxidant activity of Pistacia lentiscus berries ethanolic extract growing in Algeria", Mediterranean Journal of Nutrition and Metabolism, 2017

Publication

31

Siddig Ibrahim Abdelwahab, Faridah Qamaruz Zaman, Abdalbasit Adam Mariod, Muhammad Yaacob et al. "Chemical composition, antioxidant and antibacterial properties of the essential oils of Etlingera elatior and Cinnamomum pubescens Kochummen", Journal of the Science of Food and Agriculture, 2010

<1 %

Publication

32

Xu, Wei, Kedan Chu, Huang Li, Yuqin Zhang, Haiyin Zheng, Ruilan Chen, and Lidian Chen. "Ionic Liquid-Based Microwave-Assisted Extraction of Flavonoids from Bauhinia championii (Benth.) Benth.", Molecules, 2012.

<1 %

Publication

33

Farzad Deyhimi, Rahman Salamat-Ahangari, Massood Arabieh, Lida Parvin. "Application of response surface methodology for modelling the enzymatic assay of hydrogen peroxide by Emerson–Trinder reaction using 4-iodophenol", International Journal of Environmental Analytical Chemistry, 2006

<1 %

Publication

34

bmccomplementmedtherapies.biomedcentral.com

Internet Source

<1 %

35

Joyce P. Rodriguez, Jaemin Lee, Jun Yeon Park, Ki Sung Kang, Dae-Hyun Hahm, Sang Cheon Lee, Sanghyun Lee. "HPLC–UV analysis of sample preparation influence on flavonoid yield from *Cirsium japonicum* var. *maackii*", *Applied Biological Chemistry*, 2017

Publication

<1 %

36

R. Gordon, J. Chapman, A. Power, S. Chandra, J. Roberts, D. Cozzolino. "Comparison of Ultrasound-Assisted Extraction with Static Extraction as Pre-Processing Method Before Gas Chromatography Analysis of Cereal Lipids", *Food Analytical Methods*, 2018

Publication

<1 %

37

Green Chemistry and Sustainable Technology, 2014.

Publication

<1 %

38

springerplus.springeropen.com

Internet Source

<1 %

39

Hongfei Lu, Heqiang Lou, Hong Zheng, Ya Hu, Yan Li. "Nondestructive Evaluation of Quality Changes and the Optimum Time for Harvesting During Jujube (*Zizyphus jujuba* Mill. cv. Changhong) Fruits Development", *Food and Bioprocess Technology*, 2011

<1 %

-
- | | | |
|-------|---|----------------|
| 40 | <p>Jordy Kim Ung Ling, Yen San Chan, Jobrun Nandong. "Extraction of antioxidant compounds from the wastes of Mangifera pajang fruit: a comparative study using aqueous ethanol and deep eutectic solvent", SN Applied Sciences, 2020</p> <p>Publication</p> | <p><1 %</p> |
| <hr/> | | |
| 41 | <p>Xiao, Chun Ling, Ying Ren, and Hui Ling Ding. "Study on Extraction of Polyphenols from Black Soybeans and Antioxidant Activity", Applied Mechanics and Materials, 2014.</p> <p>Publication</p> | <p><1 %</p> |
| <hr/> | | |
| 42 | <p>Pooja Srivastava, P. V. Ajayakumar, Karuna Shanker. " Box-Behnken Design for Optimum Extraction of Biogenetic Chemicals from with an Energy Audit (Thermal × Microwave × Acoustic): A Case Study of HPTLC Determination with Additional Specificity Using On-line/Off-line Coupling with DAD/NIR/ESI-MS ", Phytochemical Analysis, 2014</p> <p>Publication</p> | <p><1 %</p> |
| <hr/> | | |
| 43 | <p>Tong Wang, Cheng-Bo Gu, Sui-Xin Wang, Ping Kou, Jiao Jiao, Yu-Jie Fu. "Simultaneous extraction, transformation and purification of psoralen from fig leaves using pH-dependent ionic liquid solvent based aqueous two-phase</p> | <p><1 %</p> |

44

journal.ugm.ac.id

Internet Source

<1 %

45

Yao, Huanhuan, Xinxuan Du, Lei Yang, Wenjie Wang, Fengjian Yang, Chunjian Zhao, Xiangdong Meng, Lin Zhang, and Yuangang Zu. "Microwave-Assisted Method for Simultaneous Extraction and Hydrolysis for Determination of Flavonol Glycosides in Ginkgo Foliage Using Brönsted Acidic Ionic-Liquid [HO₃S(CH₂)₄mim]HSO₄ Aqueous Solutions", International Journal of Molecular Sciences, 2012.

Publication

<1 %

46

Xiao, Xiao-Hua, Zhi-Quan Yuan, and Gong-Ke Li. "Separation and purification of steroidal saponins from Paris polyphylla by microwave-assisted extraction coupled with countercurrent chromatography using evaporative light scattering detection", Journal of Separation Science, 2014.

Publication

<1 %

47

Najmeh Mostafavi, Ali Ebrahimi. "The role of chlorine substituents in lichexanthones properties: the ionic and halogen bond interactions", Theoretical Chemistry Accounts,

<1 %

2018

Publication

48

Tuyen Thi Kim Nguyen, Chamroon Laosinwattana, Montinee Teerarak, Komkhae Pilasombut. "Potential antioxidant and lipid peroxidation inhibition of Phyllanthus acidus leaf extract in minced pork", Asian-Australasian Journal of Animal Sciences, 2017

Publication

<1%

49

Nusaibah Zahratunnisa, Berna Elya, Arikadia Noviani. "Inhibition of Alpha-Glucosidase and Antioxidant Test of Stem Bark Extracts of Garcinia fruticosa Lauterb", Pharmacognosy Journal, 2017

Publication

<1%

Exclude quotes Off

Exclude bibliography On

Exclude matches Off